

## Pleuropneumonia-like Organisms Isolated from Bronchopneumonia of Cattle

A bronchopneumonia of varying severity is invariably associated with a disease of cattle referred to as shipping or transit fever (1). The disease appears frequently in cattle transported over long distances by ship or rail. That it is a distinct infectious disease is suggested by the constancy of its pathological (2) and clinical (3) manifestations and the ease with which it is spread to unpre-disposed contact cattle.

Bacteriological examinations were conducted on 24 calves ranging in age from 4 to 12 mo, all of which had a severe bronchopneumonia either directly or indirectly related to shipment by rail. The principal bacterial species isolated from the lungs were *Pasteurella hemolytica*, *P. multocida*, and *Corynebacterium pyogenes*. Details of the bacteriological and pathological findings will be presented in a later report.

With a view to examining for the presence of organisms of the pleuropneumonia group, inocula from the lungs were streaked on "Bacto-PPLO agar" (+1 percent "Bacto-PPLO serum fraction") and "Bactotryptose agar" (+20 percent horse serum). On these mediums microscopic colonies were frequently demonstrated in primary cultures, employing the methylene blue-azure stain described by Dienes (4). However, in all but two instances subcultures were not successful and it was thought that perhaps the mediums were inadequate. This contention was strengthened by the demonstration in secondary broth cultures of forms that Freundt (5) has associated with loss of viability.

With an improved medium, essentially the same as that described by Edward (6), two additional isolations were made and maintained through indefinite subcultures without difficulty. On this medium the strains isolated on the PPLO and tryptose agar grew more rapidly and produced larger colonies. The four

strains successfully propagated were similar in regard to colonial morphology and all fulfilled Sabin's (7) criteria for admission to the pleuropneumonia group of organisms. Characteristic colonies are shown in Fig. 1.

That the pleuropneumonia-like organisms isolated are parasitic is suggested by the difficulty experienced in their isolation and propagation. Their recovery from bronchopneumonic lungs of cattle is important if only because of their resemblance to the organism that causes the serious epizootic of cattle, contagious pleuropneumonia.

G. R. CARTER

Department of Bacteriology,  
Ontario Veterinary College, Guelph, Canada

### References

1. W. Hepburn, *Vet. Record* 5, 201 (1925).
2. W. R. Graham, *J. Am. Vet. Med. Assoc.* 123, 198 (1953).
3. J. Anderson, *Vet. Record* 51, 65 (1939).
4. L. Dienes, personal communication.
5. E. A. Freundt, *Acta. Pathol. Microbiol. Scand.* 31, 561 (1952).
6. D. G. ff. Edward, *J. Gen. Microbiol.* 1, 238 (1947).
7. A. B. Sabin, *Bacteriol. Revs.* 5, 1 (1941).

26 April 1954.

## The Heparin-like Activity of Certain Inorganic Anions

Phosphotungstate, phosphomolybdate, and silicotungstate have been found to delay the clotting of blood, both *in vitro* and *in vivo*. When injected into rats, they also elicit the appearance in the plasma of lipemia "clearing factor" (1), an enzyme that catalyzes the lipolysis of chylomicrons and low-density lipoproteins. This clearing factor appears to be identical with that produced by heparin injection. Both anticoagulant activity and clearing-factor production occur following oral administration of silicotungstate.

The test animals were male Sprague-Dawley rats weighing about 180 g. The usual intravenous dose was 20 mg dissolved in 1 ml of 0.15M sodium chloride solution; it was injected either as the acid or as the sodium salt. The usual oral dose of silicotungstate was 200 mg of the salt in 1 ml of water administered by stomach tube in the fasting state. The heparinoid activities were apparent a few minutes after intravenous injection and 20 min after oral administration (Table 1). Lee-White clotting times were prolonged two- to fivefold. Clearing activity was measured by the decrease in optical density of a saline suspension of human chylomicrons incubated at 38°C with 1/10 rat plasma by volume during a period of 1 hr.

The anticoagulant activity was inhibited by protamine, both *in vitro* and *in vivo*. Clearing activity was inhibited by protamine and by high sodium chloride concentrations (2). During clearing, glycerol was produced. All three salts exhibited metachromatic activity with toluidine blue. In all these respects they resemble heparin.

These findings make it appear likely that the hepar-




Fig. 1. Colonies of the pleuropneumonia-like organism from bronchopneumonia of cattle ( $\times 120$ ).