

Technical Papers

Feather Papilla Stimulation by Progesterone*

C. S. Shaffner

Poultry Department,
University of Maryland, College Park

The presence of progesterone or a physiologically equivalent substance has recently been found in the blood of the fowl. Although in the hen there appears to be no equivalent of the mammalian corpus luteum, the presence of the corresponding secretion is clear (1). A new and interesting role may now be ascribed to progesterone in the fowl, the activation of the resting papilla of the feather.

Hens of comparable age, breed, and state of maintenance were defeathered in similarly selected regions of the breast and neck feather tracts. Eleven hens received, by intramuscular injection, 20 mg each of a progesterone solution (2), which is slowly absorbed and has a prolonged physiological action. Three comparable hens served as controls.

Egg production in the treated group ceased immediately after treatment, and upon examination 2 wk later, all treated birds showed full feather regeneration in the areas of the neck and breast that had been previously defeathered. All control birds failed to regenerate neck feathers, and regeneration was sporadic in the breast areas. The progesterone-treated hens, furthermore, showed many new incoming feathers in all regions of the body and proceeded to replace much of the plumage. An average of four primary flight feathers per wing were molted and replaced. It may be deduced from this occurrence that the new growth was from the progesterone-activated papilla and was not due to a stimulation of the resting papilla by the plucking of old feathers.

Progesterone administered to mature cocks caused the same kind of molt but to a less spectacular degree. This effect of progesterone in the male required no estrogen priming.

The initiation of new feather regeneration in birds has long been known to follow the administration of large dosages of thyroidally active material. Conceivably the observed action of progesterone could follow via a stimulation of the bird's own thyroid. Although this is recognized as a possibility, it is unlikely to be the case, since the time interval from the inception of treatment to the initiation of feather proliferation is similar in either treatment. Van der Meulen (3) reported that 7 to 8 days after hens were fed 2 to 20 g of desiccated thyroid powder they showed signs of a molt, which reached a maximal degree 14 to 25 days following treatment. In the present experiments, feather regeneration started 8 to 10 days following progesterone treatment, and the heaviest shedding of feathers occurred after about 2 wk. Furthermore,

when molt is caused by increasing the circulating level of thyroxin, the structure of the feather is also affected. This was not found to be the case with the progesterone-treated birds. For the present, the effect of the steroid is, therefore, considered to be a direct one upon the feather papilla. The relationship of progesterone-induced molt to reproduction will be described elsewhere.

References and Notes

- * Scientific article No. A464; contribution No. 2544 of the Maryland Agricultural Experiment Station (Department of Poultry Husbandry).
1. R. M. Fraps, C. W. Hooker, and T. R. Forbes, *Science* **109**, 493 (1949).
 2. Repositol Progesterone, Pitman-Moore Co., Indianapolis, Ind.
 3. J. B. Van der Meulen, *Proc. 7th World's Poultry Congr.* (1939), p. 109.

26 April 1954.

Failure of the Newborn Pig to Utilize Dietary Sucrose

D. E. Becker, D. E. Ullrey, S. W. Terrill,
R. A. Notzold

Department of Animal Science,
University of Illinois, Urbana

An inadequacy of sucrose in a synthetic milk diet for the baby pig was initially observed by Johnson (1). In a recent report Becker *et al.* (2) confirmed the detrimental effects of sucrose for the newborn pig and postulated that the difficulty resulted from an inability of the newborn pig to cope with the glycosidic linkage. The observations of Newton and Sampson (3) indicate that a failure in the metabolism of fructose may also be involved in an elucidation of the observed effects.

Newborn Duroc pigs that were allowed to nurse their dams for 24 hr following birth were used for this study. Three litters were distributed among four groups as the pigs became available. The pigs were group-fed *ad libitum* for a 9-day period on the floor of a shelter that had not been used for swine within recent years. The ingredients of the diet, except for the specific carbohydrate which varied with treatment, are presented in Table 1. Preparation of the milk diet and management of the pigs were described previously (2). Reagent grade sugars were employed, and the invert sugar consisted of an equimolecular mixture of D-glucose and D-fructose.

The results of the study are presented in Table 2. With glucose as the carbohydrate, performance was very satisfactory in terms of weight gain and survival. Although one pig succumbed on the glucose diet, it did exhibit a weight gain prior to death. Pigs receiving glucose occasionally voided soft feces, but they never exhibited a characteristic diarrhea.