

end of the animal is excited by the electric impulse is not understood. (iv) The swimming response appears to have a rather specific threshold of stimulation. But to divorce this response completely from all phasic activity one would have to establish the existence of a specific receptor-effector mechanism for this response that could operate independently of phasic activities. The existence of such a mechanism in the simple actinian nervous system does not seem probable. A possible answer might be that the swimming response is a combination of accelerated phasic activities.

Figure 1 shows that the general motions of *Stomphia* are the same motions found in slower phasic activity. Likewise, this sequence of movements closely resembles that of the feeding activities of *Metridium senile* as described by Batham and Pantin. Therefore, the swimming response may be all the typical phasic activities in sequence accelerated to a high degree by the presence of the starfish or the starfish substance.

Further work is in progress at these laboratories; special emphasis is being placed on the description of the neuromusculature system of this anemone.

C. S. YENTSCH

D. C. PIERCE

Department of Oceanography,  
University of Washington, Seattle

#### References and Notes

1. This work was partly supported by Office of Naval Research contract N8onr-520/III with the University of Washington (Contribution No. 190 from the Department of Oceanography.) Acknowledgment is extended to Richard H. Fleming, executive officer, for the use of the departmental facilities; to M. P. Wennekens and E. E. Collias for their technical assistance, and to L. M. Passano and D. L. Ray for their many helpful suggestions during the preparation of this report.
2. Four copies of this film are available, two at the Department of Oceanography and two at the Department of Zoology, University of Washington, Seattle.
3. H. M. Milligan, *Nature* 96, 619 (1916).
4. T. A. Stephenson, *The British Sea Anemones* (Ray Society, London, 1935), vol. 2.
5. E. J. Batham and D. F. A. Pantin, *J. Exptl. Biol.* 27, 377 (1950).

22 July 1955

### Variations of Nitrogen-15 Abundance in Naturally Occurring Substances

It has been shown that the isotopic abundances of many of the lighter elements are not constant (1). This paper is a report of some measurements on the variation of the  $N^{15}$  abundance in a number of naturally occurring substances. The absolute abundance of  $N^{15}$  in atmospheric nitrogen has been measured to be 0.365 percent (2). The  $N^{15}$  to  $N^{14}$  ratio in the  $N_2$  isolated from the samples has been compared directly with the ratio in

standard atmospheric  $N_2$ , and the results are expressed as parts per 1000 difference from the  $N^{15}$  to  $N^{14}$  ratio in the standard:

$$\frac{(N^{15}/N^{14})_x - (N^{15}/N^{14})_{std.}}{(N^{15}/N^{14})_{std.}} \times 10^3$$

Protein nitrogen was reduced quantitatively to ammonium salts by the standard Kjeldahl digestion, followed by oxidation to  $N_2$  by sodium hypobromite (3). Since this procedure alone did not yield sufficiently pure samples, the  $N_2$  was purified by repeated passage over Cu-CuO at 700°C in a carefully outgassed quartz furnace attached to a liquid nitrogen trap.

Inorganic nitrates were quantitatively reduced to ammonium salts by reduction with iron in dilute sulfuric acid (4), followed by oxidation to  $N_2$  and purification. The  $N_2$  from natural gas was isolated by passing the gas through a liquid nitrogen trap and then cycling the noncondensable fraction over hot CuO-Cu at 700°C until a constant value for the  $N^{15}$  to  $N^{14}$  ratio was obtained. In several cases, the sample was separated from a large excess of helium by trapping the  $N_2$  in charcoal at liquid nitrogen temperature and pumping off the noncondensed helium with a Toeppler pump.

The  $N_2$  from rocks and minerals was obtained by heating 100-g portions of the

Table 1. Variations in  $N^{15}$  abundance compared with standard atmospheric  $N_2$  of local origin.

Sample	Origin	Difference in $N^{15}/N^{14}$ ratio from standard (parts per 1000)
<i>Plant protein</i>		
Leaves, white clover	Local	- 6.5
Leaves, dandelion	Local	- 2.8
Leaves, red oak	Local	- 0.9
Leaves, cedar	Local	1.3
Leaves, American elm	Local	1.9
Weeds	Local	4.3
Oats	Unknown	6.2
Seaweed	Tokyo Bay, Japan	8.1
<i>Animal protein</i>		
Egg, domestic chicken	Local	5.8
Clam flesh	Atlantic Ocean	7.3
Lamb flesh	Unknown	5.0
Milk	Local	5.1
White rat, brain tissue	Local	5.4
White rat, lung tissue	Local	7.5
White rat, liver tissue	Local	4.5
White rat, blood	Local	6.2
White rat, skin and hair, thorax	Local	5.0
<i>Peat and coal</i>		
Peat	Eire	1.9
Peat	Junius, N.Y.	- 2.8
Lignite	Bowman, N.D.	- 1.2
Bituminous coal	Pittsburgh, Pa.	- 0.9
Cannel	Cannel City, Ky.	1.6
Anthracite	Gunnison, Colo.	- 1.2
Anthracite	Lehigh, Pa.	- 1.4
<i>Oil and gas wells</i>		
Ella Well	Hunton Lime Formation, Okla.	- 8.1
Plaisted No. 1	Marchand Formation, Okla.	- 3.5
Steve No. 1	Upper Bradley Formation, Okla.	- 8.2
Steve No. 1	Lower Bradley Formation, Okla.	2.9
Fletcher No. 10	Marchand Formation, Okla.	- 7.6
Bitt No. 1	Hart Formation, Okla.	- 11.5
Matheson 96-percent methane	Unknown	- 13.0
Natural gas	Washington County, Ark.	- 5.9
<i>Rocks and Minerals</i>		
Granite	Chelmsford, Mass.	- 0.2
Granite	Milford, Mass.	- 0.9
Pitchblende	Great Bear Lake, Canada	- 2.3
<i>Inorganic nitrogen</i>		
Chile nitratite	Tarapaca, Chile	- 2.6
Sal ammoniac	Paracutin, Mexico	13.0

powdered sample with CuO at 750°C for 4 hours in a carefully evacuated and outgassed Vycor flask that was connected to the vacuum system. The evolved gases were passed through a trap at liquid nitrogen temperature and treated with hot CuO-Cu. The standard atmospheric nitrogen was isolated from air by passing air repeatedly over cleaned copper turnings in a quartz tube at 700°C and then through a liquid nitrogen trap.

The mass spectrometer is of the design described by several workers for the measurement of small differences in isotope ratios (5). Mass spectrometry of nitrogen is troubled by the presence of background peaks in the instrument at mass 28 and 29. The background peaks were less than 0.1 percent the size of the signal peak at mass 28. The procedure of rapidly shifting from the standard nitrogen sample to the unknown sample minimized the effect of this background. It is imperative that N<sub>2</sub> samples be free of carbon monoxide, which gives an interfering mass spectrum. This problem was solved by passing the gas over hot CuO and through a liquid nitrogen trap. Several samples were prepared and their N<sup>15</sup> content was measured both before and after repeated treatments in this manner. There was no detectable change in the N<sup>15</sup> content.

In order to avoid isotopic fractionation in sample preparation, all operations were checked to insure quantitative yields. The estimated precision on each of the results listed in Table 1 is  $\pm 0.5$  parts per 1000.

THOMAS HOERING

Department of Chemistry, University of Arkansas, Fayetteville

#### References and Notes

- See K. Rankama, *Isotope Geology* (McGraw-Hill, New York, 1954) and E. Ingerson, *Bull. Geol. Soc. Amer.* 64, 301 (1953) for reviews of the work on this subject.
- A. O. Nier, *Phys. Rev.* 77, 789 (1950).
- D. Rittenberg, *Preparation and Measurement of Isotopic Tracers* (Edwards, Ann Arbor, Mich.), p. 31.
- R. C. Griffin, *Technical Methods of Analysis* (McGraw-Hill, New York, 1927).
- C. R. McKinney *et al.*, *Rev. Sci. Instr.* 21, 724 (1950); M. Dole, *Chem. Revs.* 51, 275 (1952).
- Monastero *et al.*, *J. Am. Pharm. Assoc. Sci. Ed.* 40, 241 (1951).
- C. W. Waller *et al.*, *J. Am. Chem. Soc.* 74, 4981 (1952).
- P. P. Minieri and A. G. Mistretta, *Antibiotics and Chemotherapy*, in press.
- Present address: Heyden Chemical Company, Garfield, N.J.
- Present address: Lederle Laboratories Division, American Cyanamid Company, Pearl River, N.Y.

24 June 1955

### Localized Electoretinograms from Isolated Poikilothermic Retinas with Macroelectrodes

What effect does the stimulation of some retinal elements have on the effect of stimulating others? Is there a real effect, or only an apparent one caused by stray light, chiefly from scattering in the dioptric media of the eye?

The interchangeable effects of stimulus area, duration, and intensity on the electroretinogram (ERG) of the intact frog eye were considered by Granit (1) to be evidence for interaction of retinal elements. Fry and Bartley (2), using the intact rabbit eye, showed that stray light could explain the apparent interaction and that, indeed, the ERG was mainly a response to stray light within the eye. Granit, Rubinstein, and Therman (3) obtained results apparently supporting interaction when they minimized stray light by using small stimuli of low intensity in an excised and opened frog eye. Recently, the stray light theory of Fry and Bartley has been revived and confirmed for the human ERG (4). This, together with the inconclusive character of the evidence for retinal interaction, raises several questions regarding the effects that stimulation of one retinal locus may have on the response from another locus.

We have recorded ERGs from isolated frog (*Rana pipiens* and *R. catesbeiana*) and terrapin (*Pseudemys elegans*) retinas. We removed the retina under subdued illumination and placed it flat on a black felt pad that had been soaked in Ringer's solution. The preparation was placed in a black box to minimize further any effects from stray light. Thread wicks from silver-silver chloride electrodes were led to any desired points on the surface of the retina, and another to the supporting pad. Two channels of alternating-current amplification led to a dual-beam cathode-ray oscilloscope and permitted simultaneous registration of potential changes that occurred between two pairs of electrodes, if desired, or between one pair of electrodes if the second beam was used for a time scale and stimulus marker.

Each stimulus was a 1-mm spot of light of adjustable intensity and duration. Two such spots can be presented at any

The countercurrent distribution system consisting of McIlvaine's phosphate-citrate buffer at pH 4.5 versus chloroform has been used in these laboratories for some time for the analytic separation of mixtures of the known tetracycline antibiotics. In a 50-tube distribution with this system, the peak tubes observed are as follows: chlortetracycline 26, tetracycline 39, and oxytetracycline 44. These values correspond to K aqueous/solvent values of 1.13, 3.90, and 8.80, respectively.

Pigments with a strong absorptive capacity in the ultraviolet, such as anhydrotetracycline (7), usually localize in the low-numbered tubes because of their excellent solubility in chloroform; thus their presence does not interfere with the ultraviolet determination of the peak tubes, which is carried out on the upper phase at a wavelength of 265 mμ after suitable dilution.

After the theoretical curves for each component have been calculated, the percentage composition of a mixture such as one containing tetracycline and chlortetracycline can be calculated within a few percent of the bioassay, via the peak heights as determined spectrophotometrically.

The method has been used not only on purified preparations but also on crude preparations, such as those obtained from butanol extractions of fermentation beers. Samples assaying 300 to 500 μg/mg have given satisfactory results in the identification of major components (not traces) when as little as 3 to 5 mg was placed in the first tube of a 50-tube apparatus containing 10 ml of each phase. If larger samples are used, as little as 1 percent chlortetracycline in tetracycline can be detected.

A change in pH from 3.5 to 5.5 seems to have little effect on the position of the peak tubes. Since the buffer can be made up in the range of pH 2.2 to 8.0 and since the solubility of the tetracyclines increases sharply at lower pH values, higher charges can be accommodated at a slightly lower pH (for example, 3.5). By the use of such higher concentrations in a larger apparatus, sufficient material can be handled to give a useful separation.

This method will be described in more detail in a forthcoming publication (8).

P. PAUL MINIERI\*

A. G. MISTRETTA†

American Cyanamid Company, Princeton, New Jersey

#### References and Notes

- P. P. Minieri *et al.*, *Antibiotics Annual* 81 (1953-54).
- W. T. Sokolski *et al.*, *Arch. Biochem. and Biophysics* 43, 238 (1953).
- D. Perlman, *Science* 118, 628 (1953).
- H. L. Bird and C. T. Pugh, *Antibiotics and Chemotherapy* 4, 750 (1954).
- N. Bohonos *et al.*, *Antibiotics Annual* 54 (1953-54).

### Differentiation and Separation of the Tetracycline Antibiotics by Countercurrent Distribution

Recently, the discovery of tetracycline (1) in *Streptomyces* elaboration products has increased interest in methods designed to differentiate among the members of this family of compounds.

In the past, various procedures (2-6) have been reported for carrying out such a differentiation; these methods are for detection only and have as yet had no extension into the field of separation.