




Fig. 3. Absence of correlation between the concentration of free ATP and contraction.

cium binding (9). It may be concluded that, in the case of muscle, calcium in the form of its ATP complex is not available to stabilize the muscle membrane. The action potentials so evoked set off the contractile process. Thus the effect of ATP applied to living muscle is unrelated to contraction per se.

GERTRUDE FALK

Department of Pharmacology,
University of Washington
School of Medicine, Seattle

References and Notes

1. F. Buchtal, A. Deutsch, G. G. Knappeis, *Acta Physiol. Scand.* 8, 271 (1944).
2. G. Falk and R. W. Gerard, *J. Cellular Comp. Physiol.* 43, 393 (1954).
3. A. Szent-Gyorgyi, *Chemistry of Muscular Contraction* (Academic Press, New York, 1947).
4. W. Hasselbach and A. Weber, *Pharmacol. Revs.* 7, 97 (1955).
5. W. Feldberg and C. Hebb, *J. Physiol. London* 107, 210 (1948).
6. F. Buchtal *et al.*, *ibid.* 106, 3P (1946).
7. E. D. Adrian and S. Gelfan, *ibid.* 78, 271 (1933).
8. S. W. Kuffler, *J. Neurophysiol.* 7, 17 (1944).
9. V. DiStefano and W. F. Neuman, *J. Biol. Chem.* 200, 759 (1953).
10. A. B. Hastings *et al.*, *ibid.* 107, 351 (1934).

27 September 1955

High Incidence Blood Group Found in Venezuelan Indians

To date no single blood group found exclusively in discrete geographic or ethnical human divisions has been described. It is only through the incidence of the well-known blood groups that the various human stocks can be characterized in a general way (1). We present here (2) data on the incidence of what appears to be a new blood system with possible anthropological implications.

In 1954, Levine, Koch, McGee, and Hill (3) mentioned a new "private" blood factor called "Diego," which was detected in the serum of a Venezuelan woman who had been sensitized through several pregnancies. They found that this serum produced agglutination of the hus-

band's red cells but did not agglutinate the red cells from 200 North American persons. Lately, with no further comments, this finding has been quoted in other publications (4). This year, serum was collected during a new pregnancy of the original patient, giving us an opportunity to study the incidence of the Diego factor in her husband's relatives and in various representative sections of the Venezuelan population.

In the study of the family (Ca. family) in which this factor was originally detected, we found eight positive cases out of 29 tested. In the general population from Caracas, we were surprised to find that several unrelated individuals carried this factor. Inquiry about the ancestry and physical features of the various positive cases and of the Ca. family revealed the probability that they all possessed ancestors from Carib Indian stock. The results of 826 tests in groups of people from various Venezuelan regions are given in Fig. 1. They were tested against anti-Diego serum by the indirect Coombs test.

It would seem that the Diego factor is not a "private" blood group, but rather that its incidence is high in Indians, especially in Carib Indians, and in people with mixed Indian ancestry. Since the Indians studied came originally from Brazil, it could be that this factor is prevalent in Brazil and other neighboring countries. The Indian element enters in a high proportion of the general Venezuelan population. This probably explains the positive tests found in the populations of Caracas and Barcelona. The cases found in the Negro population studied may also be explained by mixture with Indians.

From the genetic point of view, the study of the Ca. family and of several Indian families shows that the factor is inherited as a dominant Mendelian character with no sex linkage (examples in Fig. 2). The factor may be followed in some cases through several generations (four in one of the families). Apparently, the antibody is not of the naturally occurring type but is an immune "incomplete" one. Detailed study of the positive cases show that the antibody (anti-Diego




Fig. 1. Results of 826 tests.




Fig. 2. Examples of genetic studies.

serum), which can be called anti-Di^a, was developed by a homozygous Di^b/Di^b (Mrs. Ca., the original patient). The individuals who reacted against the anti-Diego serum were Di^a/Di^b or Di^a/Di^a, indicating that this blood system is formed by at least two allelic genes. We can postulate that eventually anti-Di^b will be found.

If in the future it can be demonstrated that the Diego factor occurs exclusively or predominantly in Indian populations, it would be wise to change the name of this antigen to a more correct one, such as "Indian factor," related to its anthropological implications.

MIGUEL LAYRISSE

TULIO ARENDS

Banco de Sangre, Caracas, Venezuela

References and Notes


1. A. E. Mourant, *The Distribution of the Human Blood Groups* (Blackwell, Oxford, 1954); W. C. Boyd, *Science* 112, 187 (1950).
2. Presented in part at the Venezuelan Academy of Medicine, Caracas, 28 July 1955.
3. P. Levine, E. A. Koch, R. T. McGee, G. H. Hill, *Am. J. Clin. Path.* 24, 292 (1954).
4. M. Van der Hart, H. Bosman, J. J. Van Loghem, Jr., *Vox Sanguinis* 4, 108 (1954); P. Levine, *V Congress International de Transfusion Sanguine, Rapports et Communications* (Paris, 1954).

26 September 1955

Spiral Male Nuclei in Ragweed Pollen

Stamens from ragweed (*Ambrosia*) growing on the premises of the Southern Illinois University were stained with iron-acetocarmine solution and observed under the microscope at 1000 diameters magnification. At the metaphase of the first meiotic division of the pollen mother cell, 18 round chromosomes were counted. Since the basic number of chromosomes in Compositae is nine, it is not surprising that the chromosome number of the ragweed is $n = 18$.

The chromosomes are regularly arranged on the equatorial plane of the first meiotic division (Figs. 1 and 2). At diakinesis, chiasmata were observed in



Figs. 1-12. Ragweed pollen.

every bivalent chromosome (Fig. 3). At the end of the first anaphase, the partition wall begins to appear between the two newly formed chromosome groups (Fig. 4); at the end of the second division, it is clearly apparent (Fig. 5).

When the tetrad is formed, each cell changes into a pollen grain with a thick cell wall (Figs. 6 and 7). After the first division of the pollen nucleus, a vegetative nucleus and a germ nucleus are produced, and the germ nucleus once more divides into two male nuclei (Figs. 8-10).

Each of the two male nuclei gradually change into the long, banded, sharp-pointed structures. One side of the banded nucleus is stained especially deeply. The band-shaped nuclei form spirals and resemble the spindle-shaped spermatozoids of a fern or a moss (Fig. 11). I have studied the cytomorphological features of many species of ferns (1), and a spermatozoid of *Alsophila martensiana* (Fig. 12) shows some resemblance to the male nucleus of ragweed (2). The male nucleus of Angiospermae corresponds to the spermatozoid of a fern or a moss, and it is interesting from the viewpoint of phylogeny that the male nuclei of an *Ambrosia* should show a spiral form.

In *Fritillaria* (3), in *Lilium* (4), and in *Monotropa* (5), the male nuclei become spiral-shaped after they enter the embryo sac, but this is the first instance in which the male nucleus of a higher plant has been found to exhibit a spiral form in the pollen.

AKIRA YUASA*

Biological Research Laboratory,
Southern Illinois University, Carbondale

References and Notes

1. A. Yuasa, *Botan. Mag. Tokyo* 46, 4 (1933).
2. —, *Studies Tokugawa Inst.* 4, 1 (1938); *Botan. Mag. Tokyo* 67, 6 (1954).
3. K. Sax, *Bull. Torrey Botan. Club* 43, 505 (1916).
4. E. J. Welsford, *Ann. Botany London* 28, 265 (1916).
5. A. Yuasa, unpublished.

* Fulbright grantee (1955) from the University of Tokyo.

3 October 1955

Genus *Haemagogus* in the United States

We have been engaged during the past 6 years in the study of the tropical American mosquitoes of the genus *Haemagogus* (Diptera, Culicidae) that are associated with the wave of sylvan yellow fever that passed through Panama during the period 1948-51 and in 1954 reached the north coast of Honduras.

In the course of field work in Middle America, we came to realize that this genus, which had been studied primarily in the tropical rain forests of South America, includes species characteristic of very different ecological situations. In southern Mexico, near Tuxtla Gutierrez, we found two species of *Haemagogus* at elevations in excess of 4000 feet, associated with a semiarid scrub-type of vegetation. This led us to believe that there were members of the genus that might inhabit similar situations at lower elevations to the north of the Tropic of Cancer. We have been interested in determining the northern limits of the distribution of these mosquitoes because of their implication in the transmission of sylvan yellow fever.

After reviewing available information on the physiography, climatology, and vegetation of the Mexican gulf versant, we selected several areas in the Rio Grande basin for survey in late August and early September of last year, when rainfall and temperature conditions would be most favorable for the breeding of *Haemagogus* (1). One of these areas was the delta region of the Rio Grande in the vicinity of Brownsville, Tex. This area is largely under intensive cultivation, but we were able to find occasional patches of thorny scrub vegetation along relatively moist depressions that are locally known as "resacas." Larvae and pupae of *Haemagogus equinus* were collected from water in three tree holes in a patch of thorn scrub off Texas State Highway 48 near the intersection with Farm Road 1792, 5 miles northeast of Brownsville (4 and 6 Sept. 1955); and from a tree-hole 15.7 miles east of Brownsville on Boca Chica Boulevard (6 Sept. 1955). By 8 Sept., adult males and females had already emerged. This material will be deposited in the United States National Museum and the collection of the Gorgas Memorial Laboratory. Because of the pressure of other field work scheduled in Mexico, no attempt was made to seek *Haemagogus* further north in Texas.

Haemagogus equinus, which occurs at least as far south as Colombia, is a proved vector of yellow fever in the laboratory, but virus has not been recovered with certainty from it in nature. It was, however, the only species of *Haema-*

gogus found by us in immediate association with the epizootic of yellow fever on the northern coast of Honduras in 1954 (2).

HAROLD TRAPIDO

PEDRO GALINDO

Gorgas Memorial Laboratory,
Panama, Republic of Panama

References and Notes

1. This investigation was supported by the Research and Development Division, Office of the Surgeon General, Department of the Army, under contract No. DA-49-007-MD-655.
2. H. Trapido and P. Galindo, *Am. J. Trop. Med. Hyg.* 4, 665 (1955).

27 September 1955

Loss of Sebaceous Glands in Skin of Thiamine-Deficient Mice

To study the effects of thiamine deficiency in mouse skin with resting or growing hair follicles, approximately 60 young adult C57 black mice of both sexes were used. The mice were kept in individual wire metabolism cages and offered tap water ad libitum.

Two kinds of thiamine-deficient diets were used—diet 227 (Table 1), obtained from Paul Fenton of Brown University (1) and a diet purchased from General Biochemicals, Inc. (Table 2). Pair-fed animals—that is, animals fed the normal diet in amounts equal to those consumed by their respective paired mates that were fed the thiamine-deficient diet, served as controls.

Biopsy specimens were removed on approximately the 7th, 14th, and 21st days of the deficiency regimen. The skin was shaved, and approximately 1 cm² was removed, spread on a piece of cardboard, cut in half, and fixed in 10-percent formal calcium. One half was prepared for histological study and stained in an

Table 1. Components of diet 227. In the control diet, 10 mg of thiamine was added to each 1000 g of diet 227.

Component	Amount
Casein (vitamin-free)	300 g
Dextrin	500 g
Salts (Hubbel <i>et al.</i>)	50 g
Corn oil	50 g
<i>Vitamin supplement per 1000 g of diet</i>	
Pyridoxine	10.0 mg
Riboflavin	20.0 mg
Nicotinic acid	50.0 mg
Biotin	0.2 mg
Folic acid	5.0 mg
Pantothenic acid	100.0 mg
Vitamin A	10,000 U.S.P. units
Vitamin D	1,000 U.S.P. units
Vitamin E	50.0 mg
Vitamin K	10.0 mg
Choline	1.50 gm