




Fig. 2. A comparison of the ways in which changes in the potentiating process are expressed by input-output curves (A) and curves representing the time-course of PTP (B). Two levels of excitability are shown: "Maximal" in which the potentiated discharge reaches the maximum over all degrees of PTP and "submaximal" in which the potentiated discharge never reaches the maximum. The curves are generalized from 31 spinal cord experiments.

curves (A) during PTP tend toward this ceiling, increases or decreases in the potentiation process, as revealed by these curves and reflecting changes in synaptic threshold (C), may not be fully indicated by the method of testing employed in B. Curves expressing these relationships are shown in Fig. 2. In the "maximal" region, changes which fail to lower the input-output curves from the ceiling give no evidence of an alteration in PTP when tested in the usual manner, except for small changes in the time-course of potentiation.

The curves of Fig. 2 are based on results obtained in a study of the effects of diphenylhydantoin (Dilantin) on synaptic transmission (7, 8, 13). Dilantin has negligible effects on the level of reflex excitability of the spinal cord and on synaptic threshold at rest. On the other hand, the drug consistently produces a shift to the right in the input-output curves obtained during PTP. The expected decrease in PTP, however, is revealed only when testing is performed in the "submaximal" zone.

In examining means of avoiding the pitfall just described, consider first the simplest situation, that in which the level of excitability at rest is not influenced by the drug. It is apparent that effects of drugs on PTP can be assessed in three ways: (i) Input-output relations can be determined directly, at rest and during PTP. (ii) PTP may be studied in the usual fashion after it has been determined that testing is being carried out in the "submaximal" zone. If the initial level proves unsatisfactory (that is, in the "maximal" zone), then a more desirable level may be found by raising the temperature (see, for example, Fig. 1A)

or, alternatively, by reducing the number of afferent fibers subjected to stimulation. (iii) PTP may be tested in the usual way but following a tetanus with either frequency or duration reduced so that the maximum potentiated discharge is less than that produced by maximal tetanic parameters (2).

A further difficulty in interpreting effects on PTP is introduced if the drug increases or decreases the discharge zone at rest, inasmuch as equivalent degrees of PTP are not to be expected at different levels of excitability (12, 14). Figure 1D shows the relationship between PTP ratio and maximal discharge zone at rest (pretetanic) which holds for the data of Fig. 1A. The additional complication of change in discharge zone can be dealt with in several ways. Qualitative statements regarding the effect of a drug are justified if both PTP ratio and discharge zone at rest are influenced in the same direction by the drug (Fig. 1D).

For more quantitative evaluation of the effect of the drug it is often possible to vary excitability (for example, by changing temperature) so that the discharge zone is restored to the control level. PTP may then be compared with the control determination. In altering level of excitability to restore the initial discharge zone the assumption is implicit that the procedure employed to vary excitability (for example, temperature) does not specifically affect the potentiating process. The soundness of conclusions derived in such experiments depends on the validity of this assumption.

The foregoing considerations refer specifically to 2N pathways of the spinal cord. For the most part, however, they

apply also to sympathetic ganglia; the postsynaptic neuronal pool is limited, and a similar relation exists between PTP ratio and discharge zone at rest (3, 13). Furthermore, Dilantin reduces PTP at this site only when testing is in the "submaximal" zone. It is probable that the considerations discussed apply in a general way to PTP in polysynaptic pathways of the spinal cord (4) and in all other synaptic systems in which, under the experimental conditions employed, limitations on efferent pool size exist.

It is therefore apparent that drugs studied for modification of PTP must be tested in a manner capable of revealing an effect if one exists. In addition, if the agent changes the level of excitability, a nonspecific effect on PTP and on PTP ratio is to be expected. Arguments for a specific effect of a drug on the potentiating process must take account of these relationships.

DON W. ESPLIN

Department of Pharmacology,  
University of Utah College of Medicine,  
Salt Lake City

#### References and Notes

1. J. C. Eccles, *The Neurophysiological Basis of Mind* (Clarendon Press, Oxford, 1953).
2. D. P. C. Lloyd, *J. Gen. Physiol.* 33, 147 (1949).
3. M. G. Larrabee and D. W. Bronk, *J. Neurophysiol.* 10, 139 (1947).
4. V. J. Wilson, *J. Gen. Physiol.* 39, 197 (1955).
5. J. C. Eccles and W. Rall, *Proc. Roy. Soc. B138*, 475 (1951).
6. C. N. Latimer, *Federation Proc.* 14, 19 (1955).
7. This investigation was supported by a research grant (B-872) from the National Institute of Neurological Diseases and Blindness.
8. D. W. Esplin, *Federation Proc.* 14, 337 (1955).
9. A. Jefferson and A. Benson, *J. Neurophysiol.* 16, 381 (1953).
10. D. P. C. Lloyd, *ibid.* 6, 143 (1943).
11. W. Rall, *J. Cellular Comp. Physiol.* 46, 373, 413 (1953).
12. K. Koizumi, J. L. Malcolm, C. McC. Brooks, *Am. J. Physiol.* 179, 507 (1954).
13. D. W. Esplin, in preparation.
14. W. Rall, *J. Physiol.* 125, 30 (1954).

8 October 1956

#### Correction

Two corrections should be made in the item "Blood groups and disease" [*Science* 124, 674 (12 Oct. 1956)] which reports a paper on associations between blood groups and disease read to the British Association for the Advancement of Science.

The three diseases for which the evidence is overwhelming are (i) duodenal ulcer, (ii) gastric ulcer, (iii) cancer of the stomach. As is stated, the ulcers are commoner in persons of group O, and cancer of the stomach in persons of group A. There is fairly strong evidence in connection with diabetes mellitus and pernicious anemia; but the incidence is raised in group A, not groups O and B.

It should perhaps be added that practically all the data so far come from Western Europe and North America, and the associations might be different or absent in other peoples.

J. A. FRASER ROBERTS  
London School of Hygiene and Tropical  
Medicine, London, England