SCIENCE

21 December 1956

Volume 124, Number 3234

Editorial	Society in the Scientific Revolution	1231
Articles	Why Biophysics?: A. V. Hill	1233
	Stokes' Edge Waves in Lake Michigan: W. L. Donn and M. Ewing	1238
	Proteins and Calcium in Serums of Estrogen-Treated Roosters: O. A. Schjeide and M. R. Urist	1242
	C. Neuberg, Biochemist: F. Lipmann, F. F. Nord, H. Waelsch	1244
News of Science	Statement on Hungary; NSF Employment Profile of Scientists, 1954–55; USIA University Catalog Program; Cancer Mortality Decline; Army Ionizing Radiation Center; NBS Boulder Radio Propagation Engineering Division Reorganized; FWS Study of Fish Chemistry; News Briefs; Scientists in the News; Recent Deaths; Education; Grants, Fellowships, and Awards; In the Laboratories; New Journals	1245
Reports	Influence of Adult Tissue Homogenates on Formation of Similar Embryonic Proteins: H. Walter, D. W. Allmann, H. R. Mahler	1251
	Uronic Acid of Chondroitin Sulfate B: P. Hoffman, A. Linker, K. Meyer	1252
	Anomalous Carbon-14 Content of Carbon Dioxide from Sewer Gas: R. L. Patterson, Jr., and I. H. Blifford, Jr	1252
	Modified End-Window Counting Tube for Paper Chromatograms: R. C. Fuller	1253
Book Reviews	Microscopium; Flow of Gases through Porous Media; Air Pollution Hand-Book; Determination of Organic Compounds; Computers; The Image; New Books; Miscellaneous Publications	1254
tings and Societies	High-Polymer Chemistry; World Federation for Mental Health; Darcy Centennial Symposia; Conference on Scientific Information; Meeting Notes; Society Elections; Forthcoming Events	1257
	Equipment News	1263

Mee

TERMATUBET A NEW CONCEPT IN CLINICAL CHEMISTRY!

Lyophilized, single assay, reagent tubes for determination of serum TRANSAMINASE, URIC ACID and LACTIC ACID DEHYDROGENASE

ACCURATE

SIMPLE

LOW-PRICED

(merely add water, serum and read)

Although developed for use with a spectrophotometer, a spectrophotometer is not necessary for Transaminase or LDH Determinations. The tubes may be used as cuvettes in the new, low priced COENZOMETER (pat. pending) which measures absorption at 340 mm. This instrument is currently being introduced by Macalaster Bicknell Company, 243 Broadway, Cambridge, Mass.

THE NEW, RAPID MICRO METHOD
OF GLUCOSE ANALYSIS
REQUIRING NO DEPROTEINIZATION or HEATING

Based on the coupled enzyme system of glucose oxidase and peroxidase, the reaction is specific for glucose.

A blood glucose can be obtained in 11 minutes using 0.1 ml. serum or plasma. A photoelectric colorimeter is the only piece of special apparatus necessary. Simply add serum to reagent and read color in 10 minutes.

With proper precautions, the method is suitable for urine glucose determinations.

SIMPLE SPECIFIC QUANTITATIVE

Write to us for complete information including coenzometer literature

WORTHINGTON BIOCHEMICAL CORP. FREEHOLD, N. J.

