

scended testes than in scrotal testes (4, 5), and two varieties of ovarian tumor, the tubular type of arrhenoblastoma and the dysgerminoma, are often seen in association with the pseudohermaphrodite state (6). It has also been reported that the testes in which tumors develop may exhibit aberrant differentiation in the form of tubules lined by immature Sertoli cells (7).

A study was undertaken of the nuclear sex of patients in whom testicular tumors had developed. Seventy-five cases were taken from the testicular tumor registry at the Armed Forces Institute of Pathology; the criteria of selection were that normal testicular tissue must be present in the sections and that preoperative radiation therapy must not have been employed. Most of the patients were members of the U.S. Armed Forces and were accepted both at their induction physical examination and by their fellows as males; only two showed cryptorchidism. All varieties of germinal testicular tumors were represented in the group; 29 were seminomas, 14 were embryonal carcinomas, 25 were teratocarcinomas, and 7 were of mixed pattern. This distribution of histological types corresponds to that of the much larger series of Dixon and Moore (4).

Sections were stained by hematoxylin and eosin and by the Feulgen technique. In each instance Leydig cells, Sertoli cells, and the cells of the supporting connective tissue were examined. In all cases the nuclear sex was diagnosed as male. In the 25 teratomas, sexing of the tumor was carried out, and 7 contained female elements. These findings indicate that, in the genesis of testicular tumors, little part is played by gross error of sexual differentiation, and that the "crossed sex" teratomas cannot be explained by the thesis that "female" tumors arise in "female" patients.

DAVID J. B. ASHLEY
ERICH A. THEISS

Armed Forces Institute of Pathology
and Veterans Administration Central
Laboratory for Anatomical Pathology
and Research, Washington, D.C.

References

1. M. L. Barr and E. G. Bertram, *Nature* 163, 676 (1949).
2. W. F. Hunter and B. Lennox, *Lancet* 1, 372 (1954).
3. D. B. Cruikshank, *ibid.* 1, 253 (1955); A. R. Sohval and J. A. Gaines, *Cancer* 8, 896 (1953); A. S. Tavares, *Lancet* 1, 948 (1955); K. L. Moore and M. L. Barr, *Brit. J. Cancer* 9, 246 (1955); L. Myers, in "Symposium on nuclear sex," see *Lancet* 2, 637 (1957).
4. F. J. Dixon and R. A. Moore, "Tumors of the male sex organs," in *Atlas of Tumor Pathology* (Armed Forces Institute of Pathology, Washington D.C., 1952), sect. VIII, fasc. 31b and 32.
5. J. B. Gilbert and J. B. Hamilton, *Surg. Gynecol. Obstet.* 71, 731 (1940).
6. E. Novak, *Gynecological and Obstetrical Pathology* (Saunders, Philadelphia, ed. 3, 1950), p. 402.
7. A. R. Sohval, *J. Urol.* 72, 693 (1954).

17 July 1958

5 DECEMBER 1958

Effects of Magnesium and Tetraethylammonium Chloride on the Hypothermic Heart

Abstract. Ventricular fibrillation was induced by the injection of magnesium chloride in isolated canine hearts as well as in intact animals. Ventricular magnesium tolerance was greatly reduced during hypothermia. Defibrillation was achieved by intracoronary injection of tetraethylammonium chloride and electrical shock, even at temperatures as low as 21°C, where it could not be achieved by other means.

Great advances in cardiac surgery have been achieved by the use of induced hypothermia. However, there remains the hazard of irreversible ventricular fibrillation at low temperatures.

In a series of experiments (1) on myocardial excitability in this laboratory, where both a nonfailing isolated canine heart preparation and intact animals were used (2), both at normothermic and hypothermic levels, it was found that magnesium chloride (but not sodium or potassium chloride) in amounts of 1 to 2 milliequivalents injected rapidly into the coronary system invariably produced ventricular fibrillation. The fibrillation threshold to magnesium was greatly reduced during hypothermia.

In a recent publication by Stovner (3) it was shown that a block at the neuromuscular junction produced by an excess of magnesium chloride was released by tetraethylammonium chloride and that the magnesium-induced block was temperature-dependent. This led us to examine the effects of tetraethylammonium chloride on normothermic and hypothermic hearts, with special reference to its effect on ventricular fibrillation. In 22 experiments involving 20 dogs, which included isolated heart experiments as well as single, intact animal experiments, it was found that tetraethylammonium chloride invariably allowed electrical defibrillation at temperatures as low as 21°C, and that defibrillation was achieved with great ease, whereas, without the use of tetraethylammonium chloride, defibrillation could not be achieved. Furthermore, the minimum amount of magnesium required to induce fibrillation was more than doubled after the use of tetraethylammonium chloride under these conditions.

The tetraethylammonium chloride dosages introduced into the coronary system were from 1 to 2 mg/kg of body weight. This dose is small enough to avoid those side effects, such as hypotension and gross electrocardiographic changes, which have been associated with the clinical parenteral dosage of 10 mg/kg of body weight. Other cardiovascular effects of tetraethylammonium chloride observed included decrease of coronary venous

magnesium content and increase of coronary blood flow, as well as a strong positive inotropic action; the last observation confirms previous work by Acheson and Moe (4).

Our results seem to suggest an important role for magnesium in the mechanism of ventricular fibrillation during hypothermia. Its effects might be exerted through a slowing of conduction or by interference with the effect of transmitter substances at the membrane level. Its fibrillatory action is apparently directly antagonized by tetraethylammonium chloride. In addition, tetraethylammonium chloride increases myocardial contractility, probably by increasing coronary flow as well as by acting directly on the muscle.

In four human beings undergoing cardiac surgery for correction of congenital or acquired heart defects under artificial hypothermia, electrical defibrillation of the ventricles has been achieved only after introduction of 1 to 2 mg of tetraethylammonium chloride per kilogram of body weight, by coronary perfusion (5).

PAOLO CALDINI

BARUCH BROMBERGER-BARNEA

Department of Research and
Laboratories, National Jewish Hospital
at Denver, and Department of
Physiology, University of Colorado
School of Medicine, Denver

References and Notes

1. This work is being supported by a grant from the U.S. Public Health Service (H-3844).
2. S. F. Sarnoff, R. B. Case, G. H. Welch, Jr., E. Braunwald, W. N. Stainsby, *Am. J. Physiol.* 192, 141 (1958).
3. J. Stovner, *Acta Physiol. Scand.* 41, 370 (1957).
4. G. H. Acheson and G. K. Moe, *J. Pharmacol. Exptl. Therap.* 87, 220 (1946).
5. The surgery was performed on these patients under the direction of Dr. John B. Grow, chief of surgery, and Dr. George Wittenstein, senior resident surgeon, National Jewish Hospital.

7 July 1958

Ion-Exchange Equilibria on Single Beads

Abstract. Equilibria on single ion-exchange resin beads show that large differences may exist between beads from the same batch. They may be so large that a significant contribution to deviations from ideality can be due to this heterogeneity effect. Correlation between swelling and equilibrium properties shows that bead-to-bead variations are due to differences in cross-linking.

Ion-exchange equilibria on single beads can be studied by radioactive tracer methods (1). Below are given some results for the $\text{Ag}^+ - \text{H}^+$ system on Dowex 50 X-4 resin. The beads were taken from the 20-50 mesh sieve fraction. The tracer was Ag^{110} , obtained by neutron bombardment of silver foil in