SCIENCE ^{3 July 1959} Volume 130, Number 3366

Editorial	In Perspective	7
Articles	Reasoning Foundations of Medical Diagnosis: R. S. Ledley and L. B. Lusted	9
	Mathematical Inutility and the Advance of Science: C. B. Boyer	2 2
	Should science entice the mathematician from his ivory tower into Solomon's House?	
News of Science	More Seismic Research Asked to Improve Atomic Test Detection; Strauss Rejected as Commerce Secretary; Number of Foreign Scholars in	<u>.</u>
	U.S. Increases	26
Book Reviews	Handbook of Physics, reviewed by H. E. Newell, Jr.: other reviews	31
Reports	Lactation Hair in the Asiatic Squirrel and Relationship of Lactation Hair to Mammary Hair: S. O. Landry, Jr.	37
	Voltage Clamp of Motoneuron Soma: K. Frank, M. G. F. Fuortes, P. G. Nelson	38
	Cortical Correlates of Auditory Localization: P. D. Coleman	39
	Nature of Colchicine Resistance in Golden Hamster: A. R. Midgley, B. Pierce, F. J. Dixon	40
	Adsorption of Antibody in vitro and Magnitude of the Schultz-Dale Reaction of Guinea-Pig Ileum: C. B. Nielsen, G. Terres, G. A. Feigen	41
	Variations in Style Number and Other Gynoecial Structures of Lychnis alba: H. L. Dean	42
	Incidence of Sex Chromatin in Gallus domesticus: I. L. Kosin and H. Ishizaki	43
Departments	Letters	4
	Forthcoming Events; New Products	48



photomicrography combining LEITZ quality with unlimited versatility

Leitz ORTHOLUX and ARISTOPHOT, shown with Fluorescence Unit HBO-200

ORTHOLUX *Microscope* with **ARISTOPHOT II**

Here are precision Leitz instruments perfectly matched for exacting photomicrography. They are simple to operate, yet capable of infinite variations to fit all your photomicrographic requirements.

With the Leitz ORTHOLUX with ARISTOPHOT, image sharpness is limited only by the resolution of the film you are using. Only air separates the optical system from the film. There is no loss of image sharpness because of mirrors or other devices.

Film size is not restricted: 35mm Leica, $3\frac{1}{4} \ge 4\frac{1}{4}$, $4\ge 5$, or Polaroid. Your choice of illumination, too, is virtually unlimited: transmitted or incident light, dark field, polarized or phase contrast.

With the ARISTOPHOT detached, the Leitz ORTHOLUX fulfills the most critical requirements for a research microscope. The ARISTOPHOT II may be used separately for gross photography or macrophotography.

Write for illustrated brochure, Dept. 294 SC-7.



E. LEITZ, INC., 468 FOURTH AVENUE, NEW YORK 16, N.Y. Distributors of the world-famous products of Ernst Leitz G. m. b. H., Wetzlar, Germany-Ernst Leitz Canada Ltd. LENSES • CAMERAS • MICROSCOPES • BINOCULARS

Saunders offers your students up-to-date and well organized texts in the biological sciences

New! Guyton-FUNCTION OF THE HUMAN BODY

This new text clearly covers human physiology for the college student, stressing the mechanisms of action and their interrelationships. The logical organization of function in the body is traced in step-by-step detail, progressing from cellular function through detailed descriptions of the physiology of each human organ system. The author stresses the many regulatory mechanisms that make it possible for the cells to function without too many mishaps throughout a person's lifetime. Selected abnormalities are discussed as examples of applied physiology. Practical answers are given to such questions as: What enables people to move? How is it possible for them to talk? How can they see the expanse of the world and feel the objects about them? How is energy derived from the food they eat? By what process do humans reproduce other beings like themselves?

By ARTHUR C. GUYTON, M.D., Professor and Chairman of the Department of Physiology and Biophysics, University of Mississippi School of Medicine. 584 pages, 307 illustrations. \$7.50. New l

New! Hall-FUNCTIONS OF THE ENDOCRINE GLANDS

Here is a remarkably clear presentation of the normal functioning of the human endocrine system. The author clearly coordinates all present knowledge on the physiology and biochemistry of the subject. Functions of the various glands are taken up and described in turn—the adrenals, thyroid, testes, ovaries, pancreas, parathyroids, etc. Dr. Hall points out the endocrine factors in growth, maturation and decline. He covers important relationships between the central nervous system and the endocrine glands. An entire chapter is concerned with the investigation of steroid metabolism. A separate section deals with the chemistry of endocrinology. The only book of its kind in the field, this is a text for medical students, graduate students in zoology and all who require a sound background in human endocrinology.

By PETER F. HALL, M.D., M.R.C.P., M.R.A.C.P., Assistant Physician, Sydney Hospital, Assistant Lecturer in Physiology, University of Sydney, Australia. 290 pages, illustrated. \$5.75. New !

New (2nd) Edition! Odum-FUNDAMENTALS OF ECOLOGY

A popular college text, this New (2nd) Edition presents a well-balanced synthesis of the entire field of ecology—combining the *functional* and the *descriptive*, the *aquatic* and *terrestrial*, as well as the *basic* and *applied* aspects of the subject. A truly integrated picture of plants, animals and microorganisms, it tells the student what nature "does" as well as how she "looks." New material is included on the ecosystem, energy relations, microbial ecology and terrestrial plant ecology. A new chapter on *Radiation Ecology* represents the first comprehensive treatment of this important field. 60 new illustrations have been added.

By EUGENE P. ODUM, Alumni Foundation Professor of Zoology, University of Georgia, Athens, Georgia; in collaboration with HOWARD T. ODUM, Director, Institute of Marine Sciences, University of Texas, Port Aransas, Texas. 546 pages, 160 illustrations. \$7.50. New (2nd) Edition t

Gladly sent to college teachers for consideration as texts W. B. SAUNDERS Company, West Washington, Square, Philadelphia 5

SCIENCE is published weekly by the AAAS, 1515 Massachusetts Ave., N.W, Washington 5, D.C. Entered at the Lancaster, Pa., Post Office as second class: matter under the act of 3 March 1879. Annual subscriptions: \$8.50; foreign postage, \$1.50; Canadian postage, 75¢.



A new achievement for STEREOMICROSCOPY

The WILD* M5 STEREOMICROSCOPE presents, new, important advances in versatility, optics, mechanical conveniences and physical design.

This Swiss precision instrument is equipped with a main objective component followed by pairs of vertically mounted intermediate lenses with <u>parallel axes.</u> The result is increased, uniform sharpness throughout the field, with no need for any change in accommodation.

With a constant working distance of 96 mm., standard magnifications are 6X, 12X, 25X and 50X, conveniently selected on a horizontal drum.

Accessories include a base for transmitted light observation, various light sources, photographic and measuring attachments. A matching steel hood is provided for easy storage and portability.

For full details about this yearsahead stereomicroscope, write for Booklet M5.

The FIRST name in Surveying Instruments, Photogrammetric Equipment and Microscopes



4

Letters

Terms for Temperatures

The Appleman-Braham interchange of viewpoints on nomenclature for temperature reductions that go below a predetermined reference point [Science 129, 1296 (1959)] prompts me to observe that none of the terms suggested are as free of ambiguity as they might be, and that some of them can be misleading. The point of contention was the choice between sub, super, and under as a prefix to the verb cool for description of such temperature reductions.

My first point is that the verb itself can be improved. I have found the verb *chill* to be more descriptive in speaking of temperature drops considerably below ambient levels, admittedly not to the same degree that *heating* rather than *warming* distinguishes a substantial rise in temperature, but still enough to establish a suitable distinction in the mind of the reader. English does not provide a common verb that bears quite the same relationship to *cool* that *heat* does to *warm*.

The use of *under* as a prefix for either verb conveys to me an image exactly opposite to the one intended; a system which is "undercooled" means one that experienced a temperature drop *less* than the magnitude to be expected from the context. To a lesser extent, *sub* as a prefix suffers the same handicap.

On the other hand, the prefix super implies a marked deviation from the norm, which is not true necessarily for the downward departures from freezing points or saturation temperatures that are meaningful in the systems usually encountered. Over describes such a departure better, because it can connote any deviation, no matter how small. It also avoids Braham's objection—a valid one in my opinion—to mixing words of different derivation.

To say that water or a saturated solution is "overchilled" registers in my mind a sharp impression that the system is at temperatures near a critical point (not just "cool"), and that it is to some degree (not necessarily great), below this reference temperature.

H. LEROY THOMPSON Birmingham, Alabama

European Degrees

I should like to support Seiden's opinion [Science 129, 933 (1959)] that it would be useful from the standpoint of international scientific contacts if some professional group would attempt to standardize the anglicization of continental degrees. May I suggest that it be considered whether the American Association for the Advancement of Science, representing if possible also the American Medical Association, American Institute of Biological Sciences, American Institute of Chemists, American Chemical Society, and other interested U.S. organizations, could undertake such an endeavor, specifically in collaboration or consultation with the analogous British societies.

D. A. A. MOSSEL Central Institute for Nutrition and Food Research, Utrecht, Netherlands

Marine Fungi and Limnoria

The recent work by D. L. Ray and D. E. Stuntz [Science 129, 93 (1959)] contains some suggestions and implications concerning our beliefs that need to be corrected.

First, we have not claimed that "marine wood-boring animals do not attack wood or become established in it unless the wood is first invaded and 'conditioned' by marine fungi." However, certain of our studies referred to have indicated that a thorough examination of the role of marine fungi in the destruction of wood certainly is due.

Second, we have not made any statement or implication that "Limnoria is unable to attack sterilized wood." There are so many unknown factors in the evidence so far presented that any statement based on this concept would be drawing essentially unsubstantiated conclusions.

Third, we have not expressed a belief that "Limnoria will not attack wood until its surface has been 'conditioned'." This "conditioning," if it is of any effect, surely involves physical, chemical, and biological factors, none of which has yet been given adequate study.

A reasonably careful reading of our two articles would show clearly that we were presenting specific observations and results of experimentation concerning marine fungi and were calling attention to the need for careful studies of the interrelationships between the marine organisms which infest submerged wood. We do believe that deterioration is a composite process in which the contribution made by any organism involved in the biologic complex, at any stage in the process, should be given critical evaluation.

> Ernest S. Reynolds Samuel P. Meyers

Marine Laboratory, University of Miami, Miami, Florida

(Continued on page 46)



NEW B&L STEREOMICROSCOPES Exclusive StereoZoom continuously variable power. Erect views in natural 3-D.





B&L "ST" MICROSCOPES Standard size and operation. Exclusive pre-focus gage saves time.

> **B&L MACROSCOPES** Ideal for gross studies, dissection, field trips. $10\times$, $20\times$ or $30\times$.

B&L TRI-SIMPLEX MICRO-PROJECTOR Projects mounted specimens or live organisms, on screen or tracing pad.

B&L SPECTROSCOPES Basic chemical analysis tool; shows spectra of elements.



B&L BALOPTICON® PROJECTORS Big bright views of slides, opaque objects, even chemical reactions!

Suddenly... SCIENCE MAKES SENSE!

With Bausch & Lomb educational instruments, students actually *see* what you're talking about. Science becomes easier to understand. They learn faster, more thoroughly, for keeps. See for yourself how B&L instruments—the science teaching standard—help you cram more real learning into every hour.

WRITE for Catalog E-152. Demonstration, too, on your request. Bausch & Lomb Optical Co., 64243 St. Paul St., Rochester 2, New York.



-(Connaught)-

STARCH-HYDROLYSED BORATE BUFFER

STARCH-HYDROLYSED is specially prepared by the Connaught Medical Research Laboratories for use in the starch-gel method of zone electrophoresis developed by Dr. O. Smithies (Biochemical Journal 61, 629, 1955 and 71, 585, 1959). The method is specially adapted for investigation of protein solutions and is particularly valuable for the study of serum proteins.

A concentrated BORATE BUFFER is now available to facilitate the preparation of suitable gels. It should be diluted with distilled water to provide the specific concentration shown on the label of each bottle of Starch-Hydrolysed.

Starch-Hydrolysed Borate Buffer Code Word GEL

BORATE

Container Size

2,000-gram bottle 500-cc plastic bottle

CONNAUGHT MEDICAL RESEARCH LABORATORIES University of Toronto Toronto, Canada

Established in 1914 for Public Service through Medical Research and the development of Products for Prevention or Treatment of Disease.

NEW MICRO-STROPPER ELECTRIC MICROTOME KNIFE STROPPER*

STROPS KNIFE TO KEEN, UNIFORM CUTTING EDGE



Much faster and more accurate than old handstropping methods. Merely run blade over special compounded wheel four or five times each side. Knife holder and support plate adjustable for differences of bevel on various knives. For further data write for Technical Bulletin 23-122.

60070. Grey finished housing, $13\frac{1}{2}$ inches wide, 11 inches deep, 9 inches high. For 115 volts A. C. Complete with charged stropping wheel and extra supply of stropping abrasive for recharging. Each \$275.00.



5655 KINGSBURY, ST. LOUIS 12, MO. 14 STOCKED DIVISIONS COAST-TO-COAST A NEW USSR ACADEMY OF SCIENCES JOURNAL



beginning with January 1959 issue at new low rates

Publishing results of experimental and theoretical investigations by leading Soviet Scientists. Articles in all branches of optics and spectroscopy, including X-ray, ultraviolet, visible, infrared and microwave, thin layer optics, filters, detectors, diffraction gratings, electro-luminescence, thermal radiation backgrounds, infrared polarizers and many applications to other branches of science and to industry.

Translated and published by the OPTICAL SOCIETY OF AMERICA, this branch of Soviet Science is now made available to all interested individuals and organizations along with the Journal of the Optical Society of America at the rate of a single journal alone. This was made possible by a grant-in-aid from the NATIONAL SCIENCE FOUNDATION to the OPTICAL SOCIETY OF AMERICA. Comments on the Soviet articles will appear in the Letters to the Editor column of the Journal of the Optical Society of America.

Associate membership dues Optical Society of America Non-member subscription	U.S. & Canada (both journals) \$13.00 \$25.00	Elsewhere (both journals) \$13.00 \$28.00
---	--	--

For membership in the OPTICAL SOCIETY OF AMERICA please write to Dr. K. S. Gibson, Secretary, Optical Society of America, National Bureau of Standards, Washington 25, D. C. Qualification for membership consists of "an interest in optics."

For non-member subscriptions write to the AMERICAN INSTITUTE OF PHYSICS, 335 E. 45th Street, New York 17, New York.

The Beginnings of Embryonic Development

AAAS Symposium Volume No. 48

Published July 1957

Edited by Albert Tyler, California Institute of Technology R. C. von Borstel, Oak Ridge National Laboratory Charles B. Metz, The Florida State University

6 x 9 inches, 408 pages, 132 illustrations, references, subject and author index, clothbound

Price \$8.75, AAAS members' prepaid order price \$7.50

A symposium on "Formation and Early Development of the Embryo", held 27 December, 1955, at the Second Atlanta Meeting of the AAAS, served as the basis for this volume. Emphasis was placed on the problems of early development and of the initiation of development. The investigations presented in the various communications cover both descriptive and experimental work on the biological and chemical levels. Apart from their intrinsic interest and the measure of progress that they provide, the specific discoveries and analyses presented serve to exemplify various approaches toward the understanding of the manner in which sperm and egg contrive to produce a new individual.

> British Agents: Bailey Bros. & Swinfen Ltd., Hyde House, West Central Street, London WC1, England

AAAS, 1515 Massachusetts Avenue, NW, Washington 5, D.C.

Letters

(Continued from page 4)

Whether, as Reynolds and Meyers claim, we have erred in our interpretation of the meaning and implications of their work (1, 2) can best be judged by reference to what we said, and to their published reports, on which our remarks were based. We wrote (3):

"Believing that fungal infection always occurs prior to attack by marine wood borers, especially *Limnoria*, they [Meyers and Reynolds] *suggested* that there *might* be a relationship between fungi and wood-destroying animals." (Italics added).

Both parts of this sentence require comment in the light of Reynolds and Meyers' objections. First, the only *belief* that we credit to these authors is that of holding that fungal infection of submerged wood always precedes marine borer attack. Our statement that this is their belief is based upon the following three quotations. In the concluding paragraph of their article in *Science* (1) they said: "The vigorous fungal infestation of

"The vigorous fungal infestation of submerged wood prior to borer attack represents a biological phenomenon that investigators of marine wood destruction should not ignore. In northern areas, winter fungal infestation of wood is evident. Hence, in the early spring, when borer activity increases rapidly, the animals have available a wood substrate thoroughly infected by a variety of marine fungi. The interrelationships within this biota are being studied in our laboratory."

And in the same article they wrote: "Vigorous attack [by fungi] upon submerged wood in boreal and northern temperate areas during winter months is accompanied by no, or very slight, borer damage. A similar situation occurs in subtropical localities, however, with a considerably shorter period of fungal attack prior to borer infestation."

Finally, the following sentences appear in another article (2, p. 10, paragraph one):

"The fungal infection that occurs before the borers attack the wood has interesting biological implications. In northern areas, especially, it may facilitate the activities of the borers, not only by making it easier for them to enter the wood, but also by providing them with a source of food."

We still think that it is fair to say that Reynolds and Meyers indicated a belief that fungal infection always precedes attack by marine borers. Concerning the second half of our sentence, we believe that the excerpts quoted above and other comments in the two articles cited do support our statement that Reynolds and Meyers suggest that there may be a relationship between marine fungi and marine wood borers.

With reference to the three specific objections enumerated in their letter, we will respond to the second one first, for here Reynolds and Meyers have legitimate grounds for complaint. In our article in Science, we inadvertently cited the report that "Limnoria is unable to attack sterilized wood" as reference number 3 [Reynolds and Meyers (2)], but it should have been cited as reference 4 [Schafer and Lane (4)]. We apologize for this typographical error in the bibliographic citation. To set the record straight, Schafer and Lane (4), reporting work done at the Marine Laboratory of the University of Miami, state in their abstract: "Limnoria of both sexes and all ages did not survive when they were allowed to feed only on sterile wood in sterile sea water. No fecal pellets were produced under the conditions of this experiment." Supporting evidence is given in the text of the paper.

Reynolds and Meyers' first listed objection is that we included their work among the cases where we said it has been ". . . stated, suggested, or implied that marine wood-boring animals do not attack wood or become established in it unless the wood is first invaded and 'conditioned' by marine fungi" (3). Immediately preceding this comment we cited

work from three different groups of investigators. Perhaps it would have been more precise to say that Becker, Kampf and Kohlmeyer (5) stated, Schafer and Lane (4) suggested, and Reynolds and Meyers (1, 2) implied.

It is our opinion that Reynolds and Meyers' work reported in their two papers does indeed convey the implication that marine wood-boring animals do not attack wood or become established in it unless the wood is first invaded and "conditioned" by marine fungi. A careful rereading of these papers has not altered this opinion. Independent evidence that we are not alone in this interpretation may be found in the table of contents of the issue of ONR Research Reviews in which one of their papers (2) appears. Following the title is this explanatory statement (authorship unknown):

"Investigations now underway show that marine fungi are one of the chief contributors to the deterioration of submerged wood. They not only vigorously degrade wood, but also prepare it for the entrance of other wood destroyers."

Finally, regarding the third point made by Reynolds and Meyers in their letter, we did not claim that they "expressed a belief" that Limnoria will not attack wood until its surface has been "conditioned." (We do not believe this either!) Our reference to their work in this connection was only to the observation that a period of time elapses between the exposure of wood to sea water and its invasion by Limnoria.

The many questions of keen interest that arise from the results we have submitted, conflicting as they do with those from other investigators, deserve fuller discussion than is appropriate in this letter or was possible in our report. Until such further discussion can be engaged in, we wish only to make our position abundantly clear. We have undertaken a critical evaluation of just one point: whether there is any relationship between marine fungi and Limnoria attack. Our results indicate that there is not. We do not believe that this closes the case; it is necessary now to determine why our results differ from those of other workers.

D. L. RAY

D. E. Stuntz

Departments of Zoology and Botany, University of Washington, Seattle References

- S. P. Meyers and E. S. Reynolds, Science 126, 969 (1957).
 E. S. Reynolds and S. P. Meyers, Office Naval Research, Research Revs. (Dec. 1957), pp. 6-11.
 D. L. Ray and D. E. Stuntz, Science 129, 93 (1969)
- (1959). 4. R. D. Schafer and C. E. Lane, Bull. Marine
- K. D. Schult and Caribbean 7, 289 (1957).
 G. Becker, W.-D. Kampf, J. Kohlmeyer, Naturwissenschaften 17, 473 (1957).

"IDENTIFICATION AND CHEMICAL ASSAY OF NOREPINEPHRINE IN BRAIN AND OTHER TISSUES"*

"A simple, rapid fluorometric method for the estimation of catechol amines in brain and other tissues is described

The procedure involves extraction of catechol amines from tissue homogenates into butanol. After returning the amines to an aqueous phase, they are oxidized to form fluorescent derivatives which are measured in a spectrofluorometer."



Activation spectra (left) and fluorescence spectra (right) of various catechols and rabbit brain extracts. To obtain the activation spectra, the fluorescence

monochromator of a Farrand Recording Spectrofluorometer was set at 520 m μ and the spectra from the activating monochromator were scanned. To obtain the fluorescence spectra, the activating monochromator was set at 400 $m\mu$ and the spectra from the fluorescence monochromator were scanned.

*Ref: Parkhurst A. Shore and Jacqueline S. Olin, Journal of Pharm. and Experim. Therapeutics, Vol. 122, No. 3.



Spectrofluorometric assay has distinct advantages over physical methods using light absorption. Fluorescent methods, where applicable, are far more discrim-inating than either Colorimetric or Spectrophotometric assay and yield a sensitivity at least two orders of magnitude greater than Spectrophotometry.



Bulletin No. 820 Supp. 1 sent on request

RADE MAD



