

- ff. The growth of the understanding of the realm of the explainable, from the end of the 13th century on, can be traced through almost every chapter of this book.
3. C. F. von Weizsäcker, *Z. Astrophys.* **22**, 319 (1944); S. Chandrasekhar, *Rev. Mod. Phys.* **18**, 94 (1946).
 4. An interesting and well-understood case is that of "focusing collisions" in which neutrons, having velocities which are rather high but with random orientation, are converted into lower-velocity neutrons but with preferential directions of motion. See R. H. Silsbee, *J. Appl. Phys.* **28**, 1246 (1957); C. Lehmann and G. Leibfried, *Z. Physik* **172**, 465 (1963).
 5. See, for instance, the section "What is the state vector" in E. Wigner, *Am. J. Phys.* **31**, 6 (1963).
 6. The possibility of an invariance principle in which velocities are replaced by positions, and conversely, was studied by M. Born, *Nature* **141**, 327 (1938); *Proc. Roy. Soc. London, Ser. A* **165**, 291 (1938); *ibid.* **166**, 552 (1938).
 7. The crossing relations were established by M. L. Goldberger, *Phys. Rev.* **99**, 979 (1955); M. Gell-Mann and M. L. Goldberger, *ibid.* **96**, 1433 (1954). For further literature, see, for instance, M. L. Goldberger and K. M. Watson, *Collision Theory* (Wiley, New York, 1964), chap. 10. The relations of the various types of symmetry principles were considered in two recent articles: *Nuovo Cimento Suppl.*, in press, and *Phys. Today* **17**, 34 (1964). See

- also E. Wigner, *Progr. Theoret. Phys.* **11**, 437 (1954).
8. V. A. Fock, *The Theory of Space, Time and Gravitation* (Pergamon, New York, 1959). The character of the postulate of invariance with respect to general coordinate transformations as a geometrical invariance was questioned already by E. Kretschman, *Ann. Phys. Leipzig* **53**, 575 (1917).
 9. M. A. Melvin, *Rev. Mod. Phys.* **32**, 477 (1960).
 10. A. Einstein, *Zur Elektrodynamik bewegter Körper*, *Ann. Phys. Leipzig* **17**, 891 (1905).
 11. — and S. B. Preuss, *Akad. Wiss.* pp. 778, 799, 844 (1915); *Ann. Phys. Leipzig* **49**, 769 (1916). Similar results were obtained almost simultaneously by D. Hilbert, *Nachr. Kgl. Ges. Wiss. Göttingen*, p. 395 (1915).
 12. J. Schwinger, *Phys. Rev.* **76**, 790 (1949). See also S. S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Row, Peterson, New York, 1961), sec. 15, where further references can also be found.
 13. See A. S. Wightman, "Quelques problèmes mathématiques de la théorie quantique relativiste" and numerous other articles in *Les Problèmes Mathématiques de la Théorie Quantique des Champs* (Centre National de la Recherche Scientifique, Paris, 1959).
 14. G. Hamel, *Z. Math. Phys.* **50**, 1 (1904); G. Herglotz, *Ann. Physik* **36**, 493 (1911); F. Engel, *Nachr. Kgl. Ges. Wiss. Göttingen*, p. 207 (1916); E. Noether, *ibid.*, p. 235 (1918); E. Bessel-Hagen, *Math. Ann.* **84**, 258 (1921).

- The quantum theoretical derivation given by E. Wigner, *Nachr. Ges. Wiss. Göttingen*, p. 375 (1927), contains also the parity conservation law which was shown, in reference 1, to be only approximately valid. See also the article of reference 16.
15. I heard this remark, for the first time, from C. N. Yang, at the centennial celebration of Bryn Mawr College.
 16. See E. P. Wigner, "Unitary representations of the inhomogeneous Lorentz group including reflections," in *Elementary Particle Physics*, F. Gürsey, Ed. (Gordon and Breach, New York, 1964) for a systematic discussion of the reflection operations.
 17. See E. P. Wigner, *Gruppentheorie und ihre Anwendung auf die Quantenmechanik der Atomspektren* (Friedr. Vieweg, Braunschweig, 1931) or the English translation by J. Griffin (Academic Press, New York, 1959).
 18. H. A. Kastrup, *Phys. Letters* **3**, 78 (1962). The additional invariance operations probably form the conformal group. This was discovered by E. Cunningham [*Proc. London Math. Soc.* **8**, 77 (1909)] and by H. Bateman [*ibid.* **8**, 223 (1910)] to leave Maxwell's equations for the vacuum invariant, that is, the equations which describe light, always propagating at light velocity. For more recent considerations, see T. Fulton, F. Rohrlich, L. Witten, *Rev. Mod. Phys.* **34**, 442 (1962), and Y. Murai, *Progr. Theor. Phys.* **11**, 441 (1954); these articles contain also more extensive references to the subject.

one plots properties against either the number of neutrons, or the number of protons in the nucleus, rather than against the mass number.

Magic Numbers

One of the main nuclear features which led to the development of the shell structure is the existence of what are usually called the magic numbers. That such numbers exist was first remarked by Elsassner in 1933 (3). What makes a number magic is that a configuration of a magic number of neutrons, or of protons, is unusually stable whatever the associated number of the other nucleons. When Teller and I worked on a paper on the origin of elements, we stumbled over the magic numbers. We found that there were a few nuclei which had a greater isotopic as well as cosmic abundance than our theory or any other reasonable continuum theory could possibly explain. Then we found that those nuclei had something in common: they had either 82 neutrons, whatever the associated proton numbers, or 50 neutrons. Eighty-two and fifty are "magic" numbers. That nuclei of this type are unusually abundant indicates that the excess stability must have played a part in the process of the creation of elements.

We then read Elsassner's papers written in 1933. In the year 1948 much more was known about properties of nuclei than was available to Elsassner. The magic numbers not only stood up

The Shell Model

Maria Goeppert Mayer

There are essentially two ways in which physicists at present seek to obtain a consistent picture of the atomic nucleus. The first, the basic approach, is to study the elementary particles, their properties and mutual interaction. Thus one hopes to obtain a knowledge of the nuclear forces.

If the forces are known, one should in principle be able to calculate deductively the properties of individual complex nuclei. Only after this has been accomplished can one say that one completely understands nuclear structures.

Considerable progress in this direction has been made in the last few years.

The work by Brueckner (1), Bethe (2), and others has developed ways of handling the many-body problem. But our knowledge of the nuclear forces is still far from complete.

The other approach is that of the experimentalist and consists in obtaining by direct experimentation as many data as possible for individual nuclei. One hopes in this way to find regularities and correlations which give a clue to the structure of the nucleus. There are many nuclear models, but I shall speak only of one and leave the others to the next lecture, by Professor Jensen.

The shell model, although proposed by theoreticians, really corresponds to the experimentalist's approach. It was born from a thorough study of the experimental data, plotting them in different ways and looking for interconnections. This was done on both sides of the Atlantic Ocean, and on both sides one found that the data show a remarkable pattern. This pattern emerges if

The author is professor of physics in the School of Science and Engineering, University of California, San Diego. This article is the lecture she delivered in Stockholm, Sweden, 12 December 1963, when she received the Nobel prize in physics, a prize which she shared with Eugene P. Wigner and J. Hans D. Jensen. It is published with the permission of the Nobel Foundation. Copyright © 1964 by the Nobel Foundation. It will also be included in the complete volumes of Nobel lectures in English, published by the Elsevier Publishing Company, Amsterdam and New York.




Fig. 1. Beta decay energies in the neighborhood of $N = 20$.

in the new data but they appeared more clearly than before, in all kinds of nuclear processes. It was no longer possible to consider them as due to purely accidental coincidences.

The magic numbers, as we know them now, are:

2, 8, 20, 28, 50, 82, 126

and, most importantly, they are the same for neutrons and protons. Table 1 shows the magic numbers and, below them, the stable nuclei containing magic numbers of protons or of neutrons.

Tin ($Z = 50$) is the element with the largest number of stable isotopes, namely 11. There are six stable nuclei

with 50 neutrons, and seven with 82 neutrons, whereas normally there are only two or three nuclei with the same number of neutrons.

It has long been known that helium, with two neutrons and two protons, is very tightly bound. An extra nucleon cannot be attached to the helium core—that is, Li^5 and He^5 do not exist. The number 8 is encountered at ^{16}O . It takes an unusual amount of energy to remove a neutron from this nucleus. On the other hand, the ninth, the extra neutron beyond the 8–8 shell, in ^{17}O , is very weakly bound.

For nuclei heavier than Ca^{40} the number of protons is less than that of neutrons, and only then does it become clear that the stability is connected with the neutron number or the proton number, and not with the total number of both.

Let me give just two examples. The first one is taken from the work of Jensen and Suess (4) and is derived from the energy changes in β -decay. Figure 1 shows the energy difference between pairs of isobaric nuclides with neutron excess 3 and 1, with the common mass number as abscissa. The light nuclei, for which the energy difference is positive decay by β^- emission to the nuclei with $N - Z = 1$. For the heavier nuclei, the neutron excess of 3 is the stable isobar, the energy is negative.

One would expect to find a smooth curve, sloping downward. Except for one point, it is indeed so. This point is

Ar^{39} , with 21 neutrons and 18 protons. From a smooth interpretation of the curve one would predict that Ar^{39} is stable, and that its isobar K^{39} is unstable against β^+ emission. However, Ar^{39} is unstable against β^- emission by about 0.5 Mev. The explanation of this anomaly is the low binding energy of the 21st neutron in Ar^{39} , while the 19th proton, into which it is transformed, has the higher binding energy of the proton shell which closes at 20. That the energies drop again sharply is due to the fact that now $Z = 20$ is involved.

These types of discontinuity occur at all magic numbers. Figure 2 shows it at the magic number $N = 50$, where it occurs for various numbers of the neutron excess.

The other example is that of the highest magic number, 126, which occurs only for neutrons, and which was noticed long ago. Again, the prediction is that it would be easy to remove the 127th or 128th neutron, but that it takes a considerable amount of energy to remove the 126th or 125th neutron, whatever the associated proton number. Fortunately, this is the region in which α -decay occurs—in which two neutrons are lost by the nucleus, along with two protons. And the prediction is simply borne out by the facts.

Figure 3 shows the experimental data of the kinetic energy of the emerging α -particle, with the number of neutrons as the abscissa. Isotopes of the same elements are connected by lines. The trend of the curves for the neutron-




Fig. 2. Beta decay energies in the neighborhood of $N = 50$.

rich nuclei is easy to understand. But for all elements the energy reaches its peak at 128 neutrons and then drops sharply when the 126th, and then the 125th, neutron is removed from the nucleus.

From these and similar data one can estimate that the discontinuity of the binding energy at the magic numbers is about 1.5 to 2 Mev.

The Atomic Analogue

The strong binding of a magic number of nucleons and the weak binding for one more immediately brings to mind a similar, but relatively much stronger, effect which occurs in the electronic structure of atoms. The energy required to remove an electron from an atom is measured by the ionization potential. The closed electron shells occur in the noble gas atoms, which have a very high ionization potential. The atoms with atomic number larger by one unit, the alkali, have a very low ionization potential. For instance, for argon, with atomic number 18 and 18 electrons, the energy needed to remove one electron is 15.69 ev, whereas the energy needed to remove the 19th electron from potassium is only 4.32 ev. That is, the binding energy of the last electron in argon is about $3\frac{1}{2}$ times that in potassium. In the case of the nucleons, the change in binding energy across a magic number is only 2 Mev out of an average value of about 6, which is only about 30 to 40 percent. Yet the experimental facts leading to the detection of magic numbers were quite marked, and the numbers could hardly arise from accident. It seemed to be worth while to attempt to explain them in the same way that the noble gases are explained. Indeed, one might try to copy the essential features of the atomic structure for nuclear structure.

The simplest atom is hydrogen, in which one electron is subjected to the spherically symmetrical attraction of one proton. The quantum mechanical levels are characterized by two numbers, of which one, n , is called the principal quantum number. The other one, l determines the angular momentum. By accident, due to the fact that the potential is proportional to the reciprocal of the distance, the energy depends only, or almost only, on the principal quantum number n .

Classically, in a field of spherical symmetry the angular momentum is a constant of the motion. In quantum me-

Table 1. Magic number nuclides.

2	8	20	28	50	82	126
<i>Protons</i>						
He ⁴	O ¹⁶ O ¹⁷ O ¹⁸	Ca ⁴⁰ Ca ⁴² Ca ⁴³ Ca ⁴⁴ Ca ⁴⁶ Ca ⁴⁸	Ni ⁵⁸ Ni ⁶⁰ Ni ⁶¹ Ni ⁶² Ni ⁶⁴	Sn ¹¹² Sn ¹¹⁴ Sn ¹¹⁵ Sn ¹¹⁶ Sn ¹¹⁷ Sn ¹¹⁸ Sn ¹¹⁹ Sn ¹²⁰ Sn ¹²² Sn ¹²⁴	Pb ²⁰⁴ Pb ²⁰⁶ Pb ²⁰⁷ Pb ²⁰⁸	
<i>Neutrons</i>						
He ⁴	N ¹⁵ O ¹⁶	S ³⁶ Cl ³⁷ A ³⁸ K ³⁹ Ca ⁴⁰	Ca ⁴⁸ Ti ⁵⁰ V ⁵¹ Cr ⁵² Fe ⁵⁴	Kr ⁸⁶ Rb ⁸⁷ Sr ⁸⁸ Y ⁸⁹ Zr ⁹⁰ Mo ⁹²	Xe ¹³⁶ Ba ¹³⁸ La ¹³⁹ Ce ¹⁴⁰ Pr ¹⁴¹ Nd ¹⁴² Sm ¹⁴⁴	Pb ²⁰⁸ Bi ²⁰⁹

chanics, the orbital angular momentum is quantized, so that its magnitude in units of Planck's constant $\hbar$ is an integral value l . A level of given l contains $2l + 1$ discrete states of different orientation in space, characterized by an integer m_l with $-l \leq m_l \leq l$. These numbers give the projection of the angular momentum on some axis in space. The states of given l and different values of m_l always have the same energy in any potential of spherical symmetry, even with potentials other than r^{-1} .

It is customary to designate the levels of different l by letters, in the following way:

$$l = 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5$$

$$s \quad p \quad d \quad f \quad g \quad h$$

Finally, the electrons have an intrinsic spin of $\frac{1}{2}$ about their own axes which can only have two directions in space. The direction of the spin can be

described by a quantum number m_s , with $m_s = \frac{1}{2}$ for spin "up" and $m_s = -\frac{1}{2}$ for spin "down." Thus every one of the $2l + 1$ states of given l is now double.

The basic assumption for the explanation of the periodic table is the following: In considering one particular electron, arbitrarily chosen, we shall assume that the action on it of all other electrons, as well as of the nucleus, can be approximated by a spherically symmetrical potential $V(r)$. Since this potential is no longer proportional to the reciprocal distance, the levels in it will be shifted, compared to hydrogen, and in such a way that the energy now depends on the angular momentum, measured by l , which is still quantized. The structure of the periodic system then follows from the Pauli principle: A quantum state of given n , l , m_l , m_s , can be occupied by only one electron.




Fig. 3. Energy release in alpha decays. [After Seaborg and Perlman]




Fig. 4. Energy levels in a square well.

In other words, an energy level characterized by l can be occupied by no more than $2(2l + 1)$ electrons. One builds up the periodic table by increasing the nuclear charge Ze , and with this the number of electrons, Z . To get the ground state of the atom we have to fill the lowest individual electron levels successively with as many electrons as the Pauli principle permits. When two successive levels are far apart

in energy, we speak of closing an atomic shell at the element for which the lower of these is filled. At the next element the next electron can only be brought into the atom at a much higher level, with much less binding energy.

This description of atomic structure may be termed the individual orbit model.

Individual Orbits in the Nucleus

In analogy with atomic structure one may postulate that in the nucleus the nucleons move fairly independently in individual orbits in an average potential which we assume to have spherical symmetry. The value of the angular momentum, l , is quantized and contains $(2l + 1)$ states, $-l \leq m_l \leq l$.

The occurrence of clear individual orbits of neutrons and protons in the nucleus is open to grave doubts. In the atom, there is firstly the dominant attraction of the nucleus. The Coulomb repulsion between the electrons is of long range, so that the potential acting on one electron does not depend sensitively on the precise position of the others. In the nucleus, on the other hand, the forces are of short range, so that the potential on one nucleon should depend strongly on the position of the others. In other words, one would expect that a nucleon would collide with another one long before it had traversed its orbit even once.

Actually, perturbation by collisions is not as severe as one would at first expect, since the Pauli principle forbids collisions that deflect nucleons into already filled orbits, and therefore most of the intuitively expected collisions do not occur. We shall pursue the description of the nucleus by the independent orbit model. It still remains surprising that the model works so well.

There are several differences between the nucleus and the electrons in the atom. Firstly, the average potentials in the two cases are quite different. Coulomb forces have a long range, in contrast to the nuclear forces. Thus, the atomic-shell numbers and the nuclear magic numbers will be entirely different from each other. One expects to find that the average nuclear potential has the form of a trough in three dimensions, where the potential is negative and rather constant inside the nucleus, rising abruptly to zero at the edge.

The second difference is that the nucleus contains two kinds of particles, neutrons and protons, each with intrinsic

spin $\frac{1}{2}$. We shall assume that the nuclear potential is the same for protons and neutrons. This assumption is now known to be in agreement with the evidence of many high-energy experiments, but at the time of the nuclear-shell-model development it was supported most strongly by the fact that the magic numbers were the same for neutrons and protons. The Pauli principle requires that, just as in the case of electrons, a level of given l , m_l , and m_s can be occupied by no more than $2(2l + 1)$ nucleons of one kind.

In a potential trough the lowest level is $1s$, $l = 0$, with room for two neutrons and two protons of one kind. Two protons and two neutrons in this level make He^4 . The next level is $1p$, $l = 1$, which has six states, so that the $1s$ and $1p$ levels together have room for eight nucleons of one kind. Since there are two kinds, neutrons and protons, altogether 16 nucleons can be accommodated, leading to O^{16} . Thus, the uniquely stable numbers are easily explained for the light nuclei.

This is by no means new, but is based on Wigner's (5) pioneering work on the light nuclei. Wigner's theory is able to explain with good approximation all the properties of light nuclei, spins, magnetic moments, transition probabilities, and so on.

Its natural extension, however, failed in predicting the properties of heavy nuclei, and somehow the theory of individual orbits in the nucleus went out of fashion. But nobody who has read Wigner's articles will ever forget them.

Figure 4 shows some types of average potentials, a square well in three dimensions, a well with rounded edges, and a three-dimensional harmonic oscillator. The three-dimensional oscillator has equally spaced levels, which are highly degenerate, but which split up into several levels of different angular momentum l in the square well. I shall frequently use the term *oscillator shell* by which I mean the group of levels which for the harmonic oscillator would have the same energy. All levels of one oscillator shell have the same parity—that is, they contain either only odd or only even values of l .

The right-hand side of Fig. 4 shows the order of the levels with different values of l , and the number of nucleons of each kind which fill these levels, in agreement with the Pauli principle.

The magic number 8 corresponds to filling all levels up to the oscillator shell $n = 1$. The magic number 20 is still explained as filling the oscillator shells




Fig. 5. Schematic level diagram.

up to $n = 2$. But beyond that the system breaks down. There is no trace of a gap in the level system at 28, 50, or 82.

The oscillator shells correspond to different numbers—namely, to 40, 70, and 112, which are not magic. Actually, for a potential which has not the oscillator shape but is a “square well” in character, the gap in energy at the oscillator shells is no longer marked.

Nuclear Shells

Elsasser had tried to explain the magic numbers by assuming that the nuclear potential in heavier nuclei is quite different from a square well. Subsequent work showed quite conclusively that a change in the shape of the potential, even a change which was quite unrealistic, could not explain the magic numbers. It was a kind of jigsaw puzzle. One had many of the pieces (not only the magic numbers), so that one saw a picture emerging. It seemed that if we had just one more piece everything would fit. The piece was found, and everything fell together.

At that time Enrico Fermi had become interested in the magic numbers. I had the great privilege of working with him, not only at the beginning, but also later. One day as Fermi was leaving my office he asked: “Is there any indication of spin-orbit coupling?” Only if one had lived with the data as long as I could one immediately answer: “Yes, and that will explain everything.” Fermi was skeptical, and left me with my numerology.

I do not know how many false starts my German colleagues made, but I had certainly made many. This was not one of them. The magic numbers from 28 on can definitely not be obtained by any reasonable extrapolation from the lower numbers, 2, 8, 20, but form a different sequence. There are two different series of numbers, 2, 8, 20, 40, . . . , of which 40 is no longer noticeable, and another, 6, 14, 28, 50, 82, 126, of which the first two, 6 and 14, are hardly noticeable. The second series is due to spin-orbit coupling. In 10 minutes the magic numbers were explained, and after a week, when I had written up the other consequences carefully, Fermi was no longer skeptical. He even taught it in his class in nuclear physics.

At about the same time Haxel, Jensen, and Suess had the same idea.

Let me explain what spin-orbit cou-

pling—or, more correctly, coupling of spin and orbital angular momentum—means. Earlier I have spoken somewhat vaguely about the quantum number of the intrinsic spin, m_s , which is $\frac{1}{2}$ for “spin up” and $-\frac{1}{2}$ for “spin down.” Up and down with respect to what? If one has just one nucleon in a shell, the only preferred direction is that of the orbital angular momentum. So spin, which is an angular momentum, can be parallel or antiparallel to the orbital angular momentum. The total angular momentum has then the magnitude $j = l + \frac{1}{2}$ or $j = l - \frac{1}{2}$. The number of states in each of two levels is $2j + 1$, due to differing orientation of the total angular momentum. There is no longer a factor 2, since the spin is now fixed. Notice that $[2(l + \frac{1}{2}) + 1] + 2(l - \frac{1}{2}) + 1 = 2(2l + 1)$, so that there are

still the same total number of states. I shall refer to the half integer j of a nucleon in a given state as its spin in this state.

The basic assumption of the shell model is that there is a strong interaction between spin and orbital angular momentum, giving the level $j = l + \frac{1}{2}$ a considerably lower energy. Since the splitting is proportional to l , and presumably goes down somewhat with nuclear size, prominent gaps in the level spectra will always occur when a high orbital angular momentum occurs for the first time. This explains the magic numbers. Let me show how this works for the number 28. The oscillator shell closes at 20. The next levels are $1f$ ($l = 3$) and $2p$ ($l = 1$), in that order. The $1f$ level splits into $j = 7/2$ and $j = 5/2$, with $7/2$ lower. Since the energy dif-




Fig. 6. Realistic level diagram for protons.

ference is large, and the $7/2$ level contains eight states, we find the gap at $20 + 8 = 28$ nucleons. All the magic numbers are explained in the same way. And since they are explained and no longer magic, I shall from here on call them shell numbers.

The assumption of a strong spin-orbit coupling contradicted the earlier tradition, which assumed that spin-orbit coupling was very weak. Our attitude was, "We know so little about nuclear forces." By now, there is ample evidence for the fact that spin-orbit interaction in nuclei is indeed an important effect. Figure 5 shows a very schematic level scheme. At the left are the numbers and levels of the oscillator shell. At the right is the level scheme with strong spin-orbit coupling. A magic number of neutrons or protons is obtained when the states of all oscillator shells up to a given one are filled with one each, and in addition the level of highest spin of the next oscillator is also filled with its complement of $2j + 1$ nucleons.

Figure 6 shows a fairly realistic level scheme for protons. It shows the fairly small splitting of the $1p$ or $l = 1$ level. The splittings of the $1f$ ($l = 3$) $1g$ ($l = 4$), and $1h$ ($l = 5$) levels are increasingly larger. Within the shells the level order is harder to predict. It depends on the relative strength of spin-orbit coupling and the deviation from the oscillator potential. The detailed order in which we put levels is dictated by experiment. For instance, in the shell with oscillator number 3 we find that the 29th proton, after the $7/2$ shell is filled, is in a $3/2$ orbit. So the level p ($l = 1$) $3/2$ is lower than the level f ($l = 3$) $5/2$, the partner to the $7/2$ state.

For neutrons, the level scheme is the same as for protons for the light nuclei up to neutron number 50. Above this, the Coulomb energy makes itself felt. It has the effect that the repulsion of protons favors orbits with higher angular momentum. Thus, for neutrons, for instance, the 51st neutron is in the d level of $j = 5/2$, whereas the proton is in the g level of $j = 7/2$. This effect is never large enough to affect the shell number.

Predictions of the Shell Model

To be a reasonable model of nuclear structure the shell model must be able to explain and predict nuclear properties

other than just a half dozen numbers. It is indeed able to do this.

Let me first consider the angular momenta, or nuclear spins, not of the individual nucleons but of the whole complex nuclei, which I shall designate by capital J . Hundreds of these have been measured. A closed shell, or a filled level, has angular momentum zero, since all states of different direction of the angular momentum contain one nucleon. Hence, nuclei with one nucleon outside (or one nucleon missing from) a closed shell of neutrons and of protons, or even of filled levels of both, should have a nuclear spin corresponding to the level of the single last nucleon, and the spin of the individual particle orbit is predicted by the shell model. This is quite a severe test, since we find there would be no possible way to explain a disagreement with the model. Happily, all known nuclei of this type have indeed the predicted spin and parity.

One example is ^{17}O , with one nucleon outside the doubly closed shell of ^{16}O , which has a spin of $5/2$ and positive parity, in agreement with the prediction. Another is ^{209}Bi , which has a nucleon outside the closed shells of 82 protons and 126 neutrons and has a spin of $9/2$, in agreement with the predictions.

In nuclei where both neutrons and protons fill shells incompletely, the individual nucleons add their spin vectors to a total spin vector $\mathbf{J}$. Even with the restriction of the Pauli principle, very many states of total angular momenta exist. For instance, if there are three identical nucleons in the $7/2$ shell there are six levels of different magnitude of one total angular momentum, ranging from $3/2$ to $15/2$. It is very fortunate that of the vast number of complicated levels only the simplest ones occur as the ground state of nuclei.

There are further regularities. For instance, in bismuth there exist five isotopes of odd mass number in which the neutron number is even. All have a nuclear spin measured to be $9/2$ —namely, that of the 83rd proton. Thus the even number of neutrons, ranging in this case from 116 to 126, does not influence the spin.

Another example is the region where the first $7/2$ shell is being filled. Here we know the spins of eight nuclei with an even number of protons and odd number of neutrons ranging from 21 to 27. Seven of these have nuclear spins $7/2$, one has spin $5/2$. There are also

five nuclei with an even number of neutrons and an odd number of protons ranging from 21 to 27, of which four have spins $7/2$, one has spin $5/2$. The numbers 21 to 27 correspond to 1, 3, 5, 7 nucleons in the $7/2$ shell. So, for nuclei in which both neutrons and protons fill shells incompletely, there emerge rules by which one may predict how the individual nucleons couple their spins to the total nuclear spin J . In a nucleus with an even number of neutrons and odd number of protons the neutrons couple their spins to zero and don't influence the nuclear angular momentum. The protons usually couple their spins to a total angular momentum J which is equal to the angular momentum j of the level being filled, and only rarely less by one unit. The same statement holds when the words *neutron* and *proton* are interchanged.

These rules are sometimes expressed in a different way. It is an experimental fact that all nuclei with an even number of neutrons and of protons have angular momentum zero. Thus, in a nucleus of even neutron number N , odd proton number Z , there is an even-even nuclear core with N neutrons and $Z - 1$ protons. The last proton occupies an orbit around the spinless core, and this orbit is prescribed by the shell model. All properties of nuclei—spin, magnetic moment, and so on—are entirely due to the last odd particle.

These coupling rules, considerably less complex than those for atoms, have some theoretical basis—namely, a simplified calculation of energies predicts them. If one considers just several particles of the same kind in the same level j , and assumes that they interact with each other with a very short range force, one finds indeed that for an even number of nucleons the ground state has spin zero. For an odd number, the ground state has the spin J which is equal to the j of the level being filled. The eigenfunctions of $J = 0$ for even, and of $J = j$ for odd, particle number are those of lowest seniority.

With these rules we should be able to explain or predict the spins of all nuclei. Up to neutron or proton number a little above 50 this simple theory and experiment are in excellent agreement. Beyond this, there are very many levels in the shells $Z = 50$ to $Z = 82$, and these levels lie close together in energy, so that one can explain just about anything. Besides, nuclei with more than 90 neutrons are highly deformed, and the assumption of a poten-

tial with spherical symmetry is no longer the best starting point. However, as the closed shells $Z = 82$ and $N = 126$ are approached, there is no longer a large deformation, and the predicted and measured spins again agree.

Another quantum number which the model predicts is the parity. We not only predict the spin but also the angular momentum l of each level. A level with odd l has odd parity, one with even l has even parity. Parity can be measured in various ways, and there is again complete agreement with the predictions.

Besides the ground states of nuclei one can also investigate the excited states. Excited states of one type are the isomeric levels, which are levels of a very long lifetime—hours, days, or even years. The explanation of this

phenomenon is that the spins of the isomeric state and the ground state are very different, so that the return to the ground state by the emission of a light quantum is greatly hindered, since the light quantum has to take up the difference in angular momentum. The transitions are not dipole but octupole or 2^4 -pole transitions, which are very slow. In nuclei of odd mass number an excited state can be produced by raising the last odd nucleon into an adjacent higher level. Now there are only very definite regions where low and high spins are close in energy—namely, at the end of the shells, where the lowest angular momenta of one oscillator shell occur, and immediately above them, where the states of highest angular momentum of the next oscillator level occur. Thus, isomerism should occur only

if the number of last odd particle is between 38 and 50, or between 64 and 82, or between 100 and 126. In addition, the shell model predicts that all these transitions involve a change in parity. This is a rather strong statement and ties isomerism to the neutron or proton number.

Some of the best work on isomerism has been done in Sweden (6) and has led to one of the nicest confirmations of the shell model. The three regions of isomerism are now called islands of isomerism. Long-lived and low-lying isomeric levels in nuclei of odd A occur only in the three islands. If one considers the mass number only, no regularities appear, since different islands of proton isomerism and neutron isomerism overlap in mass numbers.

For instance ^{115}In in the first island




Fig. 7. Magnetic moments of nuclei with odd proton, even neutron number.

has an isomeric state with a half-life of 5.1 hours. This is due to the transition of a proton from the $j = 1/2$ level to the ground state, which has spin $9/2$. For mass numbers higher by 2, one finds ${}_{50}\text{Sn}^{117}$ with an isomeric state of half-life 14 days. This is due to the odd neutron, which goes from the excited level $j = 11/2$ to a level $j = 3/2$, a transition which is expected to happen in the second island.

Failures of the Shell Model

After all this praise of the shell model, it is high time to emphasize its shortcomings. Even a crude nuclear model should be able to explain quantum numbers, like the spin, which is either integer or half integer but never in between, or parity, which is either even or odd. The shell model, as I have presented it, can indeed do this, and in this form has the advantage that it can explain or predict these quantum numbers for most nuclei.

However, the rules for the coupling of the spins of individual nuclei, which essentially postulate that everything depends only on the last odd nucleon, can be at best a very rough approximation to the truth. This becomes obvious when one tries to calculate nuclear properties which are not integers but can be measured to seven significant figures. One would hope to get approximate agreement, say to within 10 percent. Unfortunately, the agreement is not this good. For example, take the magnetic moments of nuclei. For a nucleus with odd proton, even neutron

number, the magnetic moments, according to the shell model, should depend only on the state of the last odd proton and are easy to compute. For any value of the spin, we calculate two different values of the magnetic moment, for the two different values of l , $l = j - 1/2$ and $l = j + 1/2$. In Fig. 7 the magnetic moments of odd proton, even neutron nuclei are plotted with the nuclear angular momentum as abscissa. The lines at the two extremes are the calculated ones. The middle lines are what one would obtain if the proton were a simple Dirac particle, and are added merely to emphasize the division into two groups. The difference between calculated and measured values is distressingly large. Only one general trend remains. The nuclei in the upper group, nearer to the line for $j = l + 1/2$, are indeed those for which we found that spin and orbital angular momentum are parallel, those in the lower group were assigned antiparallel orientation.

This shows that much more careful calculations of the interaction between the nucleons are required to get better numerical agreement. For individual nuclei, or special groups of nuclei, such calculations have been made by many people, using the shell model as first approximation and using different procedures to compute higher approximations. In particular, Talmi (7) has made great progress in developing a more refined shell model.

Finally, even the assumption of strong spin-orbit coupling is open to criticism, at least for the light nuclei. For these the model can easily be refined by taking into account both protons and neu-

trons in the nucleus. One should compare the results obtained to those of Wigner's (5) calculations. Although Wigner also used the independent particle model, his method is in some sense the direct antithesis of the shell model. In Wigner's theory, spin-orbit coupling is assumed to be very weak, whereas in the shell model spin and orbital angular momentum are rigidly coupled.

Actually, Wigner's values for all nuclear properties agree better with the experimental results for the light nuclei. It seems that the truth is in the middle: spin-orbit coupling is present, but not predominant. The calculations for "intermediate" coupling are more involved than either extreme, but they have been made by many people (8) for different nuclei, and have led to much closer agreement between theory and experiment.

The shell model has initiated a large field of research. It forms the starting point for more refined calculations. There are enough nuclei to investigate that the shell modelists will not soon be unemployed.

References

1. K. A. Brueckner, A. M. Lockett, M. Rotenberg, *Phys. Rev.* **121**, 255 (1961).
2. H. A. Bethe and J. Goldstone, *Proc. Roy. Soc. London* **A238**, 551 (1957).
3. W. Elsasser, *J. Phys. Radium* **4**, 549 (1933).
4. H. E. Suess and J. H. D. Jensen, *Arkiv Fysik* **3**, 577 (1951).
5. E. Feenberg and E. Wigner, *Phys. Rev.* **51**, 144 (1937).
6. E. Wigner, *ibid.*, p. 947. See, for instance, references in M. Goldhaber and A. W. Sunyar, *ibid.* **83**, 906 (1951).
7. A. de Shalit and I. Talmi, *Nuclear Shell Theory* (Academic Press, New York, 1963).
8. For references, see D. Kurath, in "Alpha, Beta and Gamma Ray Spectroscopy," K. Siegbahn, Ed., in preparation.