presented here on the Ivory Coast tektites do not clarify the significance of this isochron.

However, the plotted points for the Ivory Coast tektites lie essentially on the Birrimian-age isochron of 1.97 \times 109 years. Only three of the tektite points fall exactly on the isochron; the others all occur, at varying distances, slightly to the left of the line. This selective scatter to the left of the line can be ascribed to fractional volatilization of Rb relative to Sr during the fusion process which formed the tektites. Several recent studies suggest that selective volatilization of Rb relative to Sr took place during the fusion process that produced tektites (13) and, further, that the SiO₂ content decreases (hence the density increases) with increasing volatilization (14). The tektites listed in Table 1 (in order of increasing specific gravity) could be considered, therefore, as being listed roughly in order of increasing degree of volatilization and increasing loss of rubidium relative to strontium. There is an inverse correlation of rubidium with density, and a direct correlation of density with the distance of the points to the left of the line. That is, the samples at the top of the table, which according to the referenced studies should have suffered the least fractionation, lie on the line, while samples with increasing density (that is, less SiO₂ and more volatilization) fall, in general, further to the left of the isochron.

The age relationships presented here for the Ivory Coast tektites and the Birrimian rocks support the hypothesis of terrestrial origin for these tektites. The probability that these tektites, if extraterrestrial, would have just the right combination of rubidium, strontium, and strontium isotopic composition to lie on the isochron of the country rock on which they fell seems rather remote. Because much of the basement of West Africa is approximately 2×10^9 years old, this age value does not point unequivocally to the Bosumtwi crater as their place of formation. However, chemical similarities between the Ivory Coast tektites and glass from the Bosumtwi crater support this hypothesis of common origin. Gentner et al. (7) report the close similarity between the Bosumtwi crater glass and the Ivory Coast tektites for seven trace elements. Our Rb, Sr, and strontium isotopic analyses support this close similarity between the two types of materials. In addition, preliminary

18 FEBRUARY 1966

chemical analyses of these samples show a close similarity for the major elements examined: Fe, Ti, Ca, Mg, K, Na, and Mn (15). The concentrations of all these elements, except Ca, in the one piece of Bosumtwi glass available, lie within the rather restricted range of the 11 analyzed Ivory Coast tektites.

These three lines of evidence-the Rb-Sr age correlation (age of parent material), the K-Ar age correlation (age of fusion), and the chemical correlation-strongly support the hypothesis that the Ivory Coast tektites are related to the phenomenon that formed the Bosumtwi crater.

Note added in proof. An independent Rb-Sr isotopic study of three Ivory Coast tektites and five Bosumtwi crater glasses has recently been completed by Lippolt and Wasserburg (Z. Naturforsch., in press). Their data and conclusions are essentially in agreement with this paper.

C. C. SCHNETZLER Goddard Space Flight Center, Greenbelt, Maryland

> W. H. PINSON P. M. HURLEY

Department of Geology and Geophysics, Massachusetts Institute of Technology, Cambridge

References and Notes

- C. C. Schnetzler and W. H. Pinson, Geochim. Cosmochim. Acta 28, 953 (1964).
 M. Maclaren, Geog. J. London 78, 270 (1931).
 J. Littler, J. J. Fahey, R S. Deitz, E. C. T. C. T
- Chao, Geol. Soc. Am. Spec. Papers 68, 218
- (1961). (1961). H. P. T. Rohleder, *Centralbl. Mineral. Geol.* 316 (1934).

5. A. J. Cohen, J. Geophys. Res. 66, 2521 (1961).

- 6. J. Zahringer, Radioactive Dating (International Atomic Energy Agency, Vienna, 1963),

- tional Atomic Energy Agency, Manual Atomic Energy Agency, Manual Atomic Energy Agency, Manual P. 289.
 W. Gentner, H. J. Lippolt, O. Müller, Z. Naturforsch. 19a, 150 (1964).
 R. L. Fleischer, P. B. Price, R. M. Walker, Geochim. Cosmochim. Acta 29, 161 (1965).
 A. Lacroix, Compt. Rend. Acad. Sci. Paris 199, 1539 (1934).
 V. E. Barnes, Texas Univ. Publ. No. 3945, 477 (1939); E. C. T. Chao, in Tektites, J. O'Keefe, Ed. (Univ. of Chicago Press, Chicago. 1963), p. 51. O'Keefe, Ed. (Univ. of Chicago Press, Chicago, 1963), p. 51.
 N. R. Junner, Gold Coast Geol. Surv. Bull.
 8, 5 (1937); H. P. T. Rohleder, Geog. J. London 87, 51 (1936); John Saul, personal 11.
- communication.
- 12. M. Bonhomme, Ann. Fac. Sci. Clermont 5,
- M. Bonhomme, Ann. Fac. Sci. Clermont 5, Geol. Mineral. (1962); M. Vachette, Comp. Rend. Acad. Sci. Paris 258, 1569 (1964).
 J. A. Philpotts and W. H. Pinson, Geochim. Cosmochim Acta, in press; W. H. Pinson, C. C. Schnetzler, J. A. Philpotts, H. W. Fair-bairn, Trans. Amer. Geophys. Union 46, 118 (1965). Preliminary work on heating of tek-tites to 1800° to 2000°C, in both air and vacuum, indicates loss of Rb of between 15 and 35 percent while the Sr. ramains account. percent, while the Sr remains essenand 35 tially constant.
- tally constant.
 14. J. A. Philpotts and W. H. Pinson, Geochim. Cosmochim. Acta, in press; L. S. Walter and M. K. Carron, *ibid.* 28, 937 (1964); L. S. Walter, Science 147, 1029 (1965).
 15. W. H. Pinson, J. A. Philpotts, J. Hwang, un-published data on analyses by x-ray fluores-cence and actomic abservation
- cence and atomic absorption.
- We are indebted to the following for donating 16. samples: John Saul and Elliot Miller, of Saul Miller, Consulting Geologists; E. C. T. Chao of the U.S.G.S.; R. M. Walker and R. Laboratory; and the Société de Développe-ment Minéral de la Côte d'Ivoire. Financial aid for this research was furnished in part by NASA through the Center for Space Re-search at the Massachusetts Institute of Tech-pology and in part by the U.S. dtout Fichard nology and in part by the U.S. Atomic Energy Commission. Field work for the collection of the tektites in the Ivory Coast by John Saul and Elliott Miller was sponsored by the Na-tional Geographic Society. We also thank W. R. Shields of the National Bureau of Standards for the use of mass spectrometers, H. H. Thomas, M. L. Bottino, and P. D. Ful-lagar for aid in analytical work, and members the GSFC Geochemistry Laboratory for helpful discussions.
- 18 October 1965

Imidonitrogen in Chlorella "Polyphosphate"

Abstract. "Polyphosphate" and fragments isolated from acid hydrolysis of polyphosphate have an infrared absorption band at 1400 cm⁻¹, which is characteristic of imidodiphosphate linkages. Complete hydrolysis of purified "polyphosphate" releases 1 to 2 moles of phosphate per mole of ammonia. The polymer must contain subunits which are cyclic and which contain both imidodiphosphate linkages and phosphate anhydride linkages.

The polyphosphate in Chlorella is not a simple phosphate anhydride polymer (1-3); nor is it, because of its stability in water, one of the many polymers containing phosphorus which are unstable in water. The properties of trimetaphosphoimidate (4) prompted a study to determine whether metaphosphoimidate groups occur within the material of high molecular weight commonly designated as polyphosphate. Imidodiphosphate linkages have not been reported in natural products.

Sodium trimetaphosphoimidate was synthesized (5) in order to study its properties in more detail. Its specific metachromasy (2) is 0.3 compared to 0.0 for trimetaphosphate. Thus metaphosphoimidates are more metachromatic than metaphosphates. This is significant because Chlorella polyphosphates have been reported to have very high specific metachromasies, which could not be explained if the polyphosphate contained only anhydride linkages (2). Trimetaphosphoimidate released 30 percent of its phosphorus as orthophosphate when treated for 7 minutes with 1N hydrochloric acid at 100°C. (This release may be abbreviated as follows: $\Delta_7 P = 30$ percent.) A phosphate anhydride is completely hydrolyzed under these conditions ($\Delta_7 P$ = 100 percent). Purified *Chlore!la* polyphosphate is intermediate ($\Delta_7 P = 85$ percent) (3). When trimetaphosphoimidate is refluxed for 1 hour in 18N sulfuric acid, it releases all of its nitrogen as ammonia and all of its phosphorus as orthophosphate.

Trimetaphosphoimidate was hydrolyzed in 1N HClO₄ (15 minutes at 100°C), neutralized with KOH, and chromatographed on a column containing Dowex-1-formate resin (3). Four fractions were resolved. The first fraction (61 percent of the total phosphorus applied to the column) was orthophosphate. The second fraction (19 percent of the total phosphorus) was eluted in the diphosphate position and was a mixture of pyrophosphate and imidodiphosphate ($\Delta_7 P = 69$ percent). The third fraction (8 percent of the total phosphorus) was eluted in the triphosphate position and was mostly imidotriphosphate, a known (4) major intermediate of acid hydrolysis ($\Delta_7 P = 50$ percent). The fourth fraction (12 percent of the total phosphorus) was either unaltered trimetaphosphoimidate or, more likely, the product formed by the substitution of one trimetaphosphoimidate nitrogen by oxygen ($\Delta_7 P = 10$ percent).

Trimetaphosphoimidate was also hydrolyzed in 0.5N KOH (18 hours at 38°C) neutralized with HClO₄, and chromatographed on a column containing Dowex-1-formate resin in the same manner. Only three fractions were resolved. The first fraction (4 percent of the total phosphorus applied to the column) was orthophosphate. The second fraction (60 percent of the total phosphorus) was eluted in the diphosphate region and was mostly imidodiphosphate ($\Delta_7 P = 10$ percent). The third fraction (36 percent of the total phosphorus) was eluted in the triphosphate region and was mostly imidotriphosphate ($\Delta_7 P = 50$ percent).

Thus hydrolysis in 1N acid at 100° C for 15 minutes of trimetaphosphoimidate releases orthosphosphate and diand triphosphates which are more resistant to acid hydrolysis than pyro- and tripolyphosphate. Under these conditions *Chlorella* polyphosphate releases 95 percent orthophosphate and 5 percent di- and triphosphates which are



Fig. 1. Some possible repeating units in the structure of *Chlorella* "polyphosphate."

relatively acid resistant (3). Alkaline hydrolyzates of both trimetaphosphoimidate and *Chlorella* polyphosphate contain only a few percent of orthophosphate and a large amount of diand triphosphates. In the case of polyphosphate these are predominantly pyro- and tripolyphosphate. In the case of trimetaphosphoimidate they are predominantly imidodiphosphate and imidotriphosphate.

Chlorella polyphosphate was treated with venom phosphodiesterase and it was then chromatographed on a column of diethylaminoethylcellulose resin to remove the polyphosphate from the RNA components. These polyphosphate fractions and the di- and triphosphates isolated from acid hydrolyzates of polyphosphate by column chromatography on Dowex-1-formate resin were analyzed on a Perkin-Elmer infrared spectrophotometer, model 521. Samples were prepared by the potassium bromide pellet technique (2). Sodium trimetaphosphoimidate; hydrolytic fractions obtained from trimetaphosphoimidate such as imidodiphosphate and imidotriphosphate; ortho-, pyro- and tripolyphosphate; yeast RNA; Graham's salt, a synthetic true polyphosphate (5); and reagent blanks were analyzed in the same way. All compounds which contained imidodiphosphate linkages, and only these compounds, exhibited an absorption at 1400 cm^{-1} . Intact Chlorella polyphosphate and the acidresistant di- and triphosphates derived from it also had an absorption band at 1400 cm^{-1} .

Purified fractions of Chlorella poly-

phosphate were also assayed for ammonia nitrogen which was released by refluxing one hour in 18N sulfuric acid. The released ammonia was determined by a modified Nessler reaction (6) after micro-Kjeldahl distillation. The ratio of moles of phosphate per mole of ammonia released varied from one to two in a variety of fractions tested.

Thus Chlorella polyphosphate, as isolated, contains both phosphate anhydride and imidodiphosphate bonding. However, oxygen is easily substituted for nitrogen in polyphosphoimidates yielding mixed bonding or even true polyphosphates. A certain amount of this probably happens during isolation since this replacement takes place in water from pH 1 to 14 (4, 7). Also, cyclic groups or at least branching must occur in the polymer, since both linear polyphosphoimidates and polyphosphates are stable to alkaline hydrolysis (5, 7). In contrast cyclic (meta) phosphate anhydrides or phosphoimidates are cleaved at alkaline pH(4). The phosphorus atoms at which branching occurs must have at least one imidolinkage, since a phosphate trianhydride is immediately hydrolyzed in the presence of water (5). Not enough is known about the stabilities of other potential branching combinations to make further conclusions. Some possible repeating units in the structure of Chlorella polyphosphate are shown in Fig. 1. In such a polymer the ratio of phosphorus to nitrogen would be 1.5. Variations, which are not excluded by my data, include eight-membered rings, higher or lower ratios of phosphorus to nitrogen, the presence of mono-, di-, or triphosphate connections between rings, or branching brought about by connecting more than two rings to some rings. Upon alkaline hydrolysis such polymers might be expected to cleave within each repeating unit to open the ring and between units to yield low-molecularweight products, but little orthophosphate. The composition of the hydrolyzate would depend upon slight, but as yet unknown, differences in the stability of the bonds in each unit. Acid hydrolysis of such polymers would cleave the anhydride linkages quickly and would carry out two competitive reactions at the imidodiphosphate sites, substitution of oxygen for nitrogen or cleavage (4). Anhydride bonds produced by oxygen substitutions would then be cleaved. Any amido groups produced by cleavage would be quickly hydrolyzed in acid to give ammonia. Thus hydrolysis would be expected to yield a large amount of orthophosphate and small amounts of imidodiphosphate and imidotriphosphate. Polymers of the type shown in Fig. 1 would be highly metachromatic due to the presence of few secondary phosphate ionizations, which inhibit metachromasy (8). They might also exhibit a completely different affinity for RNA than a true polyphosphate, thus resembling the naturally occurring "polyphosphate."

DAVID L. CORRELL Radiation Biology Laboratory, Smithsonian Institution, Washington

Hemagglutination by Fava Bean

of hemagglutination by fava bean.

Agglutination of normal human

erythrocytes by substances other than

blood group-specific isoantibodies is

well known (1). Inhibition of this

hemagglutination by various complex

sugars has been reported (2). The ef-

fects of simple sugars on hemagglutina-

tion by saline extract of fava bean are

d-fructose, or maltose, but not by

5-percent d-galactose or lactose (Ta-

hydroxyl group prevents inhibition of

Hemagglutination by fava bean was

5-percent d-glucose,

the subject of this report.

inhibited by

Extract Inhibited by Simple Sugars

References and Notes

- 1. D. L. Correll and N. E. Tolbert, Plant Physiol. 37, 627 (1962).
- -, Plant Cell Physiol. 5, 171 (1964). 2
- D. L. Correll. *ibid.*, in press.
 A. Narath, F. H. Lohman, O. T. Quimby,
- A. Ivataui, F. H. Lonman, O. T. Quimby, J. Amer. Chem. Soc. 78, 4493 (1956).
 J. R. Van Wazer, Phosphorus and its Com-pounds (Interscience Publishers, New York, 1958), vol. 1.
- 6. H. S. Faber, W. D. Hatfield, M. H. McCrady, Eds., Standard Methods for the Examination of Water, Sewage, and Industrial Wastes (Wav-erly Press, Baltimore, 1955), p. 144.
- V. R. Klement and G. Biberacher, Z. Anorg. Allgem. Chem. 285, 74 (1956). 7.
- K. K. Tewari and P. S. Krishnan, Arch. Bio-chem. Biophys. 82, 99 (1959). 9. Supported in part by a grant from J. B.
- Bredin. 17 November 1965

enhanced by gum acacia (3). Similar studies with dextrans of varying molecular weights (5) show augmented titers (1:640) with dextran of high molecular weight (75,000), but not with dextran of lower molecular weight (41,500). The hypothesis that enhancement of agglutination by gum acacia is due to the gum's molecular size is strengthened. Agglutination by dextrans of high molecular weight also is inhibited by 5-percent d-glucose or normal human serum.

CONSUELA B. PERERA

Abraham M. Frumin Departments of Laboratories and Research, Albert Einstein Medical Center, Southern Division, Philadelphia, Pennsylvania 19147

References and Notes

- W. C. Boyd, Vox Sanguinis 8 (1963).
 W. M. Watkins and W. T. J. Morgan, Nature 169, 825 (1952); 175, 676 (1955); H. M. Johnson, Science 146, 548 (1964); L. D. Gelb and A. M. Lerner, *ibid.* 147, 404 (1965).
 W. P. Creger and H. Gifford, Blood 7, 721 (1952)
- (1952) K. L. Roth and A. M. Frumin, J. Lab. Clin. Med. 56, 695 (1960).
 Microdextran obtained from Pharmachem
- Pharmachem Corp., Bethlehem, Pa.; macrodextran, from Abbott Laboratories, North Chicago, Ill. Preparation of bean extract and hemagglutination techniques accorded with Creger and Gifford's (3). Supported by a grant from the Wagman Foundation.

18 August 1965

Encephalitogenic Activity of Bovine Basic Proteins

Abstract. Two basic proteins isolated from bovine white matter in connection with a study of the protein-bound phosphoinositides of central nervous system tissue have been tested for encephalitogenic activity. The biological activity of these proteins, which is equivalent to that of basic encephalitogenic proteins isolated in other laboratories, suggested that they are identical.

Table 1. Effects of simple sugars on hemagglutination by saline extract of fava bean. Degree of agglutination indicated by number of plus signs.

Abstract. Hemagglutination by extract of fava bean was inhibited by 5-percent d-glucose, d-fructose, or maltose, but not by 5-percent d-galactose or lactose. Failure to inhibit seems to reflect the presence of a hydroxyl group at the carbon

No. 4 position. Hemagglutination was enhanced by dextran of high molecular

weight, but not by dextran of low molecular weight. The finding supports the

hypothesis that large molecular size explains the enhancement by gum acacia

Sugar concentration (%)		Extract titer				
		1:10	1:20	1:40	1:80	1:160
None		++++	+ + + +	+ + +	+ +	0
Lactose	5	++++	+ + + +	+ + +	+ +	0
d-Galactose	5	++++	+ + + +	+ + +	+ +	0
Maltose	5	0	0	0	0	0
d-Fructose	5	0	0	0	0	. 0
d-Glucose	5	0	0	0	0	0
d-Glucose	0.6	0	0	0	0	0
d-Glucose	.55	+ +	+ +	0	0	0
d-Glucose	.4	+ + +	+ + +	+ +	+ +	0

18 FEBRUARY 1966

favism (4). These observations suggest ble 1). The determining factor appears to reside in the configuration of the that induced hyperglycemia or trans-No. 4 carbon atom: presence of a fusion of plasma may have a thera-

peutic effect in human favism.

human serum of hemagglutination by

fava bean has been reported (3); the

concentration of glucose in normal hu-

man serum, 0.6 to 1.05 mg ml⁻¹, can-

not explain such inhibition. Inhibition was complete with 0.6-, partial with

0.55-, and none with 0.4-percent d-glu-

cose. Agglutination disappeared if nor-

mal serum or 5-percent d-glucose was

added after agglutination occurred. The

inhibition in vitro by normal serum of

hemagglutination by fava bean did not

occur in the serum of a child with

hemagglutination. Inhibition by normal Hemagglutination by fava bean is

> The basic protein fraction of central nervous system (CNS) myelin has been investigated intensively because it can induce experimental allergic encephalomyelitis (EAE) in various animals (1). Although the encephalitogen is part of the myelin proteolipid, it is not found in the chloroform-methanol (2:1 mixture) extract of the whole tissue. Its absence from the chloroformmethanol extract of whole tissue is presumably caused by some interaction between the encephalitogenic proteolipid and other constituents of the tissue