

of estradiol fused on stainless steel rods. One of these was placed in the pectoral region, the other under the scruff of the neck. This procedure, we find, produces a female whose receptivity remains constant over many months (6). The rats were maintained in cages with large-mesh flooring (12 mm) so all droppings would immediately fall through onto a paper-covered tray below. Every morning between 8:30 and 10:00 a.m. a record was made of the number of copulation plugs lying on the paper. These data were used to calculate the number of copulations and the percentage of days on which copulations occurred.

Following a 2-week control period lesions were made with platinum electrodes 0.36 mm in diameter (28 gauge), with the insulation cleared from about 0.5 mm at the tip. With the Grass S-4 stimulator a direct current of 1 ma (anodal) was applied for 15 seconds; the cathode was a large metal rod placed in the anus. The coordinates used were those of the de Groot (7) atlas, with the middle hypothalamic lesions made at A 6.0 and 5.6, midline and depth -3.0 mm. The mammillary lesions were made at A 4.2, midline and depth -4.0 mm; for the lateral lesions the same coordinates were used, except that they were 1.5 mm on either side of the midline. All lateral lesions were made bilaterally.

These findings clearly do not support the workers who found cessation of sexual activity following mammillary lesions. They are in agreement with those who found no change in sexual behavior following mammillary lesions. One report has appeared which indicated that fairly large lesions at the junction of the diencephalon and mesencephalon can result in changes in the behavior pattern making up a complete copulatory sequence (8). Following such lesions about 50 percent of the animals showed a decreased number of intromissions to ejaculation and a shortened latency period between an ejaculation and the beginning of the next copulatory sequence. Thus in a 30-minute test period the animals ejaculated a greater number of times than the controls. These changes were found to be stable during 2 to 6 months of testing. Similarly, for the female there is a report that lesions in the pre-mammillary region resulted in five of six animals mating at diestrus (9).

In the two studies reported above iron-bearing electrodes were used; thus

they might be open to interpretation from the standpoint that the results were dependent upon irritative foci set up by deposits of metal salts in the brain, especially in light of the fact that Everett and Radford got ovulation only when stimuli were passed through iron-bearing electrodes, and then only when a certain mass of nervous tissue had been affected by the iron deposits. Similar stimulation with platinum electrodes failed to produce ovulation (4).

Our study was made with platinum electrodes to obviate the possible problems inherent due to the possibilities of chronic stimulation resulting from metal deposits in the brain. Using the platinum electrode, we were able to produce striking changes in total sex behavior in the male as the result of placing small lesions in the mammillary bodies.

Therefore inhibitory structures important in regulation of the occurrence and frequency of sexual behavior must be assumed to exist in the mammillary body region in the male rat. In the male, lesions in the mammillary region lead to increased total copulatory performance, while in the female lesions in the mammillary region result in a release of overt sex behavior from the controlling influence of the hormonal milieu as expressed in the normal estrous cycle.

ROBERT D. LISK

*Department of Biology,
Princeton University,
Princeton, New Jersey 08540*

References and Notes

1. J. M. Brookhart and F. L. Dey, *Amer. J. Physiol.* **133**, 551 (1941); K. Larsson and L. Heimer, *Nature* **202**, 413 (1964).
2. M. L. Soulaire, *Ann. Endocrinol.* **24**, No. 3 Suppl. (1963).
3. L. Heimer and K. Larsson, *Acta. Neurol. Scand.* **40**, 4 (1964).
4. J. W. Everett and H. M. Radford, *Proc. Soc. Exp. Biol. Med.* **108**, 604 (1961); R. W. Reynolds, *Amer. J. Physiol.* **204**, 60 (1963).
5. Rats obtained from Carworth Farms.
6. Constant behavioral estrus can be maintained over long time intervals in the female rat via appropriate implants of a solid ball of estradiol fused to the end of 27-gauge hypodermic tubing. In some experiments a single estradiol implant was made via stereotaxic methods, the hormone being placed in the anterior hypothalamus. Subsequently we found two such implants placed subcutaneously were equally effective in maintaining the female in a constantly receptive state over a few months. Castration was not performed before placement of the estradiol implants.
7. J. de Groot, *J. Comp. Neurol.* **113**, 389 (1959).
8. L. Heimer and K. Larsson, *Experientia* **20**, 460 (1964).
9. T. Law and W. Meagher, *Science* **128**, 1624 (1958).
10. I thank Alice Suydam and Linden Vargish for excellent technical assistance. Study supported by NSF grant GB 914.

2 December 1965

Neutron-Activation Analysis

Schroeder, Kraner, Evans, and Brydges [*Science* **151**, 815 (1966)] describe the use of lithium-drifted germanium detectors for neutron-activation analysis when destruction of the specimens must be avoided. I wish to suggest a rather obvious modification of the technique, not mentioned by the authors or elsewhere so far as I am aware.

The results reported pertain to irradiation by the whole neutron spectrum of the particular reactor used, in this case primarily a thermal neutron spectrum with a higher-energy tail. It is noted that the presence of Na may make observations of elements with similar half-life difficult, but that, in general, the data from the presence of elements with dissimilar half-lives can be enhanced or suppressed by choice of irradiation time. What is not mentioned is that this can also be done by spectral modification of the neutron flux. The modification consists in suppressing the number of neutrons in a given energy region of the spectrum—the region responsible for the activation. This can be done by surrounding the specimen by a suitable absorber of "Dagwood sandwich" of absorbers or by making a suitable geometry for the use of materials which selectively scatter neutrons away from the sample so that those impinging on it have the desired spectrum. The first method uses only the dependence of absorption cross section on neutron energy and the second uses the total cross section. The simplest arrangement is to use the element which it is desired to suppress. However, there are so many resonances among the many stable nuclides, and overlapping ones, that a large number of possibilities exists. Sometimes one may wish to employ a different primary spectrum by using a different reactor.

The suppression of Na is probably not the easiest case, though certainly some gain can be achieved by a scattering-out arrangement, for Na is a weak absorber. Undoubtedly, the bulk of activation occurs in the $1/v$ region for practically all nuclei, so one would wish strong filtering in this region with ^{10}B or ^3He in order to take advantage of the more selective spectral modification of the resonance region. Whenever the activation results from resonance capture, it should be possible to suppress it almost wholly by filter-

ing the spectrum. A reactor with a nonthermal spectrum would be advantageous.

Schroeder *et al.* remarked that lead is not readily measurable by activation analysis and hence they were not able to test its presence in the old printed matter they analyzed. Silver is a common impurity in commercial lead even today. Perhaps the silver content of old lead objects could provide a clue in correlation with the appearance of silver in printed matter.

J. H. MANLEY

Los Alamos Scientific Laboratory,
Los Alamos, New Mexico 87544

10 March 1966

Statistical Analysis in Toxicity Experiments

Three points should be made about Lindsay and Kullman's "Pentobarbital sodium: variation in toxicity" [*Science* **151**, 576 (1966)].

First, their statistical analysis gives no appreciable basis for the notion that dependence of toxicity upon hour of the day varies from experiment to experiment. Only one of eight tests reached the conventional 5-percent significance level. This should be expected by chance in 34 percent of such analyses, and in 14 percent of such analyses with the actual level reached, 0.019. Further, the comprehensive test of this interaction reaches only the probability of 0.092. If this slight observed tendency toward significant interaction needs explanation, one is readily available in that the arcsin transformation would not preserve the shape of the curve as dose (or sensitivity) changes.

Second, there is appreciable evidence that time of day has *some* effect, which is apparently roughly quadratic, with a maximum near the middle of the interval studied. (The "significant" term of seventh degree is subject to discussion parallel to that above.) Thus the concluding paragraph is misleading at best: no substantial evidence has been presented that the "full complexity of the toxicity curve" is more than that of the diurnal fluctuation which is familiar in many phenomena.

Third, it is a serious deficiency in the design of these experiments that "Two cages were taken in sequence for the experiment at each time period." Any observed effect might have been a conse-

quence of differences between cages, and it cannot be argued logically that the effect is even partly ascribable to time of day, unless this possibility is taken into account in the statistical analysis and interpretation.

H. W. NORTON

Department of Animal Science,
College of Agriculture,
University of Illinois, Urbana

11 February 1966

Quasar Red Shifts: Can They Be Due to Implosion?

Halton Arp [*Science* **151**, 1214 (1966)] has given significant observational evidence for believing that at least some of the quasars are not as distant as has been generally supposed. If his evidence can properly be extrapolated to embrace all the quasars, their power output will no longer pose a formidable problem. But, as he points out, a new problem will arise because we shall no longer be able to attribute their very large red shifts to extreme distance.

As one possibility, he has suggested that the red shifts may be due to high collapse velocity of material toward the centers of the quasars regarded as very compact objects. It is the purpose of this note to mention two possible difficulties with this suggestion. The first is suggestive rather than compelling. It has to do with the size and longevity of the quasars. Fluctuations in brightness place an upper limit on the size of at least some of them. If their outer radiating matter is falling inward with radial speeds that are significant fractions of the speed of light, and if the these speeds are not ephemeral, then the quasars would have to have been much larger in the past. For example, 3C273 has been traced back to the year 1888, and it exhibits fluctuations in brightness, one period being of length 15 years and the smallest being of length of about a year. If we assume that the red shift corresponding to a recession velocity of approximately $0.15c$ has been substantially constant since 1888, and if we assume that the red shift is due to the implosion suggested by Arp, then 3C273 will have shrunk in diameter by about an order of magnitude, and one has to ask whether it is likely that the periodicity of the fluctuations in brightness would have been maintained during so great a shrinkage in size.

The second difficulty is more precise.

Think of a quasar as an imploding sphere presenting a visual appearance of an imploding disk. The light from the center of the observed disk will indeed exhibit the full Doppler red shift, but light from other parts of the disk will exhibit smaller shifts, since the luminous matter remitting the light will be moving toward the center of the sphere and thus will not be moving away from us along the line of sight. Indeed, light from the edge of the disk will come from matter moving transversely to the line of sight and will exhibit at most only the small second order Doppler effect of $\frac{1}{2}v^2/c^2$ (plus such gravitational and Hubble shifts as may be present, but these, on the present hypothesis, are small compared with the observed shift). Consequently, there would be not a red *shift* but rather a red *smear*. The spectral lines would not be displaced while remaining relatively sharp but would be spread out, stretching from their gravitational-plus-Hubble-shifted positions. Photos of the shifted lines do not seem to exhibit such a broadening.

If there is merit in either of the foregoing points, it would seem that the red shifts of the quasars cannot be attributed to the implosion process tentatively put forward by Arp, and this leaves only his other alternative of an unknown process.

Arp does remark that fairly narrow spectral lines could result if only parts of the imploding shell were luminous at a given time, though he does not specifically link this to the foregoing Doppler-shift argument. Since, according to that argument, different parts of the shell should exhibit shifts ranging over a very broad interval, the existence of two localized luminous regions at different distances from the center of the disk would result in a doubling of the spectral lines, and this effect has not been reported.

The situation thus remains as puzzling as before, with only the location of the puzzle having changed. If we take the red shifts to be proof of extreme distance, we are confronted with the problem of their power output and also of their visual relation to the peculiar galaxies. If we accept Arp's results as evidence for the relative nearness of the quasars, we are confronted by the problem of the red shifts themselves.

BANESH HOFFMANN

Department of Mathematics
Queens College, City University of
New York, Flushing 11358

15 March 1966