

for intermediate rocks against the observations, it can be shown that intermediate rocks are not excluded by their observations. In addition, as the authors themselves have emphasized, the necessity of allowing for background corrections, which are very uncertain and which account for 90 percent of the measured values, makes it somewhat difficult to draw firm conclusions from this pioneering effort.

JOHN A. O'KEEFE

PAUL D. LOWMAN, JR.

WINIFRED S. CAMERON

Goddard Space Flight Center,
Greenbelt, Maryland

References and Notes

1. H. Jeffreys, *The Earth* (Cambridge Univ. Press, Cambridge, ed. 3, 1952), p. 352; J. Green, *Proc. Lunar and Planetary Exploration Colloquium* (North American Aviation, Downey, Calif.), 1, 1 (1959); Douglas Paper 4038 (Douglas Advanced Research Laboratory, Huntington Beach, Calif., 1966); G. Fielder, *Lunar Geology* (Lutterworth, London,

- 1965), chap. 9; G. J. H. McCall, *Ann. N.Y. Acad. Sci.* 123, 843 (1965); W. E. Elston, *ibid.* p. 817.
2. R. L. Smith, *U.S. Geol. Surv. Prof. Papers* 424D, 145 (1961).
3. C. A. Cotton, *Volcanoes as Landscape Forms* (Whitcombe and Tombs, Christchurch, N.Z., 1952), chap. 11.
4. H. Jeffreys, *Phil. Mag. Ser. 6* 49, 793 (1925).
5. R. H. Nichols, *J. Geol.* 47, 290 (1939).
6. E. A. Whitaker, *Science* 153, 1550 (1966).
7. C. K. Wentworth, M. H. Carson, R. H. Finch, *J. Geol.* 53, 94 (1945).
8. E. E. Wahlstrom, *Introduction to Theoretical Igneous Petrology* (Wiley, New York, 1950).
9. C. Wahrhaftig and A. Cox, *Bull. Geol. Soc. Amer.* 70, 383 (1959).
10. R. Baldwin, *The Measure of the Moon* (Univ. of Chicago Press, Chicago, 1963), p. 248.
11. A. K. Lobeck, *Geomorphology* (McGraw-Hill, New York, 1939), p. 81.
12. A. E. Scheidegger, *Theoretical Geomorphology* (Prentice-Hall, Englewood Cliffs, N.J., 1961), p. 72.
13. O. D. von Engel, *Geomorphology* (Macmillan, New York, 1942), p. 245.
14. A. P. Vinogradov, Iu. A. Surkov, G. M. Chernov, S. S. Krinozov, *Geokhimiya* 8, 891 (1966).
15. We thank the Lunar Orbiter Project Office at Langley Research Center, Hampton, Va., and Dr. M. J. Swetnick for the pictures.

22 November 1966

Pressure-Induced Phase of Sulfur-Selenium

Abstract. *Crystals of a fibrous phase of sulfur-selenium obtained at 20 kilobars and 280°C are trigonal, the most probable space groups being $P3_1$ and $P3_2$, with $a = 7.85$, $c = 4.62 \pm 0.01$ Å. The unit cell contains nine atoms, and the measured density of 3.20 g/cm³ implies five sulfur and four Se atoms. The structure contains mixed atom helices of 1.54 Å pitch and 0.91 Å average radius.*

In a continuing investigation of group VIA elements a new pressure-induced sulfur-selenium phase has been found. The phase is fibrous but is not isostructural with the fibrous sulfur phase (II) (1). In fact, we have also found that some selenium does dissolve in the fibrous sulfur phase.

Starting materials were pure (99.999+ percent) Se and S (American Smelting and Refining Company). A one-to-one mixture (atom percent) was put into a fused silica tube, evacuated, and sealed. The mixture was melted and kept at 250°C for 2 hours and annealed at 80°C for 110 hours. It was then removed from the tube and ground and mixed thoroughly in an attempt to insure homogenization. Some of this material was then packed into tantalum containers and subjected to pressure and heating in furnaces and piston cylinder devices similar to those described by others (1, ref. 1). The fibrous S-Se phase reported here was prepared in a furnace (2.54 cm diameter) at 20 kb. The temperature was raised to 550°C and held there for 10 minutes; the temperature was then reduced to 280°C and maintained

there for 56 hours. The resulting material was not homogeneous but that part of the sample in the bottom portion of the sample capsule was reddish-brown, crystalline, and fibrous. A measurement of the density of isolated crystals of the fibrous form by the flotation technique gave 3.20 g/cm³.

An apparently single crystal of the fibrous sulfur-selenium was aligned along the fiber axis with oscillation photography; $\text{CuK}\alpha$ radiation was used, and Weissenberg photographs were taken. Lattice constants were determined from Buerger precession camera photographs ($\text{MoK}\alpha$ radiation). The diffraction symmetry of all the photographs is $6/m$, the only systematic absences being those reflections ($00l$) for which l is not equal to $6n$. The lattice constants of the particular crystal photographed are: $a = 7.85$, and $c = 4.62 \pm 0.01$ Å. Hexagonal selenium has $a = 4.355$, $c = 4.949$ Å. The sublattice obtained by a 30° rotation from the unit cell of the sulfur-selenium phase has lattice constants $a = 4.53$ Å and $c = 4.62$ Å. It appears then that the sulfur-selenium unit cell must contain nine atoms. A cell content of five S and

four Se atoms gives an x-ray density of 3.20 g/cm³ equal to the measured density.

Any space group giving diffraction symmetry $6/m$ satisfying the conditions for these helices must contain screw axes. Further, because of the length of the c -axis, the helices in the sulfur-selenium phase must have three atoms per turn as in hexagonal selenium itself. No hexagonal space group giving diffraction symmetry $6/m$ can satisfy the requirements for this structure. Thus it appears that the $6/m$ is only an apparent diffraction symmetry; the more probable diffraction symmetry is $\bar{3}$. When crystals with this symmetry are 120° rotation-twinned, they give the apparent symmetry observed. This is analogous to the case of selenium itself (2) in which the twinning of crystals with diffraction symmetry $\bar{3}m$ leads to apparent symmetry $6/mmm$.

Thus the most probable space groups to which the fibrous $\text{S}_{0.555}\text{Se}_{0.444}$ belongs are $P3_1$ or $P3_2$. It is possible also that the two enantiomorphs are cocrystallizing in the twinned crystals.

Thus far the preliminary refinement of the x and y parameters with the use of the Busing-Martin-Levy (3) program (modified for use on the IBM 360 computer) and only the $hk0$ intensity data (for which there is no overlapping of nonequivalent reflections) indicates that the helix radius is close to 0.91 Å; the pitch, given by $c/3$ is 1.54 Å. This implies an average S-Se distance (4) of 2.20 Å as compared with a calculated one of 2.18 Å based on a value of 2.34 Å for an Se-Se distance and 2.05 Å for an S-S distance.

For Se, the pitch and radius of the helix are 1.65 and 0.95 Å, respectively (4). Thus the larger a -axis of the subcell (see above) implies poorer packing efficiency of the sulfur-selenium phase than of the hexagonal Se phase.

Spacings were calculated with the lattice constants determined from the Buerger precession camera photographs. It is seen in Table 1 that the calculated spacings compare well with those measured on an x-ray powder photograph of the material. All nonequivalent sets of indices are given.

There appears to be a range of solid solutions having the same fibrous structure, but the limits have not yet been determined. The new phase is not nearly as stable as the fibrous sulfur phase, in which case a specimen 15 months

Table 1. Powder data for pressure-induced fibrous sulfur-selenium ($S_{0.66}Se_{0.44}$); $CuK\alpha$ radiation. Abbreviations: w, weak; m, medium; s, strong; v, very.

hkl	d(Å)		I_{rel}
	Calc.	Obs.	
100	6.80	6.80	vw
110	3.93	3.93	vs
111, 121	2.99	2.99	vs
201, 021	2.74	2.73	s
210, 120	2.57	2.56	m
300	2.27		
211, 121	2.25	2.25	ms
231, 131			
102, 012	2.19	2.19	m
301, 031	2.03	2.03	s
220	1.96	1.95	w
202, 022	1.91	1.91	w
310, 130	1.885	1.879	w-m
221, 241	1.806	1.803	m
131, 311	1.746	1.742	s
141, 341			
212, 122	1.718	1.718	s
132, 232			
302, 032	1.618	1.615	w
401, 041	1.595	1.591	w
320, 230	1.560	1.555	m
103, 013	1.502	1.505	w-m
410, 140	1.483		
321, 231	1.478	1.477	w-m
251, 351			
203, 023	1.403	1.404	vvw
402, 042	1.369	1.370	vvw
330	1.308		
501, 051	1.304	1.305	w
420, 240			
412, 142	1.285	1.283	w
152, 452	1.248	1.248	w
511, 151	1.180	1.177	vw
161, 361			
403, 043	1.141		
104, 014	1.139	1.141	vw
422, 242			
262, 462	1.123	1.122	vw
114, 124	1.108	1.106	w-m
521, 251	1.060		
271, 571			
214, 124	1.053	1.056	w
134, 234			

old and having been irradiated with 35-kv x-rays for about 1500 hours still remains unaltered in any observable manner. When examined about 3 months after it was made, the "single" crystal of the sulfur-selenium phase on which x-ray data had been collected had altered. It had gone partially to the fibrous sulfur (II) type phase.

S. GELLER
M. D. LIND

North American Aviation Science
Center, Thousand Oaks, California

References and Notes

1. S. Geller, *Science* **152**, 644 (1966).
2. R. E. Marsh, L. Pauling, J. D. McCullough, *Acta Cryst.* **6**, 71 (1953).
3. W. R. Busing, K. O. Martin, H. A. Levy, *Oak Ridge National Laboratory Report ORNL-TM-305* (1962).
4. The values given for the helix radius and average S-Se distance should be taken as tentative. Limits of error for these are now approximately ± 0.05 Å.
5. We thank P. B. Crandall for technical assistance.

22 November 1966

Leukocyte Mitosis: Suppression *in vitro* Associated with Acute Infectious Hepatitis

Abstract. *Inhibition of mitosis in vitro was observed in leukocytes from patients with acute infectious hepatitis. Similarly, in cultures of normal leukocytes, after the addition of small amounts of serum from patients with hepatitis, mitosis was suppressed. Although the incidence of mitosis became normal in leukocytes from convalescent patients, there were chromosomal abnormalities.*

The effect of infectious hepatitis on the chromosomes of cells in human peripheral blood was studied during a recent epidemic of this disease, in which more than 100 cases were recognized. Many were symptomatic; others were discovered during a survey of tests for liver functions [primarily for serum glutamic oxalacetic transaminase (SGOT)]. Specimens of blood and serum were obtained from 16 patients, some of whom had been previously karyotyped. Additional samples were obtained from patients at the Massachusetts General, Boston City, and St. Elizabeth's Hospitals. Serums from eight patients with noninfectious hepatic disease and comparable abnormalities of liver function served as controls. Normal specimens were obtained from healthy students and employees in the same institutions.

The standard method of Moorhead *et al.* (1) for culturing leukocytes and preparing chromosomes was used, with two modifications; eight drops of whole blood were added to the culture medium in place of the 1.0 ml of plasma, and the cells were exposed to colcemid for 2 hours instead of the 6 hours suggested by Moorhead. In every case all stained cells were studied. The percentage of leukocytes in metaphase was derived from a count of at least 200 cells.

The initial studies were performed with preparations of peripheral leukocytes obtained from patients with acute infectious hepatitis (hereafter referred to as the direct method). In another method (indirect) 0.1 ml of the serum to be tested was added to cultures of leukocytes obtained from healthy individuals. Preparations to which no serum was added served as culture controls.

No metaphase figures, as judged by the direct method, were seen in specimens obtained from 12 patients

with acute infectious hepatitis. Most of the leukocytes present were contracted and deeply stained or macerated. Chromatin clumping occurred in a few cells, but there was no other sign of mitosis. Eight to 20 percent of cells taken from patients before they developed hepatitis had metaphase figures. Thirteen to 20 percent of leukocytes from convalescent patients after liver function tests had become normal had metaphase figures. However, these chromosomes showed an unusual sticky quality as well as multiple breaks, deletions, and additions (Fig. 1).

Serums from nine patients with infectious hepatitis repeatedly inhibited the development of metaphase figures in normal leukocytes. The incidence of metaphase figures in these preparations ranged from 0 to 0.5 percent of the cells examined. In contrast, control cultures revealed 8 to 20 percent of the leukocytes in metaphase.

Using the indirect method, dilutions of four serums that inhibited leukocyte mitosis were tested. In each case, the serums diluted up to one part in 1000 inhibited mitotic activity. Serums from normal, healthy individuals and eight patients with noninfectious hepatic disease did not suppress metaphase figures in leukocyte cultures; 12 to 20 percent of the leukocytes were in metaphase.

The blood and serum of patients with acute infectious hepatitis have a factor that inhibits leukocyte mitosis and mitosis of normal leukocytes in culture. The inhibition of leukocyte mitosis does not seem to be mediated by elevated concentrations of SGOT or




Fig. 1. Increased stickiness and other chromosomal aberrations. Several breaks are indicated by arrows ($\times 1200$).