


a new micro electrode

for biomedical
stimulation
and recording



the microtrode™

Composite structure of thin film metal layer bonded to drawn glass probes

- Reproducible performance
- Infinite shelf life
- Low resistance
- One micron tip diameter
- Available uninsulated or with resin insulation
- Gold plated pin connector with mating cable assembly
- May be cleaned and reused
- Special structures quoted on request

The Microtrode, connector and cable assembly under \$10.00.

See us at Booth 31 at the ACEMB, Houston, Nov. 18-21.



**TRANSIDYNE
GENERAL
CORPORATION**

462 Wagner Road • Ann Arbor, Michigan 48103

Dom. & For. Pat. Pend.

SC1168

Exogenous and endogenous progesterone stimulates embryo growth up to a certain point, but is deleterious to embryonic survival thereafter. For example, embryonic mortality in sheep increases as the total quantity of progesterone per embryo increases. In rats, steroid administration corrects the effects of reduced caloric intake on pregnancy and enhances placental growth, but does not improve nitrogen balance or liver protein levels.

The time when estrus and ovulation occur may be controlled readily with progestins given orally, intravaginally, by subcutaneous implant, or by injection. The manner of administration, and the type and dose of steroid used, affect the precision of the response, the frequency of ovulation without estrus, and subsequent fertility. All treatments cause some subfertility at the first estrus after treatment due to faulty sperm transport and survival in the female reproductive tract, and to endocrine imbalance. The stimulatory effect of chorionic gonadotropin on urinary steroid output during periods of inhibition of estrus suggests that progestins act at the hypothalamic-pituitary level, suppressing gonadotropin secretion. Pregnant mare's serum gonadotropin can be used to advance the breeding season of sheep. However, the problem of breeding sheep during anestrus or lactation remains unsolved.

The fertility of bull semen improves for a time after collection, reaches an optimum, and then declines as the spermatozoa age. Embryo mortality is inversely related to semen fertility. The optimum fertility, and thus minimum embryonic mortality, for sperm stored at 4°C is on the day after collection. The early improvement in fertility may be due to selective death of spermatozoa containing aberrant chromatin. The decrease in fertility and the increase in early embryonic death found after long storage periods may be due to disturbances of the genetic information system.

Several techniques have been developed for freezing ram semen in colored plastic straws, and bull and stallion semen in pelleted form. Ram semen may also be frozen by the pelleting method if freezing is rapid, if a low concentration of glycerol in a simple medium is used, and the equilibration period is short. There are striking seasonal and individual variations in the freezability of stallion semen. Polyols like xylitol are used for the protection of bovine semen during freezing.

Bull and buffalo semen is preserved at room temperature in a coconut water diluent. The coconut water is collected in Kenya, lyophilized in ampoules, and sent to Germany, where it is stored up to 1 year before it is used. Such semen is reliable for at least 4 days if sufficient attention is paid to laboratory hygiene and the semen is not kept at temperatures below 12°C. This method is being used to increase animal protein supplies of developing countries by eliminating the need for refrigerated semen in artificial insemination programs.

The proceedings of the congress, edited by Professor Charles Thibault, will be published by Station Physiologie Animale, C.N.R.Z., 78 Jouy-en-Josas, France. The site of the seventh congress in 1972 will be West Germany.

E. S. E. HAFEZ

Washington State University,
Pullman, Washington

Calendar of Events—Courses

Mass Spectrometry—Principles and Applications, Ledgewood, N.J., 2-6 December. Will provide a basic understanding of the theory of operation and function of mass spectrometers. Will include functions of major assemblies; mechanisms of ion formation; application in inorganic, forensic medical, and geological fields; principles of focusing; and interpretation of cracking patterns. *Fee*: \$190. (Center for Professional Advancement, P.O. Box 66, Hopatcong, N.J. 07843)

Oceanography—Tour to Coasts of the World: South Pacific, 11 January-1 February. This is a noncredit course which will cover intertidal and nearshore environment, Great Barrier Reef, coral atolls, coastal engineering, tropical marine biology, seismic sea waves, volcanoes, fjords, and beaches. (University of Washington, Division of Evening and Extension Classes, Seattle 98105)

Photochemistry, Moffett Field, Calif., 11-13 December. Intended for scientists, engineers, medical researchers, students, and teachers, the program will emphasize a basic understanding of chemical and physical processes affected by light, including the methods of kinetics and quantum mechanics. The latest techniques for generating, detecting or measuring visible, ultraviolet, and vacuum ultraviolet radiation will also be summarized, with descriptions drawn from research in polymers, medicine, space sciences, and atmospheric photochemistry. (Letters and Science Extension, University of California, Berkeley)

Communication Systems, Washington, D.C., 9-13 December. Modern communication theory and its applications to communication systems, such as radar, satellite communications, and point-to-point digital transmission. (J. E. Mansfield, School of Engineering and Applied Science, George Washington University, Washington, D.C. 20006)