

"Newer or Upper Eocene." Just where in the Vicksburg Group the specimen Pannella, MacClintock, and Thompson analyzed came from is not clear.

A specimen of *Cardita planicosta* (Lamarck), said to come from the Claiborne Formation at Bells Landing, Alabama, is used for a middle Eocene point. The Claiborne Group is indeed middle Eocene; however, none of its formations crop out at Bells Landing (3, 4). This is the type locality of the Bells Landing Member of the Tuscaloosa Formation, which is latest Paleocene in age (5). If the locality data are assumed to be correct, the species they used is probably *Venericor ap-smithii* (3).

For radiometric data, Pannella, MacClintock, and Thompson refer to Kulp (6) rather than to the more recent scale in Harland *et al.* (7). Use of the latter scale alters the ages of some of their points. For example, their only late Tertiary point is based on *Mercenaria campechiensis ochlockoneensis* (Mansfield), which they correctly dated as late Miocene but to which they assign a radiometric age of 18 million years. Assuming that their identification of the specimen is correct, this subspecies is known only from the *Cancellaria* zone of the northern Florida upper Miocene, which is considered to be youngest Miocene on the basis of its molluscan fauna (8). Harland *et al.* date the beginning of the Pliocene at 7 million years ago and the beginning of the late Miocene at 12 million years ago. Therefore, the Miocene point of Pannella, MacClintock, and Thompson would have been better plotted at about 8 rather than 18 million years ago. Also, by reference to the Harland scale, the *Crassatella mississippiensis*, which we suggest is Oligocene rather than late Eocene, would probably be nearer to 31 than 40 million years ago. The specimen called *Cardita planicosta* (Lamarck), probably late Paleocene rather than middle Eocene, would be dated at about 55 million years ago.

Biological and statistical weaknesses are indicated by the tendency of Pannella *et al.* to use single specimens to represent synodical-month patterns [6 out of 11 time intervals in their Table 1 (1)]. If poor preservation and ambiguity of growth patterns make the counts of growth increments highly subjective, as they state, it would be essential to derive magnitude of the synodical month from a statistical sample of a

fossil population rather than from an individual specimen.

Replotting the increments per month at the corrected absolute ages changes the curve for the Tertiary somewhat. Instead of a more or less steady decrease from the Maestrichtian (Fox Hills Sandstone, which is probably better plotted at about 70 rather than 72 million years ago) to the present, there is no significant change in the length of the synodical month from the Maestrichtian to the late Paleocene, but then there is a rapid decrease from the late Paleocene to the present.

JOSEPH E. HAZEL

U.S. Geological Survey,
Washington, D.C. 20242

THOMAS R. WALLER

Smithsonian Institution,
Washington, D.C. 20560

References

1. G. Pannella, C. MacClintock, M. Thompson, *Science* **162**, 792 (1968).
2. T. A. Conrad, *Proc. Acad. Natur. Sci. Phila.* **3**, 280 (1847).
3. J. A. Gardner and E. O. Bowles, *U.S. Geol. Surv. Prof. Paper* 189-F (1939).
4. F. S. McNeil, *U.S. Geol. Surv. Oil Gas Prelim. Map No. 45* (1946).
5. W. A. Berggren, *Micropaleontology* **11**, 279 (1965).
6. J. L. Kulp, *Science* **133**, 1105 (1961).
7. W. B. Harland, A. G. Smith, B. Wilcock, Eds., "The phanerozoic time-scale," *Quart. J. Geol. Soc. London* No. 120 S (1964), whole volume.
8. A. A. Olsson, in A. A. Olsson and R. E. Petit, *Bull. Amer. Paleontol.* **47**, 217 (1964).

11 December 1968

Hazel and Waller checked the information which we should have verified with some of our museum specimens. The fact that they did not need to look at the specimens to discover the taxonomic errors shows how patent these are. Concerned as we were with the chronological information recorded in the shells we neglected taxonomic questions.

We selected the data from the safest counts, and, if these happened to be from one specimen only, we accepted them as preliminary figures and weighted them accordingly for mathematical analysis. The use of a single specimen was unfortunate, but, when chronological data are sought, the suggested analysis of a population of probably contemporaneous specimens is not more sound from a statistical viewpoint. One would risk repeated counts of the same time interval and might think that he is gathering a random sample. In theory the best paleontological clocks would be continuous suites of not entirely

contemporaneous but slightly overlapping individuals that reach the age of Methuselah.

We ran the program again using the same polynomial formula (1) and another program to obtain the chi-square statistic in order to determine the statistical significance of the age corrections suggested by Hazel and Waller. The best-fitting curves were, again, the polynomial of order 4; the chi-square showed statistically no better fit with the corrected data than with the uncorrected data. Since the magnitude of the suggested changes is small in relation to the uncertainties associated with each point, the fit of the data is not significantly affected. When we lump together counts from specimens coming from lower and upper parts of an entire period, as we have for the Pennsylvanian figure, the corrections are not critical, and it is premature to conclude that the events which brought about the change in slope occurred later than the speculated time in our paper. However, as more data become available (we are concentrating on the Mesozoic-Cenozoic interval) the suggested corrections will probably become statistically important.

GIORGIO PANNELLA

COPELAND MACCLINTOCK

Department of Geology and
Geophysics, Yale University,
New Haven, Connecticut 06520

Reference

1. G. Pannella, C. MacClintock, M. Thompson, *Science* **162**, 792 (1968).

20 February 1969

Bird Feathers and Radiation

Working independently, I (1) reported the results of experiments paralleling the feather-reflectance measurements made by Lustick (2) on white and dyed zebra finches. We were both led to similar conclusions by our results—an increase in surface feather temperature due to absorbed radiation leads to a decreased thermal gradient from the skin to the feather surface.

FRANK H. HEPPNER

Department of Zoology, University
of Washington, Seattle 98105

References

1. F. H. Heppner, thesis, University of California, Davis (1967); *Condor*, in press.
2. S. Lustick, *Science* **163**, 387 (1969).

4 February 1969