


Autoclavable

...the extra dimension

Use the new Nalgene® Carboy to prepare, store, and autoclave 5-gallon batches of culture media, distilled water, and other solutions. This carboy (Cat. No. 2250) is precision molded of heavy-duty polypropylene to withstand repeated autoclaving. Only 2½ pounds light, has screw closure and two carrying handles for maximum safety and convenience.

Specify Nalgene Labware from your lab supply dealer. Ask for our Catalog or write Dept. 21101, Nalgene Labware Division, Rochester, N.Y. 14602.



It may turn out, then, that Europe seems to have good solutions to problems because she hasn't really had the problems. Certainly there are many others in which Europe has lived with the situation longer than we have, and really has the answers. But I suspect we will find fewer of these, and more of the former, than either we or the Europeans expect at the present time.

ALFRED B. MASON
136 Flushing Avenue, New York 11251

Lead Poison in Putty

Oberle's article "Lead poisoning: A preventable childhood disease of the slums" (5 Sept., p. 991) contains some errors of omission. Not only has titanium dioxide been used as an opacifier since the 1940's, but previously zinc sulfide pigments (such as lithopone) were used for a number of years. Kalsomine has also been used for many years for very cheap paint jobs.

However, the most serious omission is the failure to mention the most important lead hazard to all children: old sticks of glazing putty. Anyone who has repainted housing, where windows have been installed in wooden frames, has had to replace defective putty, which is brittle and comes loose from the window, often in sticks 1 to 4 or 5 inches long. This putty contains a high percentage of lead; the dry sticks resemble lollipops or candy bars and children will suck on them or chew them. I think that a thorough study of this problem will show that glazing putty is far more of a hazard than paint as a cause of lead poisoning in children.

ALBERT M. ARONOW
P. O. Box 3548, Terminal Annex,
Los Angeles, California 90054

Sequoias' Dependence on Fire

We were greatly pleased to read Oberle's commentary on the ecological drawbacks of forest suppression (8 Aug., p. 568), and we appreciate your assistance in changing the long-standing American philosophy that all fire in the forest is bad. The concept of fire as a natural environmental factor before the advent of man has been a most difficult one to establish. Perhaps this has been due to man's role in starting forest fires and perhaps to the fact that man has too long been inclined to see trees as

individuals rather than as dynamic communities in which fire has always played a significant role.

Inasmuch as we are engaged in a 10-year study of the relationship of fire to sequoia regeneration, some of our findings may be of interest. It became evident that long-time fire prevention and suppression had created two conditions in many of the national park sequoia groves, both of which were contrary to established park policy:

1) Fuels, once consumed by regular fires, had accumulated to unprecedented amounts, presenting uncommonly great fire hazards.

2) Plant succession, which was previously reversed by fire, had progressed in most groves to the point that the giant sequoia was almost completely unsuccessful in reseeding itself. Where protection is complete, there is a tendency for the sequoia to disappear.

We recommended to the National Park Service that fire be used on an experimental basis, primarily to determine just what the optimum conditions for sequoia regeneration were. In March 1964, four study areas were approved by NPS for our use in the Redwood Mountain Grove of Kings Canyon National Park. Few young sequoias were growing on any of the areas. It became abundantly evident that accumulations of fuel on three of the four areas were so great as to preclude broadcast burning. At these areas, it was necessary to pile logs and limbs by machine. In addition to increasing the costs, the machinery greatly disturbed the natural soil conditions we wished to study. We discovered, however, that where light and soil moisture conditions were adequate, seedling establishment tended to be proportional to the intensity of the fire. In short, seedling survival was better in burn pile soils where incineration temperatures penetrated several inches into the soil. Such temperatures probably aided in sterilization against pathogenic fungi, reduced competition with established plants, improved the soil wetability and structure, and may have removed ectocrines. But this created a problem of controlling fires of sufficient intensity to provide conditions for sequoia regeneration while still protecting the primary resources—the parent giant sequoias and other valuable species making up the community.

Recently, the administration of Sequoia and Kings Canyon National Park began a fuel reduction program in one of the critical areas of the Redwood Mountain Grove. Under carefully prescribed

conditions for burning, ground fuels could be reduced effectively and cheaply and, at the same time, create sufficiently hot spots of fire to provide good reseeding conditions for sequoia. This initial plan involves about 35 acres. While this may seem like a modest beginning, it is a beginning and the NPS stands to learn much, both about burning techniques, and the ecology of the giant sequoia.

The NPS is now recognized as one of the leaders in accepting fire as a manipulative tool in land management. While we do not advocate fire as a panacea for forest problems, we do feel that it has useful applications in many situations.

RICHARD J. HARTESVELDT
H. THOMAS HARVEY
HOWARD S. SHELLHAMMER
RONALD E. STECKER

*Department of Biological Sciences,
San Jose State College,
San Jose, California 95114*

Comparative Pathology


The Registry of Comparative Pathology has been established at the Armed Forces Institute of Pathology as a part of the American Registry of Pathology. This new registry is operated jointly by the Armed Forces Institute of Pathology and Universities Associated for Research and Education in Pathology. The objective of the center is to serve as an information exchange for scientists interested in the study of animal models of human disease and other aspects of comparative pathobiology. Information will be gathered about primates, domestic animals, such as horses and cattle, zoo animals, laboratory animals, fish, birds, and, to some extent, invertebrates.

A news bulletin will be published, beginning in October 1969, to describe the activities of the Registry of Comparative Pathology. This will be available without charge to scientists throughout the world. Those interested in receiving this bulletin should request it from the Registrar, Registry of Comparative Pathology, Armed Forces Institute of Pathology, Washington, D.C. 20305.

Members of the advisory committee of the Registry of Cooperative Pathology are: Robert W. Wissler, T. C. Jones, Kurt Benirschke, Robert W. Leader, Donald B. Hackel, and Robert W. Squire.

ROBERT W. LEADER
Rockefeller University, New York 10021

Gas Exchange Studies? THE WARBURG RESEARCH MODEL UFL



— the newest answer from Bronwill

- 15 stations with stationary manometers, shaking flasks
- flexible stainless steel capillaries couple fixed manometers to shaking flasks — you can read the manometers while the flasks are shaking
- all components of each manometric system are interchangeable — and pre-calibrated
- complete system is impermeable to all gases
- cold light illumination for studies of photosynthesis
- shaking frequency and amplitude continuously variable
- bath uniformity better than $\pm 0.01^\circ\text{C}$
- fully solid state electronics
- all controls — including contact thermometer — grouped at eye level

Ask your Bronwill dealer for a demonstration and for your copy of the new catalog which describes the complete line of Warburg instrumentation. Or write directly to us.



BRONWILL SCIENTIFIC, INC.

A leader in scientific instrumentation
Box 7824 — Rochester, New York 14606