Letters

Mortality and Fertility: Causal Link

Frederiksen's article "Feedbacks in economic and demographic transition" (14 Nov., p. 837) lends substance to what was previously only a "strong clinical impression" of mine. While working at the Pakistan SEATO Cholera Research Laboratory in East Pakistan, I was concerned, and in fact told by many people, that the control of mortality from cholera and choleralike diarrheal illness or any other cause was not a good thing in such an overpopulated area. In fact the neo-Malthusian party line would have it that we were actually hurrying the day of a catastrophic famine in the area. The village people I talked with and treated who had experienced the loss of one or more of their children as a frequent and expected occurrence seemed to be little inclined to control their fertility. In fact in this context the woman who was unable to bear children was a very unhappy individual indeed. It was also a myth that grief over the loss of family members was any less real or intense simply because of the frequency with which this event occurred. Thus in one particular overpopulated area at a personal level it seemed crystal clear that the regular loss of a very large number of offspring provided an urgent drive toward a high birth rate. This should not be surprising to any biologist since it is not unique to the human animal but is a rather general rule that populations of organisms with high rates of attrition will have high birth rates.

Fortunately, some answers will become available in the next 5 to 10 years as improved practices in preventive and curative medicine reduce loss of life in babies and children in the vaccine trial area of the Cholera Research Laboratory in rural East Pakistan. In the meantime, however, in lieu of experimental evidence which may be forthcoming in the future, Frederiksen's article certainly gives substance to those who argue for a closer look at the actual

effects of mortality control in the real world. As in other biological systems the simplest way to control fertility may in the long run be to guarantee survival of larger numbers of progeny. Such an approach is also more consistent with the humanities of medicine and civilization as aspired to by our culture.

WILLIAM B. GREENOUGH, III Department of Medicine, Baltimore City Hospitals, 4940 Eastern Avenue, Baltimore, Maryland 21224

What's behind a Label?

We have observed an increasing carelessness among suppliers of radioactive biochemical products. Some examples are:

- 1) A compound, labeled as 98 percent radiochemically pure C¹⁴-phenylalanine, was shown to consist of about 2 percent C¹⁴-phenylalanine with the remaining radioactivity distributed among three other components.
- 2) A compound, packaged as 99 percent radiochemically pure C¹⁴-de-oxyadenosine, was in fact rather pure C¹⁴-adenine.
- 3) A wipe, taken on the outside of an intact vial of P³²-orthophosphate, was counted by liquid scintillation and showed over 20,000 count/min. The package showed no gross evidence of spillage or leakage. Investigators working with low-level radioisotope counting equipment eventually become aware, to their chagrin, of the havoc that can be caused by a few nanocuries of P³² introduced into a lab as a dust from a package.

Until recently, such slovenliness was confined mostly to companies that enjoyed earned reputations for unreliability. The worrisome aspect of this problem is that the more reliable suppliers are now allowing this decline of standards. What to do? The problem of contaminated packages is handled here by

requesting our investigators to assume that all incoming radiochemical packages are contaminated until a wipe test proves otherwise.

Not so easily solved is the problem of questionable purity or reliability of descriptive labels of radiochemicals. However, it may save some users of radiobiochemicals many frustrating hours if they are made aware of these facts. It might help all concerned if investigators would report such incidents promptly to their colleagues and forcefully to their suppliers.

IRVING GOLDMAN
Salk Institute for Biological Studies,
Post Office Box 1809,
San Diego, California 92112


Herbicides: No Margin of Safety

Bryce Nelson's otherwise accurate article on 2,4,5-T (21 Nov., p. 977) contains one error. I estimated that, after a spray operation of 27 pounds per acre of 2,4-D and 2,4,5-T and a 1-inch rainfall, 2,4,5-T in Vietnamese drinking water would reach a concentration of about 50 milligrams per liter. A pregnant woman consuming 1 liter of water per day would thus ingest 50 milligrams per day of this teratogenic agent, not 50 milligrams per kilogram of body weight, as Nelson had me saying. Even so, the more than 1 milligram per kilogram of body weight ingested per day is not far enough below the minimum teratogenic dose in mice and rats to give us any margin of safety. It is for this reason that many of us are pressing for further investigations on the use of the phenoxyacetic acid herbicides at home and in Vietnam. In the meantime, prudence would dictate further restrictions on the use of these compounds until we know more about their teratogenicity in humans.

ARTHUR W. GALSTON
Department of Biology,
Yale University,
New Haven, Connecticut 06520

Purified Transfer RNA's

A number of purified transfer RNA's from *Escherichia coli* have been produced in gram-size quantities and are available without charge from Oak Ridge National Laboratory, as a result of a continuing National Institute



of General Medical Sciences project for the separation of biological macromolecules. The following transfer RNA's from E. coli K-12 MO7, fMet (97 percent), Arg (70 percent), Phe 2 (78 percent), and Glu (~95 percent), and Glu (~85 percent) are fMet (~95 percent), Arg (100 percent), and Glu (~85 percent) are ready for distribution. The values in parentheses are the activities, expressed as amino acid acceptance per terminal adenosine. The arginine tRNA's contain more than one iso-accepting species; all of the others represent a single chromatographic peak. In addition, valine tRNA from E. coli K-12 MO7 will be available shortly.

Qualified investigators in the United States or other countries may request samples of these tRNA's by writing to G. David Novelli, Oak Ridge National Laboratory, P.O. Box Y, Oak Ridge, Tennessee 37830. The letter should include a brief justification and explanation of planned experiments as well as an indication of the quantity needed. These requests will then be reviewed by a committee appointed by the National Institute of General Medical Sciences, and samples will be shipped in accordance with their recommendations.

A. D. KELMERS M. P. STULBERG

Oak Ridge National Laboratory,
Post Office Box X,
Oak Ridge, Tennessee 37831

Well-Rounded Egghead

Bohannan's letter entitled "Fest me no schriften" (14 Nov.) reminds me that the custom of Festschriften comes from Continental university systems along with the more common tradition (thank goodness!) of conferring honorary degrees within the framework of commencement and convocation ceremonies in modern colleges and universities. . . . Bohannan is right that Festschriften are expensive, space-consuming, and may leave long-lasting effects on the inches (or centimeters) of library shelves to which they are relegated. Perhaps they should all be reduced to microfilm so that they could be retrieved on demand by consulting published index lists.

Bohannan's negative approach to these historic and traditional customs leaves a little to be desired. He says, "Because I am the kind of egghead who cannot look at only one article in a book, I also looked at the rest of them (I did not say read—I am not that kind of an egghead)." But what kind of an egghead is he? The reader is entitled to know since he chooses that appellation, and at Northwestern, Bergen Evans takes seriously his thesis that words mean what he means them to mean. I would only point out that most truly macroscopic eggs have two ends, a rounded one and a pointy one. While the latter gets there first (when laid), it is the well-rounded one that has in it the most stuff.

S. R. M. REYNOLDS 933 Olde Hickory Road, Lancaster, Pennsylvania 17601

Eskimos' Language Links

In Dumond's article on the prehistory of the Alaskan Eskimo (28 Nov., p. 1108), he has elected to propagate the myth that the Eskimo-Aleut linguistic stock is "genetically distinct" from any North American Indian language. He does, however, correctly relate the Eskimo-Aleut languages to the languages of northeast Asia, presumably to the Aral-Altaic languages. His citation of Swadesh's 1962 article (1). for placing the Eskimo-Aleut languages "distinct" from North American Indian languages is interesting in that Swadesh states, in 1969:

It was long thought that Eskimo-Aleut had no relationship with American Indian languages, despite a few points of similarity (such as the dual suffix -k of Eskimoan and the -ki plural of Algonkin). Evidence was brought forward in the late 1950s, however, to show systematic agreements pointing toward a common origin, particularly with the Wakashan stock Algonkin-Wakashan phylum of the northwest coast of North America (2).

It appears that the Ural-Altaic, Korean-Japanese, Eskimo-Aleut, and Algonkin-Wakashan languages represent a vast language spectrum of greater diversity than the Indo-European language family, with a prehistoric geographic spread of three continents, and a buried insight into the histories and cultures of a multitude of races and peoples.

CHARLES F. FROMME, JR. 218-39 112th Avenue, Queens Village, New York 11429

References

M. Swadesh, Amer. Anthropol. 64, 1262 (1962).
 ———, "Eskimo-Aleut languages" in Encyclopedia Britannica 8, 706 (1969).