

AMERICAN ASSOCIATION FOR  
THE ADVANCEMENT OF SCIENCE

*Science* serves its readers as a forum for the presentation and discussion of important issues related to the advancement of science, including the presentation of minority or conflicting points of view, rather than by publishing only material on which a consensus has been reached. Accordingly, all articles published in *Science*—including editorials, news and comment, and book reviews—are signed and reflect the individual views of the authors and not official points of view adopted by the AAAS or the institutions with which the authors are affiliated.

## Editorial Board

1970

GUSTAF O. ARRHENIUS	RICHARD C. LEWONTIN
FRED R. EGAN	ALFRED O. C. NIER
HARRY F. HARLOW	FRANK W. PUTNAM
MILTON HARRIS	

1971

THOMAS EISNER	NEAL MILLER
AMITAI ETZIONI	BRUCE MURRAY
EMIL HAURY	JOHN R. PIERCE
DANIEL KOSHLAND, JR.	

## Editorial Staff

## Editor

PHILIP H. ABELSON

Business Manager: HANS NUSSBAUM

Managing Editor: ROBERT V. ORMES

Assistant Editors: ELLEN E. MURPHY, JOHN E. RINGLE

Assistants to the Editor: NANCY TEIMOURIAN, PAULA LECKY

News Editor: JOHN WALSH

Foreign Editor: DANIEL S. GREENBERG\*

News and Comment: LUTHER J. CARTER, PHILIP M. BOFFEY, NANCY GRUCHOW, SCHERRAINE MACK, SAMUEL Z. GOLDBABER, THOMAS P. SOUTHWICK

Research Topics: ROBERT W. HOLCOMB

Book Reviews: SYLVIA EBERHART, KATHERINE LIVINGSTON, CAROL BROWN

Cover Editor: GRAYCE FINGER

Editorial Assistants: JOANNE BELK, ISABELLA BOULDIN, ELEANORE BUTZ, NANCY HAMILTON, CORINE HARRIS, OLIVER HEATWOLE, ANNE HOLDSWORTH, MARSHALL KATHAN, MARGARET LLOYD, VIRGINIA NUSSLE, PATRICIA ROWE, LEAH RYAN, LOIS SCHMITT, BARBARA SHEFFER, RICHARD SOMMER, YA LI SWIGART, ALICE THEILE, MARIE WEBNER

Membership, Subscriptions, and Circulation: THOMAS BAZAN, PATRICIA CAESAR, BETT SEEMUND

\* European Office: 22 Mulberry Walk, London, S.W.3, England (Telephone: 352-9749)

## Advertising Staff

Director	Production Manager
EARL J. SCHERAGO	KAY GOLDSTEIN

Advertising Sales Manager: RICHARD L. CHARLES

Sales: NEW YORK, N.Y. 10036: Robert S. Bugbee, 11 W. 42 St. (212-PE-6-1858); SCOTCH PLAINS, N.J. 07076: C. Richard Callis, 12 Unami Lane (201-889-4873); MEDFIELD, MASS. 02052: Richard M. Ezequille, 4 Rolling Lane (617-444-1439); CHICAGO, ILL. 60611: Herbert L. Burklund, Room 2107, 919 N. Michigan Ave. (312-DE-7-4973); BEVERLY HILLS, CALIF. 90211: Winn Nance, 111 N. La Cienega Blvd. (213-657-2772)

EDITORIAL CORRESPONDENCE: 1515 Massachusetts Ave., NW, Washington, D.C. 20005. Phone: 202-387-7171. Cable: Advancesci, Washington. Copies of "Instructions for Contributors" can be obtained from the editorial office. See also page xviA, *Science*, 27 March 1970. ADVERTISING CORRESPONDENCE: Room 1740, 11 W. 42 St., New York, N.Y. 10036. Phone: 212-PE-6-1858.

## Methyl Mercury

During this century, in the United States, about 75 million kilograms of mercury have been consumed; little information is available on its final disposition or on the concentration of the element at specific points in the environment. Recently, it has become clear that compounds of mercury present a substantial hazard. Of particular significance is methyl mercury, a highly toxic substance that causes neurological damage, produces chromosomal aberrations, and has teratogenic effects. It is mainly in this form that mercury is found in food fishes. Recent studies have elucidated some of the steps in the accumulation.\*

Industrial wastes containing inorganic mercury or phenyl mercury find their way into bottom muds of lakes. There they are converted by anaerobes into  $\text{CH}_3\text{Hg}^+$  or  $(\text{CH}_3)_2\text{Hg}$ . The latter compound is volatile, and it escapes into the water column from the sediment. Though quite stable in alkaline solutions,  $(\text{CH}_3)_2\text{Hg}$  is converted to  $\text{CH}_3\text{Hg}^+$  at low pH. This ion is soluble in water, and it is concentrated by living things, usually appearing in the body lipids. In part, the concentration may come by way of the food chain, but apparently fishes may also accumulate the toxic ion directly. The concentration factor from water to pike is of the order of 3000 or more.

Substantial mercury pollution in the Great Lakes became apparent in March of this year. Mercury concentrations as high as 5 parts per million were reported in some pickerel shipped from Canada. Typical concentrations in fish taken from Lake Erie were 1 part per million or less. Further investigations have confirmed the existence of a major environmental problem traceable to the dumping of large amounts of mercury-containing liquid wastes. This discovery comes as a surprise to most scientists and apparently to federal authorities. However, there was ample reason for looking for such a phenomenon. Episodes in Sweden† and Japan had pointed to dangers arising when large quantities of mercury are discharged into the environment. In Sweden the use of methyl mercury in a seed dressing had led to a drastic decrease in wild bird populations. Near Minamata, Japan, between 1953 and 1960, 111 persons were reported to have been killed or to have suffered serious neurological damage as a result of eating fish and shellfish caught in mercury-contaminated areas. Among the 111 were 19 congenitally defective babies born of mothers who had eaten the fish and shellfish. Later, at Niigata, Japan, 26 cases of methyl mercury poisoning were noted. The affected persons and their families ate fish with a frequency of 0.5 time to 3 times a day; the fish contained mercury at concentrations of about 5 to 20 parts per million.

Physiological and cytological studies have revealed some of the behavior of methyl mercury. It tends to be associated with red blood cells and nervous tissue, and it easily passes the placental barrier, becoming moderately concentrated in the fetus. It can cause chromosomal disorders. Fruit flies consuming food containing methyl mercury at a concentration of 0.25 part per million had offspring carrying one extra chromosome.

It seems unlikely that anything approaching the Japanese observations will be seen in the region of the Great Lakes. There the concentrations of methyl mercury in fish are lower and fish are a less important part of the diet. Nevertheless, we have a substantial and long-enduring problem; even if fresh pollution were stopped, it would be many years before natural processes could cleanse the lakes.

This episode has not led to demonstrable tragedy, but it should remind us how much we risk when we convert our rivers to sewers and our lakes to cesspools.—PHILIP H. ABELSON

\* S. Jensen and A. Jernelöv, "Biological methylation of mercury in aquatic organisms," *Nature* 223, 753 (1969). † Report of an International Committee, "Maximum allowable concentrations of mercury compounds," *Arch. Environ. Health* 19, 891 (1969).