

nitrosamine, for which the lung is the principal target regardless of its mode of entry. Certain respiratory viruses behave similarly, going to lung preferentially, and these also may be important as cofactors or as initiators of DNA alterations in pulmonary cells.

The five-paper section on cellular and functional injury following inhalation exposure is not immediately relevant to carcinogenesis, but it establishes the sensitivity of radioactive thymidine indices of DNA synthesis and cell turnover in the alveolar portion of the respiratory tract, which reflect injury to the lung. Susceptibility of the lung to bacterial infection is also a sensitive measure of injury. However, no clear case can yet be made for relationships between cell turnover and carcinoma or between susceptibility to infection and carcinoma.

The results of cigarette-smoking experiments in rodents and dogs are important to hasten abandonment of such exposure by humans, to locate the carcinogenic agent, and perhaps to stimulate development of a "safer" cigarette. W. Dontenwill's experiments on long-term exposure of hamsters, mice, and rats to cigarette smoke to simulate human exposure have evolved to allow analysis of epithelial changes, including tumors in the larynx and precancerous lesions beyond, and of deposition by ^{14}C labeling of cigarette smoke. In the hamster about 48 percent of deposition is in the head and palate and 52 percent in larynx, trachea, and lungs. In these experiments and also in Auerbach's the airway epithelial changes are impressive, though invasive tumors are infrequent. Cigarette smoke causes accumulation of macrophages in alveoli, and this effect in the absence of clearance of these cells may have important sequelae. The book serves as a description of the techniques of inhalational exposure, a review of some aspects of cell injury, and a progress report on the special subject of inhalation exposure in experimental carcinogenesis. Plans pointing toward a "systems" approach to inhalation carcinogenesis by the National Cancer Institute, the National Institute for Environmental Health Sciences, and the Atomic Energy Commission are described. It should be of greatest interest to investigators in this field.

KAYE H. KILBURN

*Division of Environmental Medicine,
Duke University School of Medicine,
Durham, North Carolina*

Sponges

The Biology of the Porifera. Proceedings of a symposium, London, Sept. 1968. W. G. FRY, Ed. Published for the Zoological Society of London by Academic Press, New York, 1970. xxviii, 512 pp., illus. \$22.50. Symposia of the Zoological Society of London, No. 25.

This volume is the proceedings of a symposium on the biology of sponges which virtually all the investigators in the world currently working with this phylum attended. Approximately 50 people were there. The papers presented at the symposium are ordered under five headings: The Palaeontological Evidence, Spicules and Evolution, Ecology and Biogeography, Biochemistry and the Cell Surface, and Cells, Integration and Morphogenesis. The proceedings as a whole make interesting reading because many of the papers present conflicting views and because there are so few investigators working on sponges that everyone has to be a generalist. It is refreshing to read a paper by an embryologist that refers to a study by an embryologist and vice versa. By and large the students of sponge systematics and phylogeny handle controversial issues in a respectable manner. The investigators working on the biochemistry of the sponge cell surface, however, have gotten to the point where one party, A. S. G. Curtis, denies the existence of the species-specific cell sorting out that the other two investigators in this area, A. P. MacLennan and T. Humphreys, use as an assay in their work. This case seems more like stubbornness than controversy.

One of the highlights of the meeting was the report of W. D. Hartman and T. F. Goreau on coralline sponges. The fossil record indicates that sponges were dominant reef builders during Paleozoic times; there has been no good evidence that sponges play such a role today. Hartman and Goreau have discovered a class of sponges, initially thought to be corals, that appears to be continually consolidating the bases of reefs today in much the same way the fossil forms did.

The part of the symposium that I found the most interesting was the section on integration and morphogenesis. The work that is summarized here is forcing zoologists to change the way they look at sponges. Traditionally the Porifera have been distinguished from more complex animal phyla because they do not have a consistent organiza-

tion above a cellular level. This diagnosis has led to the view that sponges are bags of cells. The elegant experiments of H. Mergner on the formation of the osculum have shown that tissue layers are probably the most important units in the integration of morphogenetic events in sponges. This point is also made very convincingly in the papers of R. Borojević and C. Lévi, who review their excellent histological and experimental studies which define at least some of the factors that control the histogenesis of the various cell types in sponges during embryogenesis.

GARY FREEMAN

*Department of Biology,
University of California, San Diego*

Productivity in Plants

Physiological Aspects of Crop Yield. Proceedings of a symposium, Lincoln, Neb., Jan. 1969. JERRY D. EASTIN, F. A. HASKINS, C. Y. SULLIVAN, and C. H. M. VAN BAVEL, Eds. American Society of Agronomy and Crop Science Society of America, Madison, Wis., 1969. xx, 396 pp., illus. \$10; to members of the societies, \$8.

Based on 16 lectures, this conference report intends to cover the strictly theoretical aspects of the study of crop yield, from climatology to biochemistry and physiology, and selected practical measures. The task seems formidable, but by careful choice of speakers and restrictions of the topic the organizers have succeeded in it. They present a homogeneous, well-balanced volume with contributions of a uniformly high standard and hardly any overlapping.

The dominating theme is the production of organic matter in photosynthesis in relation to microclimate and crop morphology and the distribution of the organic matter leading to the practical yield. Comprehensive treatments are given of plant morphology and leaf pattern, irradiance, gas exchange, and water stress in crop stands, as well as of the biochemistry of photosynthesis and photorespiration insofar as knowledge of these processes is required for understanding production. Perhaps the most striking link between theory and practice is the demonstration of the importance of the "source to sink" principle of nutrient transport for the formation of the plant products. Strictly practical measures for increasing yield, called manipulations, are treated only by means of examples, and more could