

New Lauda C-3T Constant Temperature Circulator with dial-in temperature control.

Only \$375.



Your laboratory could be using a new Lauda Constant Temperature Circulator with dial-in temperature control, and for as little as \$375.

That's the price of the new Model C-3T, with 1,000 watt heater, 8-liters per minute pumping capacity, easy-to-set one-knob thermostatic control, built-in coil for external cooling, all stainless-steel components, reading thermometer, and 30-100°C operating range (0-100°C using external cooling) with $\pm 0.2^\circ\text{C}$ control accuracy.

Need greater control accuracy? Model C-3B has it ($\pm 0.03^\circ\text{C}$), plus pre-set temperature selection (25°, 37° and 56°C) and fine adjustment within $\pm 1.0^\circ\text{C}$, all for \$495.

For literature on these and other Lauda models, write: Lauda Division, Brinkmann Instruments, Cantiague Rd., Westbury, N.Y. 11590. In Canada: 50 Galaxy Blvd., Rexdale (Toronto), Ont.



Circle No. 324 on Readers' Service Card

LETTERS

Organic Farming Methods

The *Science* article by Nicholas Wade, "Boost for credit rating of organic farmers" (5 Sept., p. 777), accurately reflects the content of the report (1) of the Center for the Biology of Natural Systems (CBNS) of Washington University, St. Louis. However, because of the way the report is presented, I would expect a high proportion of readers to misinterpret it with respect to the most fundamental points. All of the comparisons between organic and conventional farms are on a *per acre basis*. The only valid comparison is on a whole-farm basis, inasmuch as the researchers report that the quality of soil and numbers of livestock are similar. I note that the organic farms had 14 more total acres (476 versus 462) but harvested 92 fewer acres of cropland (266 versus 358). Using the report's figures on crop production returns [table 9 in (1), p. 40], I calculate the total for the average organic farm to be \$35,644 (\$134 per acre \times 266 acres) compared to \$46,256 (\$132 \times 358 acres) for conventional farms. This is about 30 percent more for conventional farms on 2 percent fewer acres per farm, which translates into slightly more than 30 percent more food being produced on conventional farms.

The CBNS report contains a single sentence (1, p. 50) which recognizes the deficiency, but few persons are likely to catch its significance: "But because the crops produced on both samples of farms are primarily livestock feeds, not food, a study of the total output (crops and livestock) of both kinds of farms would be needed to determine the actual contribution of each kind to total food production." The same concept is recognized indirectly in item 6 (1, p. 55): "A thorough assessment of how the total level of food production would be affected if there were an appreciable degree of adoption of organic methods." Inasmuch as the authors indicate that livestock are similar in the two systems, it seems unlikely that a study of total farm production would alter my economic analysis to any great extent. Such might not be the case if there were substantial differences in the amount of "feed purchases or sales to balance crops and livestock" (1, p. 13). No data on this matter are presented.

The CBNS report states that most of the additional costs on conventional farms is for fertilizers. A difference in nitrogen requirements is the only valid difference between the two systems in fertilizer costs. The authors make a gross error when they include an additional charge for phosphorus and potassium for conven-

tional farms. An equal charge should be made against organic farms because, unless supplemented by off-farm sources of phosphorus and potassium, organic farm soils will gradually be depleted of these important elements.

The point made above also applies to the CBNS comparison of energy intensiveness. Related to the fertilizer comparison is the use of rock phosphate (energy intensiveness of 0.19 million Btu's) on organic farms versus triplesuperphosphate (energy intensiveness, 0.67 million Btu's) on conventional farms. Based upon many years of research, we expect 3 to 4 pounds of phosphorus in untreated rock to be equivalent to 1 pound in rock that has been acidulated to break the apatite bond. This is a generalized ratio that is adjusted for soil pH, type of crop, and soil test level. The failure to recognize the difference in utility of the phosphorus in rock versus triplesuperphosphate unfairly penalizes the conventional system.

In relation to the additional costs on conventional farms attributed by CBNS mainly to fertilizers, I think it significant that the difference would have been much less had the comparison been made in any of the previous 20 years.

SAMUEL R. ALDRICH

*Agricultural Experiment Station,
College of Agriculture,
University of Illinois at
Urbana-Champaign, Urbana 61801*

References

1. W. Lockeretz, R. Klepper, B. Commoner, M. Gertler, S. Fast, D. O'Leary, R. Blobaum, "A Comparison of the Production, Economic Returns, and Energy Intensiveness of Corn Belt Farms That Do and Do Not Use Inorganic Fertilizers and Pesticides" (Center for the Biology of Natural Systems, Washington University, St. Louis, Missouri, 1975).

Excess Bladder Cancer in Beauticians?

The scientific literature is replete with instances of misstatements and oversimplifications in secondary sources becoming accepted as fact. A recent example occurs in a review article on "Cancer of the urinary tract" (1), in which the following statement occurs in the course of a discussion of bladder cancer: "The Leeds, New York, and New Orleans series all noted an excess of hairdressers and beauticians, who, as indicated by Williams (1962), might have had exposure to dye-stuffs" (1, p. 321).

In fact, this conclusion is hardly supported by reference to the three original papers.

1) The Leeds study (2) reports an excess in incidence of bladder cancer among "male hairdressers" (the English term

"hairdresser" translates to "barber" in the American language—a total of 5 cases versus an "expected" value of 1.8. For female hairdressers, the observed number was 0, the expected value, 0.6.

2) In the New York study (3), the 70 female bladder cancer patients included *one* beautician; there were also four male hairdressers among the 300 men. In their conclusions, the authors state "Also under suspicion are painting, hairdressing, certain textile operations, coal mining, and perhaps plumbing; there is possible ex-

posure to dye in the first 3 occupations" (3, p. 1405). The statement implies that male hairdressers practice hair coloring, which is not common practice in this country.

3) The New Orleans study (4) does not refer to hairdressers or beauticians at all; it speaks only of barbers. It concludes that the comparatively small difference between the number of cancer patients (five) and control patients (two) may have been due to chance only.

The misstatement dissected above may seem trivial. However, in the current con-

troversy on hair dyes, it has been cited in the public press (5, 6) as raising questions concerning the safety of these dyes. One of the articles (5) has, in fact, been dignified by double inclusion in the *Congressional Record* (7).

Two points in conclusion: First, a report (8) on a recent epidemiological study in Massachusetts concludes, "Although suggested by earlier studies [presumably those just discussed] no excess risk was found for . . . [female] hairdressers (one observed, 0.9 expected)." Second, even if beauticians had been observed to incur excess bladder cancer, there would be no justification for linking this to exposure to hair dyes. Tests in which the skin of animals was painted throughout their lifetime with dye composites (9) and numerous skin painting and feeding studies in which single ingredients of dyes were tested have shown no such association.

JOHN MENKART

*Clairol, 2 Blachley Road,
Stamford, Connecticut 06902*

References

1. D. B. Clayson and E. H. Cooper, *Adv. Cancer Res.* 13, 271 (1970).
2. H. M. Anthony and G. M. Thomas, *J. Natl. Cancer Inst.* 45, 879 (1970).
3. E. L. Wynder, J. Onderdonk, N. Mantel, *Cancer* 16, 1388 (1963).
4. L. J. Durham, A. S. Rabson, H. L. Steward, A. S. Frank, J. L. Young, *J. Natl. Cancer Inst.* 41, 683 (1968).
5. J. Brody, *New York Times*, 18 March 1975, p. 26.
6. P. Toynbee, *Observer (London)*, 29 June 1975, p. 1; *Washington Post*, 6 July 1975, p. E-11.
7. T. Eagleton, *Congr. Rec.*, 19 March 1975, p. S.4462; L. K. Sullivan, *ibid.*, p. E.1247.
8. P. Cole, R. Hoover, G. H. Friedell, *Cancer* 29, 1250 (1972).
9. H. J. Kinkel and S. Holzmann, *Food Cosmetic Toxicol.* 11, 641 (1973); C. M. Burnett, B. Lanman, R. Giovacchini, G. Wolcott, R. Scala, M. Keplinger, *ibid.* 13, 353 (1975).

Succulent Stalks

So, that's what does it! Robert H. White ["Occurrence of S-methyl thioesters in urines of humans after they have eaten asparagus" (5 Sept., p. 810)] should now guide his research to a really meaningful conclusion by attempting to establish a correlation between asparagus quality and urine odor specificity, something many an asparagus devotee can do without the benefit of laboratory facilities. (One friend goes so far as to insist that the effect is modulated by the vintage of the companion wine.)

I, for one, choose to remain unconvinced. The uniqueness of the succulent stalks must be due to a certain wonderful and mysterious magic, not to some substance with a name I can't pronounce.

G. N. TSANDOULAS

*Lincoln Laboratory, Massachusetts
Institute of Technology, Lexington 02173*

THIS NEW ALL STAINLESS, ALL-IN-ONE BENCH TOP FREEZE DRYER DOES IT ALL

The new VirTis 3-liter bench top gives you *all* the options. Up to 36 port manifold for ampules and flasks. 3-shelf heated bulk dryer. Clear lucite chamber with internal stoppering for serum bottles from 5 to 125ml. Nitrogen backfill. Even tissue fixation and embedding.

It comes complete with direct reading Thermocouple Vacuum Gauge, wide range Temperature Indicator — and an extra thermocouple for product temperature readout. The stainless steel condenser pulls down to -50°C in just 5 minutes — and you can observe ice build-up through the clear lucite trap top. A built in shell freezer rounds out the system and gives you everything you need:

a complete all around freeze drying facility in just 22" of counter space — BULK · BATCH · MANIFOLD PROCESSING AND BUILT IN SHELLER!

DIRECT FACTORY SERVICE

THE VIRTIS COMPANY P.O. BOX V284 GARDINER, N.Y. 12525

Circle No. 341 on Readers' Service Card