

#### References

1. University of Wisconsin Fusion Feasibility Study Group, *UWMAK-I, A Wisconsin Toroidal Fusion Reactor Design* (Nuclear Engineering Department Report No. UWFD-68, University of Wisconsin, Madison, 1974).
2. ———, *UWMAK-II, A Conceptual Tokamak Power Design* (Nuclear Engineering Department Report No. UWFD-112, University of Wisconsin, Madison, 1974).
3. ———, *UWMAK-III, A Noncircular Tokamak Power Reactor Design* (Nuclear Engineering Department Report No. UWFD-150, University of Wisconsin, Madison, 1976).

The Wisconsin reactor studies indeed show a progression in which the third design is slightly more compact than the first. But the reactor containment building for UWMAK-III would nevertheless be 65 meters high and 75 meters in diameter, with heat exchanger and turbine buildings almost as large. Even though the Wisconsin team has succeeded in boosting the power density of a fusion reactor from 0.7 to 2 Mw/m<sup>3</sup>, there is a very long way to go before achieving the 100 Mw/m<sup>3</sup> power density that would allow a fusion reactor core to be as small as that of a light water fission reactor.

—W.D.M.

#### International Decade of Climatology

We were pleased to see from Willard F. Libby's letter of 28 May (p. 843) that an International Decade of Isotope Climatology Study is proposed, but we are concerned that only isotopic studies have been singled out. The research referred to by Libby is primarily marine in nature; the program, therefore, suffers from the exclusion of the great majority of the terrestrial record. Moreover, much of the marine sedimentary record cannot be interpreted with the degree of high resolution obtainable in the record on land where more rapid sedimentation allows annual or seasonal events to be resolved. This information is surely vital to questions of rapidity of past (and future) climatic change. In addition the marine record may be more climatically buffered than the land record.

Libby's reference to isotopic studies of the Greenland ice core raises doubts about the sole reliance on that approach for the polar regions, since alternative interpretations of the chronology of the glacial oxygen-18 record have been proposed and since complexities in correlations between ice cores have been noted (1). The ice core records do not provide adequate summer paleoclimatic data, and the tree-ring analyses are not applicable to the tundra regions. Thus very large arctic and antarctic areas which are sensitive to climatic change and were probably the first regions to experience

## Slash density gradient spin time with a Sorvall® OTD ultracentrifuge and new vertical rotor.



Sorvall® OTD ultracentrifuges with oil turbine drive and Automatic Rate Controller are ideal for density gradient work. The soft start and soft stop characteristics of the ARC and Reograd mode of deceleration prevent mixing of the gradient at speeds between 0 and 1,000 rpm.

With these features and the new Sorvall® vertical rotor, Sorvall® ultracentrifuges give high resolution with reduced spin times. The rotor holds the tube at a fixed angle of 0° while the gradient reorients from horizontal to vertical. This means the particle must travel only the width of the tube, not the length. It also improves resolution by increasing the surface area and reducing the depth of the starting zone. In fact, the K factor calculated for the highest performance ultraspeed swinging bucket rotor is 45, while the K factor for the Sorvall® TV865 vertical rotor is only 10.

The oil turbine drive eliminates failure-prone gears, belts and brushes. And Sorvall® OTD-50 and OTD-65 ultracentrifuges have self-contained cooling systems — eliminating problems with hard water as well as installation of plumbing, filters, valves and gauges. Both are built with the high quality and attention to detail that have been characteristic of Sorvall® centrifuges for years.

For more information on Sorvall® OTD ultracentrifuges, just write Du Pont Instruments, Biomedical Division, Room 23707A, Wilmington, DE 19898

## Du Pont Instruments



Circle No. 26 on Readers' Service Card

633

# Next Week:

# THE VIKING ISSUE

SCIENCE marks the historic Mars landing of 20 July in next week's special "Viking Issue," dedicated to previously unpublished reports of the mission's preliminary findings. The 13 "Viking" papers cover early results of experiments and analyses, including such topics as:

- Atmospheric Water Vapor
- Infrared Thermal Mapping
- Composition and Structure of the Martian Atmosphere
- Surface of Mars
- Preliminary Meteorological Results
- Physical Properties of the Martian Surface
- Spin Axis of Mars
- Viking Landing Sites
- Radar Observations



AMERICAN ASSOCIATION for the  
ADVANCEMENT of SCIENCE  
1515 Massachusetts Avenue, N.W.  
Washington, D.C. 20005

the onset of glaciation and "abortive glaciations" would not feature at all in the proposed studies.

As recorded, the Holocene period is of highest priority in terms of extreme conditions and the frequency characteristics of climatic fluctuations. In the present formative state of paleoclimatic research, it would seem prudent not to put all our eggs into one (isotopic) basket, but to take this opportunity to involve the entire range of paleoclimatic expertise in a program of mutual data testing and evaluation. Would not an International Decade of Climatology be a preferable research framework?

HARVEY NICHOLS  
JOHN T. ANDREWS  
ROGER G. BARRY  
JACK D. IVES

*Institute of Arctic and Alpine Research,  
University of Colorado, Boulder 80309*

## References

1. W. Dansgaard, S. J. Johnsen, H. B. Clausen, N. Gundestrup, *Medd. Groenl.* **197** (No. 2), 53 (1973); W.S.B. Paterson, paper presented at the Fourth Biennial American Quaternary Association Meeting, Tempe, Ariz., 1976.

## Statistics and Rare Phenomena

Walzer and Gerald (Reports, 19 Dec. 1975, p. 1228) correctly report (1) that, in a study in which they obtained karyotypes of about 10,000 newborn males, the difference between the children of upper-class fathers and those of lower-class fathers in frequency of occurrence of the XYY (and also the XXY) karyotype was not statistically significant (2). The authors interpreted their results as showing "no significant effect of social class upon the frequency of XYY, and of XXY, among newborn males." An Associated Press report of the study stated flatly that XYY "occurs as frequently in the higher socioeconomic classes as in the lower strata" (3).

The frequencies are not stated in Walzer and Gerald's report but can be calculated from the data in their table 2. For XYY they are .00195 and .00107 in the lower-class and upper-class groups respectively (4). This is a difference of 81 percent and would be noteworthy if it were found repeatedly; but because of the rarity of the karyotype it would not reach statistical significance except in much larger samples.

With their sample sizes and overall frequency of the karyotype, the frequency in the lower-class sample would have to be more than 3½ times that in the upper-class sample (5) to show statistical significance. For the difference between the observed frequencies to be

statistically significant at the .05 level, the sample size would have to be increased fourfold (6). In short, the usual statistician's caveat against interpreting lack of statistical significance as counter-evidence must be observed with special scrupulousness when dealing with rare phenomena such as XYY.

DOUGLAS H. BOUCHER

*Division of Biological Sciences,  
University of Michigan,  
Ann Arbor 48109*

## References and Notes

1. The statistical test used in the study was not mentioned; presumably it was a chi-square, maximum likelihood, z-, or similar test.
2. Following a common practice, the authors use the term "significant," not "statistically significant."
3. "Chromosome abnormality hits all classes," *Ann Arbor News*, 17 December 1975, p. 15.
4. These frequencies were calculated from Walzer and Gerald's table with the procedures described in their text: non-Caucasians were excluded from the analysis, maternal age was disregarded, and classes IV and V were pooled as lower-class and I, II, and III as upper-class. The corresponding frequencies of the XYY karyotype are .00117 in the lower-class sample and .00092 in the upper-class sample, a 27 percent difference.
5. A chi-square value of 5.365, for which  $P = .021$ , is given by frequencies of .00273 and .00077 for lower-class and upper-class infants respectively. These correspond to 5 XYY's among 6512 infants for the upper-class sample and 7 in 2564 for the lower-class sample, a 3.56-fold difference.
6. A fourfold increase of Walzer and Gerald's sample corresponds to 20 XYY's in 10,256 cases for the lower-class group and 28 in 26,048 for the upper-class group. These results give a chi-square statistic of 3.63, for which  $P = .057$ , and a maximum-likelihood (G) statistic of 3.96, for which  $P = .047$ .

The analysis described by Boucher was likewise performed by us during the initial examination of our data. If the frequency of individuals with the XYY karyotype in social classes IV and V is indeed as great as .00195, it would still be insufficient to account for the 10- to 20-fold excess of XYY individuals in certain institutional populations. This was the basis for our concluding that whatever effect social and economic factors might have on the institutionalization of XYY individuals, the effect was not *primarily* due to any influence on non-disjunction.

It is still reasonable to believe that the frequency of the XYY karyotype in newborns is not affected by the social class of their fathers. It should be noted that, while an excess of XYY individuals was present in social classes IV and V in the Boston study, a *deficiency* in these two classes occurred in the Edinburgh study (1).

PARK S. GERALD  
STANLEY WALZER

*Children's Hospital Medical Center,  
300 Longwood Avenue,  
Boston, Massachusetts 02115*

## References

1. S. G. Ratcliffe and H. J. Evans, *Lancet* **1975-I**, 1144 (1975).