


High, wise and handsome



The MultiRac fraction collector

LKB's new MultiRac™ fraction collector is a real space saver. On the opposite page you can see that reservoir, column, pump, monitor and recorder all mount on it easily. And you can keep adding decks to take as much more equipment as you want.

The new LKB fraction collector is bright too. It will collect from microliters to liters, adjust fraction size automatically according to OD, stop all flow as the head traverses, and on command will channel all void volume to waste.

Its good looks go well beneath the surface: solid state electronics, rugged materials of construction and a clear, bright, unambiguous LED display all make for an instrument that's safe, dependable and easy to use.

Contact LKB today for full details.

LKB

LKB Instruments Inc.
12221 Parklawn Drive Rockville, MD 20852
301: 881-2510 Telex: 8-9682

58A-311

Circle No. 180 on Readers' Service Card

LETTERS

Pontryagin's Article

In order that the readers of *Science* can assess correctly the reliability and veracity of the letter from Academician Pontryagin on page 1083 of the 14 September issue, they are invited to look at the third from the last paragraph of an autobiographical article by Pontryagin that has appeared in *Uspekhi Matematicheskikh Nauk* [33 (No. 6), (1978)]. A translation of this article, with a disclaimer and a rebuttal, will appear in *Russian Mathematical Surveys*, published by the London Mathematical Society, in November 1979.

ALEX ROSENBERG
Department of Mathematics, Cornell University, Ithaca, New York 14853

Treating Mental Disorders

Gina Bari Kolata reports (*Research News*, 5 Jan., p. 36) on the 1976 discovery by Wurtman and others that the rate at which the brain synthesizes acetylcholine can be increased by increasing the amount of choline in the blood and on the later observations that tardive dyskinesia, a side effect of antipsychotic drugs such as the phenothiazines and the butyrophenones, can be controlled by a high intake of choline or lecithin (which contains phosphatidylcholine). This article appears to indicate that the idea that dietary constituents can influence the functioning of the brain represents a new approach, discovered by Wurtman. In fact, there had been much earlier work in this field, which in recent years has been called orthomolecular psychiatry.

Kolata states that "It was thought that as long as the brain was supplied with oxygen and glucose it would make whatever it needed, independently of the metabolic and nutritional state of the body." It has, on the contrary, been known since long before 1976 that other exogenous constituents of the brain are required for its proper functioning and that, as I stated in my 1968 article (*Science*, vol. 160, p. 265), "Varying the concentrations of substances normally present in the human body may control mental disease." The most striking early work is that on the treatment of psychiatric patients with large doses of niacin, one of the B vitamins (*1*). A dozen other nutrients that affect brain function are also discussed in my 1968 article, which includes in addition a discussion of mechanisms such as increasing the rate of

formation of an important substance by increasing the concentration of a reactant. An example, not given in the article, is increasing the amount of the neurotransmitter acetylcholine by increasing the choline concentration.

I defined orthomolecular psychiatry therapy in 1968 as "the treatment of mental disease by the provision of the optimum molecular environment for the mind, especially the optimum concentrations of substances normally present in the human body." Wurtman has been quoted elsewhere as having said that the treatment of tardive dyskinesia by an increased intake of choline has nothing in common with orthomolecular psychiatry. It is clear, however, that it is a good example of orthomolecular psychiatry; especially interesting because the mechanism of its action is understood in greater detail than in the earlier examples.

LINUS PAULING
Linus Pauling Institute of Science and Medicine, Menlo Park, California 94025

References

1. V. P. Snyderstriker and H. M. Cleckly, *Am. J. Psychiatry* **99**, 83 (1941); A. Hoffer, H. Osmund, M. J. Callbeck, I. Kahan, *J. Clin. Exp. Psychopathol.* **18**, 131 (1957).

Fetal Research Ethics

The 3 November 1978 issue of *Science* (p.540) contained another chapter in the long, unhappy history of the Philipson *et al.* fetal research study (*1*). This experiment led, at the state level, to criminal charges (eventually dismissed) of grave-robbing against the principal investigators, and, at the national level, to a scrutiny of fetal research that culminated in a research moratorium and creation of the National Commission for the Protection of Human Subjects of Biomedical and Behavioral Research.

Throughout the debate on the Philipson *et al.* experiment, ethical questions concerning the fetal subjects of that research have preempted the attention of adversaries, advocates, and reporters like. Ethical questions concerning the *maternal* subjects of that research have not been raised but are just as important. Twenty-five of the 33 patient-subjects, 10 to 22 weeks pregnant, were aborted by hysterotomy (*1*). Hysterotomy is a major surgical procedure not normally used for abortion because it entails hospitalization, risk, discomfort, disfigurement, and because it precludes normal vaginal deliveries in future. From an investigator's point of view, however, hysterotomy as a method of