

16. P. P. K. Smith and P. R. Buseck, *ibid.* **212**, 322 (1981); *ibid.* **216**, 984 (1982).
17. R. Hayatsu, R. G. Scott, M. H. Studier, R. S. Lewis, E. Anders, *ibid.* **209**, 1515 (1980).
18. D. D. Clayton and R. A. Ward, *Astrophys. J.* **224**, 1000 (1978); F. Käppeler, H. Beer, K. Wisshak, D. D. Clayton, R. L. Macklin, R. A. Ward, *ibid.* **257**, 821 (1982).
19. O. K. Manuel, E. W. Hennecke, D. D. Sabu, *Nature (London)* **240**, 99 (1972); D. D. Clayton, *Astrophys. J.* **199**, 765 (1975); U. Frick and R. K. Moniot, *Proc. 8th Lunar Sci. Conf.* (1977), pp. 229–261.
20. E. Anders and D. Heymann, *Science* **164**, 821 (1969); M. Dakowski, *Earth Planet. Sci. Lett.* **6**, 152 (1969); B. Srinivasan, E. C. Alexander, Jr., O. K. Manuel, D. E. Troutner, *Phys. Rev.* **179**, 1166 (1969).
21. R. S. Lewis, J. Gros, E. Anders, *J. Geophys. Res.* **82**, 779 (1977); E. Anders, *Proc. R. Soc. London Ser. A* **374**, 207 (1981).
22. U. Frick and R. O. Pepin, *Lunar Planet. Sci.* **11**, 303 (1980); *Earth Planet. Sci. Lett.* **56**, 45 (1981); P. K. Swart, M. M. Grady, C. T. Pillinger, *Lunar Planet. Sci.* **13**, 788 (1982).
23. P. K. Swart, M. M. Grady, C. T. Pillinger, *J. Geophys. Res.* **87**, A289 (1982). The  $\delta^{13}\text{C}$  values were determined by measuring the ratios 45/44 and 46/44+45 with a VG Micromass 602-E mass spectrometer. Corrections were applied according to H. Craig [*Geochim. Cosmochim. Acta* **12**, 133 (1957)]. Minimum sample size analyzed conventionally was 1  $\mu\text{g}$ , with a standard error of  $\pm 0.01$  per mil.
24. In some temperature steps the quantities of carbon were too small for conventional isotopic measurement, but still above blank level. Such fractions were analyzed by isotope dilution, by first measuring the amount of  $\text{CO}_2$  on a capacitance manometer and then mixing the sample with a metered amount of  $\text{CO}_2$  of known isotopic composition. The reproducibility of this technique, as tested with  $\text{CO}_2$  of known isotopic composition, is  $< \pm 10$  per mil for a sample size of 100 ng.
25. K. Kvenvolden *et al.*, *Nature (London)* **228**, 923 (1970); S. Chang, R. Mack, K. Lennon, *Lunar Planet. Sci.* **9**, 157 (1978); M. M. Grady, P. K. Swart, C. T. Pillinger, unpublished results.
26. B. Srinivasan, J. Gros, E. Anders, *J. Geophys. Res.* **82**, 762 (1977).
27. These treatments removed silicates as well as the organic polymer, and left a residue (0.265 percent of the original meteorite) consisting mainly of elemental carbon and spinel. This residue (2C10) was separated into three grain size fractions by filtration through Nuclepore filters; the two samples analyzed here are designated f (fine,  $< 1 \mu\text{m}$ ) and m (medium, 1 to 3  $\mu\text{m}$ ). See (5) for further details.
28. D. N. Schramm and D. S. P. Dearborn, in preparation.
29. The small amount of dichromate-soluble Xe is isotopically similar to the surface-sited  $\text{HNO}_3$ -soluble Xe and is simply a remnant of it. Other experiments (2, 3, 21) show that it can be removed by prolonged treatment with  $\text{HNO}_3$  or  $\text{H}_2\text{O}_2$  with little loss of carbon. Hence this component does not belong to the dichromate-degradable carbon.
30. This jump—a hint of which was seen in earlier studies (2, 3, 31)—may be caused by a trace of heavy carbon associated with s-Xe (and Ne-E?). Though stepped heating failed to detect these noble gas components in Allende (21, 32), stepped combustion revealed minor amounts of both (33). Unfortunately, we took no isotopic data for B1B after the main peak at 550°C, and hence do not know whether the heavy carbon survived the dichromate treatment.
31. P. K. Swart, M. M. Grady, C. T. Pillinger, *Nature (London)* **297**, 381 (1982).
32. S. P. Smith, J. C. Huneke, R. S. Rajan, G. J. Wasserburg, *Geochim. Cosmochim. Acta* **41**, 627 (1977); B. Srinivasan, R. S. Lewis, E. Anders, *ibid.* **42**, 183 (1978).
33. D. N. Schramm and K. A. Olive, in *Symposium on the Orion Nebula to Honor Harry Draper*, A. E. Glassgold, P. J. Huggins, E. L. Schucking, Eds. (New York Academy of Sciences, New York, 1982), pp. 236–242; see also A. G. W. Cameron and J. W. Truran, *Icarus* **30**, 477 (1977); D. D. Clayton, *ibid.* **32**, 255 (1977); H. Reeves, *Astrophys. J.* **231**, 229 (1979).
34. U. Frick and R. O. Pepin, *Meteoritics* **17**, 217 (1982).
35. I. P. Wright, S. J. Norris, C. T. Pillinger, R. S. Lewis, E. Anders, *Lunar Sci.* **14**, 861 (1983).
36. Y. Kolodny, J. F. Kerridge, I. R. Kaplan, *Earth Planet. Sci. Lett.* **46**, 149 (1980); F. Robert and S. Epstein, *Geochim. Cosmochim. Acta* **46**, 81 (1982); J. Yang and S. Epstein, *Meteoritics* **17**, 301 (1982); N. J. McNaughton, J. Borthwick, A. E. Fallick, C. T. Pillinger, *Nature (London)* **294**, 639 (1981); N. J. McNaughton, A. E. Fallick, C. T. Pillinger, *J. Geophys. Res.* **87**, A297 (1982).
37. A. Bauman, J. Devaney, E. Bollin, *Nature (London)* **241**, 264 (1973). The oxidation rate of carbon below about 1000°C is proportional to surface area, so for a range of  $10^3$  in particle size (Table 1), combustion rates will vary over a range of  $10^6$ . Crystal perfection can easily introduce another factor of  $10^2$ , according to data in Gmelin's *Handbuch der anorganischen Chemie* (Springer-Verlag, Berlin, 1968), vol. 14B2, p. 804.
38. J. W. Larimer, *Geochim. Cosmochim. Acta* **39**, 389 (1975); R. C. Gilman, *Astrophys. J.* **155**, L185 (1969); T. Tsuji, *Proc. Jpn. Acad.* **40**, 99 (1964).
39. We thank L. Aiaerts for preparing the 2C10 samples, I. P. Wright for discussions, and Tang Ming for assistance with the figures. An anonymous referee provided a most helpful, constructively critical review. Supported in part by NASA grant NGL-14-001-010 and the Science and Engineering Research Council.

21 October 1982; revised 10 January 1983

## Helium on Venus: Implications for Uranium and Thorium

**Abstract.** Helium is removed at an average rate of  $10^6$  atoms per square centimeter per second from Venus's atmosphere by the solar wind following ionization above the plasmopause. The surface source of helium-4 on Venus is similar to that on Earth, suggesting comparable abundances of crustal uranium and thorium.

Studies of helium in Earth's atmosphere have a long and checkered history (1–4), providing valuable information on escape, on the composition of the solid body, and on physical processes regulating release of volatiles from the interior. The atmosphere contains  $5.3 \times 10^{-6}$  by volume of  $^4\text{He}$ , formed by decay of uranium and thorium in crustal rocks, and  $6.6 \times 10^{-12}$  by volume of  $^3\text{He}$ , for the most part primordial, emanating from tectonically active regions of the sea floor (3).

Helium escapes from the atmosphere predominately by nonthermal mechanisms, mainly as  $\text{He}^+$  along magnetic field lines open to the interplanetary medium at high latitudes—the polar wind (4). Escape is efficient and must balance production on time scales of the order of  $10^6$  years. The rate of production of  $^3\text{He}$ , 4 atoms per square centimeter per second, may be derived from measurements of the gas dissolved in the ocean, with independent information on the oceanic mixing time (5) and allowing for a small additional source from cosmic rays. The rate of escape, or equivalently production, of  $^4\text{He}$  may be obtained by scaling results for  $^3\text{He}$ , using the observed ratio of the gases in the atmosphere and correcting for minor enrichment of  $^3\text{He}$  above the turbopause (4, 6). The source derived in this fashion,  $2 \times 10^6 \text{ cm}^{-2} \text{ sec}^{-1}$ , is in good agreement with limits established by MacDonald (2) from considerations of the planetary heat flux and available data for uranium, thorium, and potassium.

Helium was first observed as a component of Venus's atmosphere by Mariner 10 (7), and more extensive measurements were made by Pioneer. The abundance of  $^4\text{He}$  in the upper atmosphere is known quite accurately ( $\pm 20$  percent) from mass spectrometric measurements

on the Pioneer orbiter (8) and bus (9). Combined data from the orbiter, bus, and probe (10) indicate a mixing ratio for the gas in the bulk atmosphere of  $1.2 \times 10^{-5}$  with an uncertainty of about a factor of 2.

Helium ionized above the plasmopause on Venus is swept off by the solar wind, as discussed by Dessler (11). The ionization rate may be calculated as described by McElroy *et al.* (12) for oxygen. Using ionization rates for helium of  $3.4 \times 10^{-7} \text{ sec}^{-1}$  for photoionization (13) and  $6.4 \times 10^{-7} \text{ sec}^{-1}$  for electron impact (14), and measured concentrations of  $^4\text{He}$ , we estimate a loss rate of  $10^6 \text{ cm}^{-2} \text{ sec}^{-1}$ . We assume that loss of helium from the nightside is trivial and use data from Brace *et al.* (15) to define an empirical model for the height of the plasmopause. Collisions between hot oxygen atoms and helium were shown by Knudsen (16) to be important for escape of helium from Mars. They are not directly important for escape of helium from Venus but contribute an additional 20 percent to the abundance of helium above the dayside plasmopause. The loss rate for helium may be defined to about the same precision as that for oxygen; in both cases the uncertainty is related primarily to definition of the mean position of the plasmopause. McElroy *et al.* (12) argued that the momentum flux of solar wind is balanced mainly by addition of mass as  $\text{O}^+$ , allowing an independent estimate for escape of oxygen. These considerations suggest that escape rates for oxygen, and by extension for  $^4\text{He}$ , are determined to an accuracy of about  $\pm 30$  percent.

The rate for escape of  $^4\text{He}$  derived here ( $10^6 \text{ cm}^{-2} \text{ sec}^{-1}$ ), combined with the abundance inferred from Pioneer ( $1.8 \times 10^{22} \text{ cm}^{-2}$ ), indicates a lifetime for  $^4\text{He}$  in Venus's atmosphere of

$6 \times 10^8$  years, about 300 times longer than that for Earth. The longer lifetime for Venus reflects the higher abundance of  $^4\text{He}$  (factor of 100) and the lower escape efficiency (factor of 3). Escape from Venus is limited by the relatively small quantities of gas which extend above the plasmopause.

Venus has lost a quantity of  $^4\text{He}$  equivalent to  $1.8 \times 10^{22} \text{ cm}^{-2}$  over the past  $6 \times 10^8$  years. The present atmosphere is unlikely to retain appreciable quantities of its initial helium—escape would lead to a reduction of the primordial abundance by a factor of  $5 \times 10^{-4}$  (17). Contemporary escape should mirror the average quantity of  $^4\text{He}$  released by the solid planet over the past  $10^9$  years. The necessary source would be supplied by decay of uranium and thorium if the abundance of these elements in Venus's lithosphere were similar to that for Earth (18).

Venus's atmosphere also includes measurable quantities of  $^{40}\text{Ar}$ , formed by decay of  $^{40}\text{K}$ . The relative abundances of  $^{40}\text{Ar}$  in the terrestrial and Venus atmospheres suggest that the averaged source of  $^{40}\text{Ar}$  at Venus's surface is less than that for Earth by about a factor of 4 (19, 20). The lower release rate could reflect either smaller abundances of  $^{40}\text{K}$  or less efficient mechanisms for transfer of gas to the atmosphere. Measurements by Venera 13 and Venera 14 (21) suggest that potassium in Venus's surface material is similar to that for typical terrestrial basalts. In this case the lower abundance of  $^{40}\text{Ar}$  on Venus could be due to less efficient degassing, perhaps reflecting a reduced role for tectonic activity in the presence of high surface temperatures (22). However, we might expect a differential effect for helium. Helium can escape by molecular diffusion from a depth of 1 km on Venus, aided by the high surface temperature, while the corresponding diffusion length for Earth is only 400 cm (23).

In summary, Venus's atmosphere contains 160 times more  $^4\text{He}$  than the atmosphere of Earth. The source of  $^4\text{He}$  is similar in magnitude for both planets, suggesting comparable abundances of uranium and thorium. Independent arguments suggest that the two planets have similar abundances of potassium, and it follows that the magnitude of the planetary heat flux should be similar for both bodies. From these results, together with information for nitrogen (24), carbon, water (25), and noble gases (19), we conclude that Venus and Earth had similar origins. The high concentration of primordial noble gases on Venus is attributed to capture of solar wind materi-

als prior to planetary formation (19, 26); the low abundance of  $\text{H}_2\text{O}$  is due to escape of H and O (12, 25, 27) evidenced by enrichment of contemporary deuterium (28).

MICHAEL J. PRATHER

MICHAEL B. McELROY

Center for Earth and Planetary  
Physics, Harvard University,  
Cambridge, Massachusetts 02138

#### References and Notes

1. M. Nicolet, *Ann. Geophys.* **13**, 1 (1957); D. R. Bates and M. R. C. McDowell, *J. Atmos. Terr. Phys.* **11**, 200 (1957); G. Kockarts and M. Nicolet, *Ann. Geophys.* **18**, 269 (1962); G. J. Wasserburg, E. Mazor, R. E. Zartman, in *Earth Science and Meteoritics*, J. Geiss and E. D. Goldberg, Eds. (North-Holland, Amsterdam, 1963), p. 219; W. J. Raitt, R. W. Schunk, P. M. Banks, *Planet. Space Sci.* **26**, 255 (1978).
2. G. J. F. MacDonald, *Rev. Geophys.* **1**, 305 (1963).
3. J. E. Lupton and H. Craig, *Earth Planet. Sci. Lett.* **26**, 133 (1975).
4. W. I. Axford, *J. Geophys. Res.* **73**, 6855 (1968); H. E. Johnson and W. I. Axford, *ibid.* **74**, 2433 (1969).
5. H. Craig, W. B. Clarke, M. A. Beg, *Earth Planet. Sci. Lett.* **26**, 125 (1975).
6. In initial calculations indicating significant thermal escape of  $^3\text{He}$ , unrealistically large values were assumed for the exospheric temperature (2).
7. A. L. Broadfoot, S. Kumar, M. J. S. Belton, M. B. McElroy, *Science* **183**, 1315 (1974); S. Kumar and A. L. Broadfoot, *Geophys. Res. Lett.* **2**, 357 (1975).
8. H. B. Niemann, W. T. Kasprzak, A. E. Hedin, D. M. Hunten, N. W. Spencer, *J. Geophys. Res.* **85**, 7817 (1980).
9. U. von Zahn, K. H. Fricke, D. M. Hunten, D. Krankowsky, K. Mauersberger, A. O. Nier, *ibid.*, p. 7829.
10. J. H. Hoffman, V. I. Oyama, U. von Zahn, *ibid.*, p. 7871.
11. A. J. Dessler, in *The Atmosphere of Venus and Mars*, J. C. Brandt and M. B. McElroy, Eds. (Gordon & Breach, New York, 1968), p. 241; P. A. Cloutier, M. B. McElroy, F. C. Michel, *J. Geophys. Res.* **74**, 6215 (1969).
12. M. B. McElroy, M. J. Prather, J. M. Rodriguez, *Geophys. Res. Lett.* **9**, 649 (1982).
13. P. M. Banks and G. Kockarts, *Aeronomy* (Academic Press, New York, 1973); M. Oppenheimer and C. Downey, *Astrophys. J.* **241**, L123 (1980).
14. See analysis in (12) with cross sections from W. K. Peterson, E. C. Beaty, C. B. Opal, *Phys. Rev. A* **5**, 712 (1972).
15. L. H. Brace, R. F. Theis, W. R. Hoegy, J. H. Wolfe, J. D. Mihalov, C. T. Russell, R. C. Elphic, A. F. Nagy, *J. Geophys. Res.* **85**, 7663 (1980).
16. W. C. Knudsen, *ibid.* **78**, 8049 (1973).
17. The largest error in our calculated lifetime for atmospheric helium is related to the uncertainty (a factor of 2) in definition of the total atmospheric abundance. If the volume mixing ratio of helium were as large as  $24 \times 10^{-6}$  the lifetime would be  $1.2 \times 10^9$  years and the present abundance could have resulted from an initial concentration as low as 0.1 percent.
18. Data from Veneras 8, 9, and 10 show a wide range of uranium and thorium abundances, similar to those observed in rocks like those on Earth [C. E. Fichtel and J. I. Trombka, *Gamma-Ray Astrophysics*, NASA Spec. Publ. SP-453 (1981), pp. 19–60].
19. M. B. McElroy and M. J. Prather, *Nature (London)* **293**, 535 (1981).
20. J. B. Pollack and D. C. Black, *Science* **205**, 56 (1979).
21. V. I. Moroz, report to COSPAR (the Committee on Space Research) (1982); see *Science News* **121**, 214 (1982).
22. D. L. Anderson, *Geophys. Res. Lett.* **8**, 309 (1981).
23. The diffusion coefficient for He is taken as  $5 \times 10^{-4} \exp(-600/T) \text{ cm}^2 \text{ sec}^{-1}$  based on W. G. Perkins and D. R. Begeal, *J. Chem. Phys.* **54**, 1683 (1971). Ten kilometers of crust with 5 ppm of thorium and 1.6 ppm of uranium would produce  $10^6$  alpha particles per square centimeter per second. Additional mechanisms, such as shallow melting in the lithosphere (22), are needed to account for a helium flux of  $10^6 \text{ cm}^{-2} \text{ sec}^{-1}$ .
24. J. H. Hoffman et al., *J. Geophys. Res.* **85**, 7882 (1980).
25. M. B. McElroy, M. J. Prather, J. M. Rodriguez, *Science* **215**, 1614 (1982).
26. G. Wetherill, *Icarus* **46**, 70 (1981).
27. L. H. Brace, R. F. Theis, W. R. Hoegy, *Planet. Space Sci.* **30**, 29 (1982); D. S. Intriligator, *Geophys. Res. Lett.* **9**, 727 (1982).
28. H. A. Taylor, Jr., H. C. Brinton, S. J. Bauer, R. E. Hartle, P. A. Cloutier, R. E. Daniell, Jr., *J. Geophys. Res.* **85**, 7765 (1980); T. M. Donahue, J. H. Hoffman, R. R. Hodges, Jr., A. J. Watson, *Science* **216**, 630 (1982).
29. Supported by NSF grant NSF-ATM-81-17009 and NASA grant NAGW-417.

12 October 1982; revised 1 February 1983

## Desickling of Sickled Erythrocytes by Pulsed Radio-Frequency Field

**Abstract.** Electric fields were found to deform sickled erythrocytes. When the intensity of applied fields exceeded a threshold value, sickled erythrocytes transformed into a spherical shape. Prolonged application of the field usually caused hemolysis of erythrocytes. Deformation of red blood cells could be partly reversed if the field was turned off at an early stage. The cause of desickling may be the interaction of the field with the erythrocyte membrane and also with gelled intracellular hemoglobin S molecules.

In patients with sickle cell anemia, red cell sickling is caused by the intracellular polymerization of deoxy-hemoglobin-S molecules. This polymerization can be inhibited by chemicals that interact with the hemoglobin molecule (1, 2) or with the erythrocyte membrane (3, 4). Chemicals such as cetiedil interact with the red cell membrane to increase water intake and inhibit red cell sickling at a concentration of 1 to 3 percent of intracellular hemoglobin (3–5).

We report a method for reversing red cell sickling without using chemicals. We found that sickled cells were converted within several minutes to a spherical shape by the application of pulsed radio-frequency (RF) fields. Experiments were carried out by placing a suspension of sickled red blood cells in an isotonic saline solution between two parallel Pt-Ir wires (see legends to Figs. 1 and 2 for details). We chose a short pulse width of 5 msec with an interval of 1 second to