

LETTERS

Antisatellite Weapons

R. Jeffrey Smith, in his article of 14 October (News and Comment, p. 140), quotes a remark of mine (incorrectly attributed to Robert Buchheim) about using rockets, balloons, and aircraft to supplant U.S. photoreconnaissance and meteorological satellites, thereby eliminating "the benefit to the Soviet Union from destruction of those satellites." He also quotes me as saying that, if the satellites were destroyed, "we would not be without information," and that "[i]f it cost you a million dollars per flight and you had to do this for 100 days it would be nothing" . . . compared with losing an Army division."

These brief excerpts are accurate, but readers might misunderstand my views, long presented to congressional committees and in my writings.

1) I would regard the destruction of U.S. satellites in peacetime by the Soviet Union as a *casus belli*. For many years I have worked to preserve both principle and reality of satellite invulnerability.

2) During nonnuclear war (that is, absent attacks by the Soviet Union on the U.S. homeland and vice versa), the Soviet Union might have an incentive to attack U.S. satellites if they were aiding in a conventional war in Europe. It is in this context that I advocate supplementing (not supplanting because the worldwide satellite capability would not be destroyed in this case) satellite capability with drone aircraft carrying radar, photographic equipment, and the like, and (indeed) penetrating active enemy territory. The "balloons," and the "meteorological rockets" would be confined to NATO territory in order to provide the equivalent of Navstar and weather satellites for the European theater, and would do as good a job.

In central strategic war, not only would low-altitude satellites be vulnerable even to the limited Soviet ABM system, but their utility would be eliminated by virtue of attacks on their ground stations.

I continue to believe that the U.S. national security would be improved by serious and urgent negotiations to ban antisatellite weapon capabilities and weapons in space, responding to the Soviet initiative on banning the use of force in space of August 1983.

RICHARD L. GARWIN

IBM Thomas J. Watson Research
Center, Post Office Box 218,
Yorktown Heights, New York 10598

EPA's Studies of the Greenhouse Effect

A recent study of the greenhouse effect by the Environmental Protection Agency (EPA), *Can We Delay a Greenhouse Warming? (1)*, concludes that significant global warming throughout the next century is likely and that fossil fuel policies designed to slow the rate of warming will not be effective until 2050 or later. Three separate items in recent issues of *Science* bear directly on this study. Since each raises important issues regarding its assumptions or conclusions, a response is in order.

The first item is an editorial by Philip H. Abelson (25 Nov., p. 879), in which he expresses considerable optimism about society's ability to reduce CO₂ emissions. He bases this optimism on the "static" level of CO₂ emissions over the last decade, the promise of research on ways to remove CO₂ from flue gases, and hopes of greatly expanding the world's use of biomass fuels. We see nothing in the historical record or in the emergence of either CO₂ control technologies or alternatives to fossil fuels that would allow us to share this optimism.

Although informed investigators can and do differ on the expected rate of CO₂ increase, both the EPA report and a report by the National Research Council (2) underscore the immense difficulty of changing emission trends. New, lower CO₂-emitting fuels are likely to take at least 50 years to significantly penetrate energy markets, and CO₂ control technologies currently suffer from major technical uncertainties and enormous economic burdens. We agree with Abelson that "careful monitoring" of CO₂ and "efforts to develop contingency alternatives" are sorely needed. But we are not hopeful that strategies which rely on market-based fuel substitutions or the emergence of CO₂ control technologies will be effective in significantly delaying a greenhouse warming.

The second item is a letter from A. M. Perry (9 Dec., p. 1072). Perry agrees with our projections of temperature rise during the first half of the next century (roughly 2°C), but questions whether a rise as high as 5° to 10°C "from full exploitation of the world's recoverable resources of fossil fuels" is even a reasonable speculation (we estimated a total rise of 5°C—3.1°C due to CO₂ alone—by 2100). Perry argues that policies to limit fossil fuels might be both effective and practical. He also takes issue with our assumptions about the growth of greenhouse gases other than CO₂. He points

out that (i) these gases (CH₄, N₂O, CFC-11, and CFC-12) are responsible for most of the projected temperature rise in our lowest CO₂ scenarios, (ii) the effect of these gases on temperature is uncertain, and (iii) some of the gases may be subject to control.

First, we agree that policies to limit the use of fossil fuels could substantially dampen the extent of warming in the long run. Our own results demonstrate this (for example, a simulated ban on shale oil and synfuels reduced the projected 5°C temperature rise in 2100 by 20 percent). We tried to maintain a clear distinction in our report between the almost universal ineffectiveness of fossil fuel policies in the medium run (by 2050) and the sometimes substantial effectiveness of these policies in the long run (by 2100). Unfortunately, this distinction typically was not maintained in press reports.

We also agree that greenhouse gases other than CO₂ are significant to global warming in our analysis. At several points in our report we underscored the sensitivity of our results to assumptions about future levels of these gases. Moreover, one of our key recommendations is that learning more about the sources, fates, and effects of these "trace" gases should be given high priority on future research agenda.

We did not formally test policies to control CH₄ and N₂O, primarily because relatively little is known about the sources of these gases. However, we were conservative in our selection of growth rates and used simple linear rates of 2.0 and 0.2 percent per year, respectively, rather than the historical compound growth rates of the same magnitude reported in the literature (3).

We were also conservative in our assumptions about growth rates for the atmospheric abundance of CFC-11 and CFC-12, since we held worldwide emission rates constant at 1980 annual levels. Unless a worldwide effort to curtail CFC's is undertaken, both aerosol and nonaerosol usage can be expected to grow in all countries that have not regulated aerosol use, and nonaerosol applications will increase in countries that have regulated aerosol use. We also did not consider the potential warming effects of increases in ozone in the upper troposphere and lower stratosphere (4, 5).

The third item is an article by Woodwell *et al.* (9 Dec., p. 1081) on biosphere contributions to atmospheric CO₂. Drawing from recent estimates of forest clearing and agricultural practices, they