

(A DIGITIZER AND INTEGRATED SCIENTIFIC GRAPHICS SYSTEM FOR ONLY \$495.) A versatile digitizing pen;

accurate, reliable and rugged.

A Turnkey Measurement System Linear • perimeter • area • x, y coordinates • scaling • statistics: standard deviations, correlations, regression lines, transforms. A Pocket Drafting System Multiple CAD-like aids auto lines, boxes, elipses, 3-d, enlarge, shrink, invert • color lift-off • symbol libraries • many type fonts.

A Quick-Slide System Bar-graphs • line graphs • all from numbers, with auto-edit • auto-scaling • sizing • conversion • labeling • coloring • programmed for Polaroid Palette. Display and Printout "Slide-Show" on screen • Printout on dot matrix or color printers • choice of size, shape.

We're the scientist's friend. Try the system for 15 days. Absolutely free. Call toll-free 1-800-874-1888 or in California 415-331-3022. The Computer Colorworks, 3030 Bridgeway, Sausalito, CA 94965.

*Apple version \$299

Circle No. 319 on Readers' Service Card

SCIENCE

Posters

The following posters of *Science* covers are available:

30 March 1979, Tropical flowering tree;

23 February 1983, Landsat photo of Detroit, Michigan;

29 July 1983, Cheetah;

2 December 1983, Snowshoe hare;

23 December 1983, Cathedral window/DNA molecule.

Combination of space covers in scroll format: 1 June 1979, 23 November 1979, 10 April 1981.

Price is \$5 each (prepaid).

Write to AAAS, Department POST, 1515 Massachusetts Avenue, NW, Washington, D.C. 20005.

LETTERS

Fusion Program

The article "Hard times in magnetic fusion" by Mark Crawford (News and Comment, 31 May, p. 1069) states that I am "...gone." This letter is to observe that my "goneness" (to paraphrase Mark Twain) has been somewhat exaggerated. In particular, my interest in the magnetic fusion program remains strong, and I have attempted, as a private citizen, to contribute to it.

This year I suggested to congressional committees that the Department of Energy (DOE) be requested to study the concept of a magnetic fusion research facility at a DOE site with remoteness, security, Nuclear Regulatory Commission and DOE supervision, tritium handling capability, available hot labs, waste disposal facilities, low-cost electric power, and scientific and engineering support.

The rationale for this suggestion is based partially on the now generally accepted realization that the future of the magnetic fusion program probably lies in any of several machines which are much smaller than today's large mainline experimental devices and which could be tested much more cheaply and quickly.

Because some small machine concepts may evolve into competitive devices for the production of electricity, it is my opinion that private industry could be drawn into R&D programs for them, if there were a facility where they could be tested with deuterium-tritium burns. (The production of 14 mev neutrons and the resultant induced radioactivity precludes such testing at a university campus or at a conventional industrial laboratory.)

A DOE fusion research facility, which could be on-line 5 to 7 years after commitment, would be easily affordable, even with a reduced fusion budget. It could give life and meaning to a program that may otherwise be in great danger.

The United States will need all the electric generating capacity it can get, starting before the year 2000 and for an indefinite period thereafter. With luck and prudent leadership now, fusion power could enter the generation market early in the 21st century. Citizen involvement from within the scientific community could help make that possible.

MIKE McCormack* McCormack Associates, Inc., 508 A Street, SE, Washington, D.C. 20003

*Former member of Congress.

Trichinosis Test

I am pleased that the ELISA (enzymelinked immunosorbent assay) test developed for swine trichinosis by the U.S. Department of Agriculture received attention in *Science* (News and Comment, 8 Feb., p. 621). Because the eventual application of this important control strategy by private industry will depend upon a realistic understanding of its capabilities and its potential cost-benefit features, several points mentioned in Gina Kolata's article should be clarified.

It is implied in the article that the "window of vulnerability" of the ELISA, that is, the discrepancy in time between development of infective muscle larvae and detection of antibody, is insignificant. However, there is evidence that the muscle larvae may reach the infective stage as early as 16 to 17 days after initial infection, rather than 24 days (1, p. 42). In contrast to heavily infected hogs, many with light infections (one larvae per gram of muscle or less) may not produce antibody levels detectable by the ELISA until 4 to 5 weeks or longer after infection; this provides an interval of 2 weeks or more in which infectious hogs may go undetected (2). Epidemiological investigations reveal that the majority of naturally infected hogs have such light infections (3, 4). How frequently this situation may arise under natural conditions is at present conjectural, and many authorities in this field feel that such lightly infected hogs are not an important public health hazard. Still, it is not clear that serological tests can provide complete detection.

Because ELISA testing of market hogs is not the only potential inspection procedure now being given serious consideration in the United States, the statement in the article that the pooled digestion test has the disadvantage of not allowing convenient tracking of individual muscle samples should be clarified. The procedure as developed in Denmark (and now officially sanctioned by the European Economic Community) easily allows testing of individuals in a positive pool. Given the very low prevalence of infected swine in the United States (about 1 hog in 1000), the probability of finding a positive pool requiring individual testing is only 1 in 50. Recent epidemiological investigations show that most infected hogs are marketed through smaller slaughterhouses (5), many of which will require relatively unsophisticated inspection techniques. Therefore the pooled digestion test could prove to be a successful alternative.

Several other points require comment. The number of deaths from trichinosis reported to the Centers for Disease Control is not several per year; such deaths occur infrequently. Also, it is not correct that pigs from southeastern farms have a high incidence of trichinosis. Except for occasional cases of swine and human trichinosis in Louisiana, the south as a whole appears to have one of the lowest incidence rates for trichinosis of any region of the United States.

Finally, it should be noted that Ruitenberg and Van Knapen and their associates in the Netherlands first published the basic ELISA method for swine trichinosis and provided a scheme for its application for large-scale testing (6).

K. DARWIN MURRELL Helminthic Diseases Laboratory, Animal Parasitology Institute, U.S. Department of Agriculture, Agricultural Research Institute, Beltsville Agriculture Research Center, Beltsville, Maryland 20705

References

- J. B. Villella, in Trichinosis in Man and Ani-mals, S. E. Gould, Ed. (Thomas, Springfield, Ill., 1970), chapter 11, pp. 19-60.
 F. Van Knapen, Vet. Parasitol. 7, 109 (1980); H. R. Gamble, W. R. Anderson, C. E. Grahm, K. D. Murrell, *ibid*. 13, 349 (1983); G. L. Sea-wright, D. Despommier, W. Zimmermann, R. S. Isenstein. Am. J. Trop. Med. Hyg. 32, 1275 (1983)

- Isenstein. Am. J. Irop. Med. Hyg. 32, 1213 (1983).
 W. J. Zimmermann and P. J. Brandly, Publ. Health Rep. 80, 1061 (1965).
 G. A. Schad, Prev. Vet. Med., in press.
 G. A. Schad et al., ibid.
 E. J. Ruitenberg, P. A. Steerenberg, J. M. Brosi, J. Buys, Bull. W. H. O. 51, 108 (1974); E. J. Ruitenberg, I. Lungstrom, P. A. Steerenberg, J. Buys, Wiad. Parazytol. 21, 747 (1975).

On Objectivity

The mind tends to categorize things as "good" or "bad" and then to become relatively impervious to information that runs counter to these initial impressions. For example, disarmament proposals are "good." Some of them would weaken us to the point that war would become more likely, but anyone saying so is called a "hawk," at the very least. Automotive pollution-control devices are "good." Cars equipped with three-way catalysts emit from 20 to 150 times more sulfuric acid than older cars, but to discover this one must read an obscure Environmental Protection Agency paper. Acid rain is "bad," but it is caused by industrial pollution, so one need not worry about cars. Removing lead from gasoline is "good." Older cars need a small amount of lead to lubricate their exhaust valves and prevent worsening of both performance and emissions, but one learns this from car magazines, not scientific journals. Moreover, the composition of unleaded gasoline must be altered to restore antiknock properties. What longterm health effects this alteration causes are unknown, but to my knowledge the question has not been raised.

It is as though doing something "good" absolves one from the responsibility of knowing about, much less preventing, any foreseeable bad results of such action. This mode of thinking is all too human, but it should be guarded against by scientists and particularly by editorial boards.

DAVID C. STOLINSKY University of Southern California, School of Medicine, Los Angeles 90033

Solitary Grazers

Roger Lewin's recent article on increased foraging efficiency in gregarious grazers (Research News, 3 May, p. 567) indicates it is "theoretically" possible for a solitary grazer to maintain a grazing lawn on its own and therefore reap the benefits of enhanced food quality. However, Lewin suggests that predation would select against such solitary grazing patterns. In fact, there are solitary vertebrate grazers that increase feeding efficiency by maintaining "grazing plots." Biorndal (1) found that the green turtle Chelonia mydas maintained grazing plots of young leaves of the abundant turtle grass Thalassia testudinum by consistent recropping. The result was consumption of significantly more digestible forage (11 percent higher in protein and 100 percent lower in lignin) than ungrazed older leaves. Therefore, enhanced nutrition through maintenance of a grazing lawn is not only selected for in some cases of gregarious grazing vertebrates, but also in some solitary grazers. In contrast to gregarious grazers, solitary grazers can avoid concomitant costs associated with increased resource competition while still enjoying enhanced foraging efficiency.

JAMES B. MCCLINTOCK Center for Marine Studies, University of California, Santa Cruz 95064


References

1. K. A. Bjorndahl, Mar. Biol. 56, 147 (1980).

Erratum: In the article, "Generics, Roche joust for Valium market" (News and Comment, 26 Apr., for Valum market (News and Comment, 26 Apr., p. 472), a word was inadvertently omitted from a quote by Wallace Mendelson, a researcher at the National Institute of Mental Health. The quote should have read that there is 'no established way to predict from Valium's effects on a normal person's EEG's, its ability to relieve anxiety, relax muscles, or treat convulsions."


Erratum: The authors of the Research Article "Chromatin structure and de novo methylation of sperm DNA: Implications for activation of the pater-nal genome" (31 May, p. 1061) were Mark Groudine and Kathleen F. Conklin. Conklin's name was mis-spelled in the Table of Contents and in the by-line.





Master polyacrylamide gel electrophoresis.

You don't need to be a polymer chemist to master polyacrylamide gel electrophoresis (PAGE). At least not if you have Bio-Rad's new Technical Bulletin 1156, "Acrylamide Polymerization—A Practical Approach." It includes page after page of useful information on such topics as · critical parameters · polymerization protocols · polymerization analysis · handling of gel reagents. Also a section on troubleshooting! With this document to help, the novice can master PAGE and the experienced user can optimize separations



Circle No. 288 on Readers' Service Card