

vaccine (15) because of the strain independence of the CS proteins. Extensive cross-reactivity has been observed between the CS proteins from different strains of *P. cynomolgi* (16), *P. vivax* (17), and *P. falciparum* (17). In the case of *P. knowlesi*, only two strains (H and P strains) have been tested, and they showed weak cross-reactivity (15). Since the repeating units constitute the immunogenic epitope, recent emphasis has been on making protective antigens by means of synthetic peptide units coupled to a carrier molecule (18). In view of the extreme differences in the repeating peptide units in the two *P. knowlesi* strains, the utility of such synthetic protective antigens may be limited. Indeed, the CS protein of the Nuri strain does not cross-react with the antibody to the repeat region of the H strain CS protein (19). Recent reports (7, 20) have indicated possible diverse repeating units and serological antigenic diversity between the CS proteins of different strains of another simian malarial parasite, *P. cynomolgi*. It remains to be determined whether such differences occur in the CS proteins of different strains of human malarial parasites.

References and Notes

- L. S. Ozaki, P. Svec, R. S. Nussenzweig, V. Nussenzweig, G. N. Godson, *Cell* **34**, 815 (1983).
- R. L. Coppel *et al.*, *Nature (London)* **306**, 751 (1983).
- J. B. Dame *et al.*, *Science* **225**, 593 (1984).
- R. L. Coppel *et al.*, *Nature (London)* **310**, 789 (1984).
- M. Koenen *et al.*, *ibid.* **311**, 382 (1984).
- J. V. Ravetch, R. Feder, A. Pavlovic, G. Blobel, *ibid.* **312**, 616 (1984).
- V. Enea *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **81**, 7520 (1984).
- A. F. Cowman *et al.*, *Cell* **40**, 775 (1985).
- R. A. Young and R. W. Davis, *Proc. Natl. Acad. Sci. U.S.A.* **80**, 1194 (1983).
- F. Sanger, S. Nicklen, R. Coulson, *ibid.* **74**, 5463 (1977).
- J. Messing and J. Vieira, *Gene (Amsterdam)* **19**, 269 (1982).
- G. N. Godson, J. Ellis, J. R. Lupski, L. S. Ozaki, P. Svec, *Philos. Trans. R. Soc. London Ser. B* **307**, 129 (1984).
- R. F. Doolittle, in *The Evolution of Proteins*, H. Neurath and R. L. Hill, Eds. (Academic Press, New York, 1979), vol. 4, pp. 1-118.
- C. Steffen and R. Timpl, *Int. Arch. Allergy Appl. Immunol.* **22**, 333 (1963).
- A. H. Cochrane, R. S. Nussenzweig, E. H. Nardin, in *Malaria*, J. P. Kreier, Ed. (Academic Press, New York, 1980), vol. 3, pp. 163-202.
- D. H. Chen, R. S. Nussenzweig, W. E. Collins, *J. Parasitol.* **62**, 636 (1976).
- E. H. Nardin *et al.*, *J. Exp. Med.* **156**, 20 (1982).
- J. Gysin, J. Barnwell, D. H. Schlesinger, V. Nussenzweig, R. S. Nussenzweig, *ibid.* **160**, 935 (1984).
- S. Sharma, unpublished results.
- A. H. Cochrane *et al.*, *Mol. Biochem. Parasitol.* **14**, 111 (1985).
- M. Goman *et al.*, *ibid.* **5**, 391 (1982).
- S. Anderson, *Nucleic Acids Res.* **9**, 3015 (1981).
- R. A. Young and R. W. Davis, *Science* **222**, 778 (1983).
- W. D. Benton and R. W. Davis, *ibid.* **196**, 180 (1977).
- We thank F. Rheinart and L. S. Ozaki for preparing the DNA and S.-C. Chiang for assistance in making the λ gt11 library. Supported by the National Institutes of Allergy and Infectious Diseases grant 1 R01 AI21496-01.

4 March 1985; accepted 29 May 1985

Selective Attention Gates Visual Processing in the Extrastriate Cortex

Abstract. Single cells were recorded in the visual cortex of monkeys trained to attend to stimuli at one location in the visual field and ignore stimuli at another. When both locations were within the receptive field of a cell in prestriate area V4 or the inferior temporal cortex, the response to the unattended stimulus was dramatically reduced. Cells in the striate cortex were unaffected by attention. The filtering of irrelevant information from the receptive fields of extrastriate neurons may underlie the ability to identify and remember the properties of a particular object out of the many that may be represented on the retina.

JEFFREY MORAN

ROBERT DESIMONE

Laboratory of Neuropsychology,
Building 9, Room 1N107,
National Institute of Mental Health,
Bethesda, Maryland 20205

Our retinas are constantly stimulated by a welter of shapes, colors, and textures. Since we are aware of only a small amount of this information at any one moment, most of it must be filtered out centrally. This filtering cannot easily be explained by the known properties of the visual system. In primates, the visual recognition of objects depends on the transmission of information from the striate cortex (V1) through prestriate areas into the inferior temporal (IT) cortex (1). At each successive stage along this pathway there is an increase in the size of the receptive fields; that is, neurons respond to stimuli throughout an increasingly large portion of the visual field. Within these large receptive fields will

typically fall several different stimuli. Thus, paradoxically, more rather than less information appears to be processed by single neurons at each successive stage. How, then, does the visual system limit processing of unwanted stimuli? The results of our recording experiments on single neurons in the visual cortex of trained monkeys indicate that unwanted information is filtered from the receptive fields of neurons in the extrastriate cortex as a result of selective attention.

The general strategy of the experiment was as follows. After isolating a cell, we first determined its receptive field while the monkey fixated on a small target. On the basis of the cell's response to bars of various colors, orientations, and sizes, we determined which stimuli were effective in driving the cell and which were ineffective. Effective stimuli were then presented at one location in the receptive field concurrently with ineffective stimuli at a second location. The monkey was trained on a task that required it to




Fig. 1. Effect of selective attention on the responses of a neuron in prestriate area V4. (A) Responses when the monkey attended to one location inside the receptive field (RF) and ignored another. At the attended location (circled), two stimuli (sample and test) were presented sequentially and the monkey responded differently depending on whether they were the same or different. Irrelevant stimuli were presented simultaneously with the sample and test but at a separate location in the receptive field. In the initial mapping of the receptive field, the cell responded well to horizontal and vertical red bars placed anywhere in the receptive field but not at all to green bars of any orientation. Horizontal or vertical red bars (effective sensory stimuli) were then placed at one location in the field and horizontal or vertical green bars (ineffective stimuli) at another. The responses shown are to horizontal red and vertical bars but are representative of the responses to the other stimulus pairings. When the animal attended

to the location of the effective stimulus at the time of presentation of either the sample (S) or the test (T), the cell gave a good response (left), but when the animal attended to the location of the ineffective stimulus, the cell gave almost no response (right), even though the effective stimulus was present in its receptive field. Thus the responses of the cell were determined by the attended stimulus. Because of the random delay between the sample and test stimulus presentations, the rasters were synchronized separately at the onsets of the sample and test stimuli (indicated by the vertical dashed lines). (B) Same stimuli as in (A), but the ineffective stimulus was placed outside the receptive field. The neuron responded similarly to the effective sensory stimulus, regardless of which location was attended.

attend to the stimuli at one location but ignore the stimuli at the other. After a block of 8 or 16 trials, the monkey was cued to switch its attention to the other location. Although the stimuli at the two locations remained the same, the locus of the animal's attention was repeatedly switched between the two locations. Since the identical sensory conditions were maintained in the two types of blocks, any difference in the response of the cell could be attributed to the effects of attention.

The task used to focus the animal's attention on a particular location was a modified version of a "match-to-sample" task. While the monkey held a bar and gazed at the fixation spot, a sample stimulus appeared briefly at one location followed about 500 msec later by a brief test stimulus at the same location. When the test stimulus was identical to the preceding sample, the animal was rewarded with a drop of water if it released the bar immediately, whereas when the test stimulus differed from the sample the animal was rewarded only if it delayed release for 700 msec. Stimuli were presented at the unattended location at the times of presentation of the sample and test stimuli, affording two opportunities to observe the effects of attention on each trial (2).

We recorded from 74 visually responsive cells in prestriate area V4 of two rhesus monkeys and found that the locus of the animal's attention in a cell's receptive field had a dramatic effect on the cell's response (Fig. 1A). When an effective and an ineffective sensory stimulus were present in a cell's receptive field, and the animal attended to the effective stimulus, the cell responded well. When the animal attended to the ineffective stimulus, however, the response was greatly attenuated, even though the effective (but ignored) sensory stimulus was simultaneously present in the receptive field. Thus when there were two stimuli in the receptive field the response of the cell was determined by the properties of the attended stimulus.

To characterize the magnitude of the attenuation, an attenuation index (AI) was derived for each cell by dividing the response (minus baseline) to an effective stimulus when it was being ignored by the response to the same stimulus when it was being attended. For the large majority of cells in V4, the outcome of ignoring an effective sensory stimulus in the receptive field was to reduce the response by more than half (median AI, 0.36 for the sample stimulus and 0.33 for the test) (Fig. 2A).

In the design described, the effective




Fig. 2. Comparison of effect of attention in area V4 (A and B), the IT cortex (C), and the striate cortex (V1) (D). An attenuation index (AI) for each cell was calculated by first subtracting its baseline firing rate from the responses to the sample and test stimuli. The responses to stimuli when ignored were then divided by responses to the same stimuli when attended. AI values less than 1 (dashed line) indicate that responses were reduced when a stimulus was ignored. The number of cells is indicated by *n*. For a few cells, irrelevant stimuli were paired only with the sample stimuli.

stimuli at one location in the receptive field always differed in some sensory quality, such as color, from the ineffective stimuli at the other location. Thus attenuation of the response to an ignored stimulus could have been based on either its location or its sensory qualities. For example, for the cell described in the legend to Fig. 1, effective horizontal or vertical red bars were presented at one location while ineffective horizontal or vertical green bars were presented at the other. When the monkey attended to the green bars, the cell's response to the irrelevant red bars might have been attenuated because they were red or because they were at the wrong location. To test whether attenuation could be based on spatial location alone, for some cells we randomly intermixed the stimuli at the two locations so that, for example, red or green could appear at either spatial location on any trial.

When the locations of the effective and ineffective sensory stimuli were switched randomly, the responses of cells were still determined by the stimulus at the attended location. Cells responded well when the effective sensory stimulus appeared at the attended location and poorly when it appeared at the ignored location (median AI, 0.57 for the sample and test stimuli). Thus attenuation of irrelevant information can be based purely on spatial location.

When attention is directed to one of two stimuli in the receptive field of a V4 cell, the effect of the unattended stimulus is attenuated, almost as if the receptive field has contracted around the attended stimulus. What, then, would be the effect on the receptive field if attention were directed outside it? To answer this, for 112 visually responsive cells (including 51 in the original sample) we placed an effective sensory stimulus inside the receptive field and an ineffective stimulus outside (Fig. 1B). The cells gave a good response regardless of which stimulus was attended (Fig. 2B). Thus, when attention is directed outside a receptive field, the receptive field appears to be unaffected. Furthermore, since the firing rates of cells were the same regardless of whether attention was directed inside or outside the receptive field, we can conclude that attention does not serve to enhance responses to attended stimuli.

To test whether the attenuation of irrelevant information also occurs at the next stage of processing after V4, we recorded from 161 visually responsive neurons in the IT cortex. As in V4, when the animal attended to one stimulus inside the receptive field and ignored another, the response to the ignored stimulus was reduced. Unlike receptive fields in V4, which were typically 2° to 4° wide in the central visual field, those in the IT cortex were so large that the responses

of cells could be influenced by attention to stimuli throughout at least the central 12° of both the contralateral and ipsilateral visual fields (the maximum distance that could be tested). The magnitude of the effect was somewhat smaller than in V4 (Fig. 2C), possibly because IT neurons generally gave weaker, more variable responses than neurons in V4.

The results from area V4 and the IT cortex indicate that the filtering of irrelevant information is at least a two-stage process. In V4 only those cells whose receptive fields encompass both attended and unattended stimuli will fail to respond to unattended stimuli. In the IT cortex, where receptive fields may encompass the entire visual field, virtually no cells will respond well to unattended stimuli.

In contrast to area V4 and the IT cortex, there was no effect of attention in V1. When relevant and irrelevant stimuli were in a receptive field (typically 0.5° to 0.9° wide), the animal could not perform the task. When one stimulus was located inside the field and one just outside, the monkey was able to perform the task, but, as in V4 under this condition, attention had little or no effect on the cells (Fig. 2D).

Our results indicate that attention gates visual processing by filtering out irrelevant information from within the receptive fields of single extrastriate neurons. This role of attention is different from that demonstrated previously in the posterior parietal cortex (3), to our knowledge the only other cortical area in which spatially directed attention has been found to influence neural responses. In the posterior parietal cortex, some neurons show enhanced responses when an animal attends to a stimulus inside the neuron's receptive field compared to when the animal attends to a stimulus outside the field.

Since parietal neurons have large receptive fields with little or no selectivity for stimulus quality, these cells may play a role in directing attention to a spatial location (4), but by themselves do not provide information about the qualities of attended stimuli. By contrast, in area V4 and the IT cortex selective attention may allow the animal to identify and remember the properties of a particular stimulus out of the many that may be acting on the retina at any given moment. If so, then the attenuation of response to irrelevant stimuli found in V4 and the IT cortex may underlie the attenuated processing of irrelevant stimuli shown psychophysically in humans (5).

References and Notes

1. C. G. Gross, in *Handbook of Sensory Physiology*, vol. 7, part 3, *Central Processing of Visual Information*, R. Jung, Ed. (Springer, Berlin, 1973), pp. 451-482; L. G. Ungerleider and M. Mishkin, in *Analysis of Visual Behavior*, D. J. Ingle, M. A. Goodale, R. J. W. Mansfield, Eds. (MIT Press, Cambridge, 1984), pp. 549-586; R. Desimone, S. J. Schein, J. Moran, L. G. Ungerleider, *Vision Res.* 25, 441 (1985).
2. Both sample and test stimuli were presented for 200 msec, with a delay between them of 400 to 600 msec. The sample and test were randomly chosen on each trial from a set of two stimuli, and the irrelevant stimuli were chosen from a different set of two. If the animal attempted to perform the task on the basis of the irrelevant stimuli, its performance would be governed by chance. The performance of the animals was 94 percent correct. The cue to the animal to switch the locus of its attention was to delete the test-time stimulus from the previously relevant location for two trials. On the first of these trials, the animals' performance dropped to 65 percent correct and their reaction time increased by 90 msec, indicating that they had been ignoring the irrelevant stimuli. The neural responses on the two cue trials were not counted. The locus of attention was switched frequently enough to achieve a minimum of ten trials per stimulus configuration. Fixation was monitored by a magnetic search coil, and trials were aborted if the eyes deviated from the fixation target by more than 0.5°.
3. J. C. Lynch, V. B. Mountcastle, W. H. Talbot, T. C. T. Yin, *J. Neurophysiol.* 40, 362 (1977); M. C. Bushnell *et al.*, *ibid.* 46, 755 (1981).
4. M. I. Posner, J. A. Walker, F. J. Friedrich, R. D. Rafal, *J. Neurosci.* 4, 1863 (1984).
5. D. E. Broadbent, *Acta Psychol.* 50, 253 (1982); D. Kahneman and A. Treisman, in *Varieties of Attention*, R. Parasuraman and D. R. Davies, Eds. (Academic Press, New York, 1984).
6. We thank M. Mishkin for his support in all phases of the study.

20 March 1985; accepted 12 June 1985

Locus of the α -Chain of the T-Cell Receptor is Split by Chromosome Translocation in T-Cell Leukemias

Abstract. *Mouse lymphoma cells were hybridized with two human acute T-cell leukemias with a t(11;14) (p13;q11) translocation and the segregated hybrids were examined for the presence of the DNA segments coding for the constant (C_α) and the variable (V) regions of the α chain (C_α and V_α) of the T-cell receptor. The C_α segment was translocated to the involved chromosome 11 (11p⁺) while the V_α segment remained on the involved chromosome 14 (14q⁻). The data indicate that the locus for the α chain of the T-cell receptor is split by the chromosomal breakpoint between the V_α and the C_α gene segments, and that the V_α segments are proximal to the C_α segment within chromosome band 14q11.2.*

JAN ERIKSON

*Wistar Institute,
Philadelphia, Pennsylvania 19104*

DOROTHY L. WILLIAMS

*Division of Pathology and
Laboratory Medicine,
St. Jude Children's Research
Hospital, Memphis, Tennessee 36101*

JANET FINAN

PETER C. NOWELL
*Department of Pathology and
Laboratory Medicine,
University of Pennsylvania
School of Medicine,
Philadelphia 19104*

CARLO M. CROCE

Wistar Institute

The locus for the α -chain of the T-cell receptor is at band q11.2 of chromosome 14 (1), a chromosome region that is involved in inversion and translocations in T-cell tumors (2). In Burkitt lymphoma, immunoglobulin gene loci are split by the three chromosome translocations associated with this tumor (3) and are involved in rearrangements with the *c-myc* locus (3), leading to deregulation of the *c-myc* gene involved (4). In this paper, we have tested whether the locus for the α -chain of the T-cell receptor is directly involved in the chromosome rearrangements observed in T-cell neoplasms, since this

locus could be similarly involved in proto-oncogene activation (1).

Cells from two (8511 and 8508) acute T-cell leukemias (ALL) that are characterized by a t(11;14) (p13;q11) chromosome translocation (5) were hybridized with BW5147 mouse T-cell lymphoma cells deficient in hypoxanthine phosphoribosyltransferase (6). The hybrids were selected in HAT (hypoxanthine, aminopterin, thymidine) medium containing 10⁻⁷ M ouabain (7). Several hybrids were derived from the fusion of BW5147 cells with 8511 leukemia cells (hybrid number 517), while the others were derived from the fusion of BW5147 cells with 8508 leukemia cells (hybrid number 515) (Table 1). In T cells from ALL patients with a t(11;14) translocation, the breakpoint on chromosome 14 has been previously assigned to band 14q13 (5), but karyologic analysis of our ALL hybrids indicates that the breakpoint is probably closer to the centromere and involves band 14q11 (Fig. 1), in agreement with other reports (2). The breakpoint on chromosome 11 may also be more proximal than the p13 location previously reported (5).

The presence of the relevant chromosomes in the hybrid clones is summarized in Table 1. Southern blot analysis of DNA from the hybrids (Fig. 2) indi-