

Herbert (2), the size but not the number of preneoplastic foci in livers of rats given aflatoxin was increased by vitamin A to an unspecified extent. Unmentioned by Herbert was the finding that selenium supplementation reduced the focal area by a factor of 5. The third citation reports the effects of ascorbic acid on the in vitro growth of human leukemic cells from bone marrow aspirates (3). From 259 cases studied, 169 aspirates were successfully cultured, and in these growth enhancement was seen in 53 and growth suppression in 28. In the discussion section the author cites six studies demonstrating in vitro tumor suppression and one demonstrating tumor enhancement by ascorbic acid.

The fourth of Herbert's references (4) is to a dietary study of human colon and rectal cancer patients and their case controls. In this work mean fiber intakes were reported as 22.4 ± 0.8 grams per day in the female colon cancer cases and 19.7 ± 0.6 grams per day in the controls, on the basis of a dietary questionnaire. The difference does not appear to be meaningful. A dose-related positive association between fiber and colon cancer was reported in women when fiber intake was broken down into quintiles with no figures given for the number of individuals in each quintile. A protective effect of vitamin C against rectal cancer was found but was not mentioned by Herbert.

I believe the four references provided by Herbert give little support to his original contentions. This area of research deserves more profound analyses of the literature before public statements are made.

BURTON KALLMAN

National Nutritional Foods Association,
125 East Baker Avenue, Suite 230,
Costa Mesa, CA 92626

REFERENCES

1. F. W. Rosa *et al.*, *Teratology* **33**, 355 (1986).
2. J. Colford and R. S. Parker, *Fed. Proc. Fed. Am. Soc. Exp. Biol.* **45**, 1089 (1986).
3. C. H. Park, *Cancer Res.* **45**, 3969 (1985).
4. J. D. Potter and A. J. McMichael, *J. Natl. Cancer Inst.* **76**, 557 (1986).

Hazardous Waste Disposal

Philip H. Abelson (Editorial, 1 Aug., p. 509) has expanded on his earlier editorial about hazardous wastes (7 June 1985, p. 1145). Incineration and biodegradation are indeed preferable to dispersion and deposition, but can be both encouraged and discouraged by technical, economic, environmental, and regulatory constraints (1). Concentrated forms of hazardous wastes are destroyed in incinerators; dilute forms are biodegraded in wastewater treatment plants.

Halogenated organics are reductively dehalogenated, dehydrohalogenated, and microbially metabolized to CO_2 , H_2O , and inorganic halides. Other examples include biodegradations of CH_2Cl_2 (2) and $\text{C}_6\text{H}_6\text{X}_6$ mixtures (3) and conversions of CH_3CCl_3 , CHCl_3 , and CCl_4 , to lower chlorinated homologs by iron (II) porphyrins (4).

Reported shortfalls in incineration capacity apply more to commercial offsite operations than to private on-site operations. Most solid wastes generated by the chemical industry over the past 30 years have been successfully managed on-site in currently active facilities under the direct management of the original generators. These include the bulk of on-site incineration capacity. Several large corporations, including the Dow Chemical Company, committed themselves to waste minimization and waste treatment before the current public regulatory focus. Recoveries of energy or material values from process intermediates minimize wastes before they are generated; incineration destroys organics in wastes after they are generated (5). Both operations reduce ultimate waste volumes and hazards. Long-term monitoring and remedial actions are mitigated by immediate destruction of combustible portions. Noncombustible residues may require secure land disposal. Wastewater treatment and air emission control are also integral to overall solid waste management.

Priority testing should be mirrored by proper data evaluation. Real-world data often are ignored, "worst-case" despoliation scenarios are assumed, technology-forcing controls are thrust upon the regulated community, and confirmations of environmental improvements are not attempted. The "troublesome" aspects of small halogenated hydrocarbons at Superfund sites are not manifestations of hazard or developments of control technologies, but "selected presence" at very low levels. Because of their unique chemical "fingerprints," they are "found" because they are "sought." Environmental ubiquity at some level above "zero" then triggers extended debate and calls for "action."

I support the resolution of socially troublesome aspects of hazardous waste disposal by multidisciplinary science and engineering. "Multimedia" approaches to defining the distributions, fates, and effects of the significant constituents of air, water, and land have recently been addressed (5). We must understand how much of which, where and when, and for how long, might do what to whom and then determine if, how, and what action is appropriate. The hazardous waste "problem" is as much definition and interpretation as it is practice and remediation. The negative perceptions of

"dumps for toxics" perpetuated by some of the public and the media must be supplanted by the positive realities of "proper hazardous waste management facilities" practiced by scientists and engineers.

STACY L. DANIELS

Advanced Technology Applications Research,
Dow Chemical Company,
Midland, MI 48667

REFERENCES

1. S. L. Daniels, paper presented at the AAAS Annual Meeting, Los Angeles, CA, 28 May 1985.
2. G. M. Klecka, *Appl. Environ. Microbiol.* **44**, 701 (1982).
3. T. D. Lickly *et al.*, *Environ. Toxicol. Chem.* **3**, 503 (1984).
4. G. M. Klecka and S. J. Gonsior, *Chemosphere* **13**, 391 (1984).
5. S. L. Daniels, paper presented at the Fifth Conference on Environmental Engineering Education, Michigan Technological University, Houghton, MI, 22 July 1986; C. R. Baillod, *Environ. Sci. Technol.* **20**, 432 (1986).

Mystery Cloud: Additional Observations

After the appearance of our article "Kaitoku Seamount and the mystery cloud of 9 April 1984" (8 Feb. 1985, p. 607), the original data base was supplemented with additional testimony from Captain Van den Berg (KLM Royal Dutch Airlines flight 868) and another pilot (Captain Presley of Flying Tiger Airlines flight 022). Huub Eggen, editor of the Dutch publication *Aarde & Kosmos*, located Van den Berg and supplied us with a transcript of his interview along with six drawings depicting the event as viewed from the cockpit window. We submitted a number of additional questions which were subsequently answered by Van den Berg (1). To summarize the drawings we divide the event into four stages: (i) a towering cumulus-like cloud appearing to rise out of the stratiform layer; (ii) fading of the cloud tower and replacement with a small semicircular halo segment; (iii) expansion of the halo to a full circle; and (iv) further expansion and dissipation. The time elapsed from (i) to (iii) was approximately 5 minutes. Stage (iv) lasted for another 10 to 15 minutes, giving a total observation time of about 20 minutes from Van den Berg's vantage point. At the time Van den Berg was flying on air route A90 near the PAWES intersection at a speed of 500 knots. The change in Van den Berg's position from the beginning to the end of his observations, along with Captain McDade's observations (Japan Airlines flight 036; 33 minutes behind Van den Berg on air route A90; approaching the intersection designated PAWES by the Federal Aviation Administration when he first sighted the cloud at 1349Z Greenwich mean time; 48 miles

south of air route A90 and abeam of PAWES at 1406, when the sphere was dissipating) and Presley's observations (air route A90; behind McDade and approaching the intersection designated SABES) allows us to estimate the size and distance of the halo.

Our conclusion is that original estimated positions were in error. Additional data, primarily from Van den Berg, place the event between the Kurils and Sakhalin. The altitude of the center of the halo at the maximum observed size is estimated to have been greater than 200 miles, and the diameter of the halo is estimated to have been at least 380 miles. It seems unlikely that a ground-based explosion could produce this kind of an effect. It is surprising to us that no official data have been provided by government agencies and that such a significant observation from a region of demonstrated military sensitivity was, and still remains, a mystery. In retrospect, we believe that erroneous assumptions regarding the presumed location of the mystery cloud may have contributed to the early dismissal of a hypothesis that now deserves intense examination. The 10 April 1984 issue of the *Anchorage Daily News* reported (2) that the Soviet Union had informed Japanese officials that missile testing would begin on 9 April in an

area west of the Kamchatka Peninsula. The 12 April 1984 issue of that newspaper reported (3) that "a Japanese aviation official confirmed Wednesday the Soviet Union had scheduled missile tests in the northern Pacific, but not on the day or in the area where the mysterious cloud burst was sighted."

DANIEL L. MCKENNA
Department of Meteorology,
University of Hawaii at Manoa,
Honolulu, HI 96822
DANIEL A. WALKER
Hawaii Institute of Geophysics,
University of Hawaii at Manoa

REFERENCES

1. H. Eggen, *Aarde Kosmos* 5, 292 (1985).
2. L. Campbell, *Anchorage Daily News*, 10 April 1984, p. A1.
3. M. Kuramitsu, *ibid.*, 12 April 1984, p. A1.

Erratum: The Research News article "How unusual are unusual events?" by Roger Lewin (26 Sept., p. 1385) cited the statement "Given a 5% probability of an unusual event in a 1-year study, one should expect a 35% probability of such an event in a 7-year study." The expected correct probability for a 7-year study would be 30.17%.


Erratum: In the article by Jane Menken *et al.* "Age and infertility" (26 Sept., p. 1389), the first sentence under the sidehead "Diagnosis of infertility by failure to conceive within a year" (p. 1390, column two, paragraph two) should have read: "The historical evidence indicates that the proportion of women who were sterile increased rather slowly and almost linearly from the 20's until the early 40's."

New standards of vibration isolation MICRO-g[®] systems

- Standard Post-Mount Isolators combined with Series 76 Honeycomb Tops achieve exceptional levels of attenuation and damping.
- Gimbal Piston isolators attenuate both vertical and horizontal vibration in a single compact module. Wide range of load ratings.
- New Series 76 honeycomb core tops offer superior damping, rigidity and structural integrity.
- Standard sizes to 4' x 12'; top thickness 2", 4", 8" and 12".


Technical Manufacturing Corporation

15 Centennial Drive, Peabody, Massachusetts 01960
Telephone: 617-532-6330 • Telex: 951408




DAMPING

(Decay rate of on-board input)



ISOLATION

(One vertical division equals 90% isolation)




TMC 14R2

Circle No. 184 on Readers' Service Card

PIPETTING? SINGLE-SHOT MICROLITER DISPENSING?

Try the all-new
FMI
micro *TT*-petter!



It provides a simple, low-cost means of dispensing very small amounts of fluids for applications such as pipetting, reagent preparation, syringing, precision sampling, repetitive and volumetric dispensing. The variable displacement, valveless RR piston design makes it possible to dispense volumes of 2 to 100 μ l per shot with incremental adjustment of less than 0.5 μ l. PiP units are also reversible for aspirating fluids or back flushing the system.

A simple, sealed pushbutton hand/foot pendant switch makes a momentary contact for one dispense per push when the mode switch is in the down (Singles) position, or it may be held down for purging or reagent rinsing when the mode switch is in the up (Repeat) position. Using a combination of forward and reverse modes makes dilutions especially easy. Wetted surfaces of ceramic and fluorocarbon provide for maximum chemical compatibility. A three-speed 110 VAC 60 Hz synchronous motor provides three shot velocities to meet your fluid and vessel requirements.

Two standard models are available at a price of \$450 each. F.O.B. FMI:

Model PiP0CKC—0 to 50 μ l.
Model PiP1CKC—0 to 100 μ l.

Available for same day shipment.

FMI FLUID METERING, INC.

P.O. Box 179
29 Orchard St., Oyster Bay, NY 11771
516-922-6050 • Telex 5101001281

Circle No. 174 on Readers' Service Card