

do not develop diabetic vascular disease (3). In contrast, GH-induced diabetes in dogs is accompanied by diabetic vascular lesions, and in diabetic rats GH injections resulted in increases in vascular basement membrane thickness (4). There is a correlation between GH levels and increased skin capillary fragility in diabetics (5). Dilman *et al.* (6) feel that "paradoxical responses" wherein GH rises during glucose loading may be significant to the clinical pathology of aging.

Pituitary tumors associated with GH production are present in high incidence in aging rodents, and age-related kidney damage (glomerulosclerosis) in the rat was greatest in those animals with such tumors (7). Pituitary tumors are frequently found clinically at autopsy in aged patients where an incidence of 20 out of 152 unselected subjects (8) has been reported.

The loss of GH (pituitary ablation) can reverse the clinical retinopathy and renal complications of diabetes (9). Hypophysectomy in rats can reverse age-related glomerulosclerosis (10), and there is evidence for a hypothalamic pituitary neuroendocrine clock that programs aging with hypophysectomy-producing levels of rejuvenation in rats and mice (1), although in contrast, GH may be responsible for maintenance of protein synthesis in aging rodents (11).

We have to be concerned about not only the potential promiscuous clinical use of GH for cosmetic effects but also the current related "health food" use of arginine, which has GH-releasing activity. The latter is purportedly used to increase muscle mass cosmetically, without exercise. The action of GH on aging is not clear, and we need good studies in animal models to determine the effects of GH or GH-releasers to avoid premature clinical cosmetic use that could lead to long-term injury such as is seen in acromegaly or diabetes.

WILLIAM REGELSON
Medical College of Virginia,
Post Office Box 273,
Richmond, VA 23298

REFERENCES

1. W. Regelson, in *Intervention in the Aging Process*, W. Regelson and M. Sinex, Eds. (Liss, New York, 1983), part B, pp. 3-52.
2. A. V. Everitt, in *Hypothalamus Pituitary and Aging*, A. V. Everitt and J. A. Burgess, Eds. (Thomas, Springfield, IL, 1976), pp. 262-281; J. A. Lindfoot, in *Endocrine Control of Growth*, W. H. Daughaday, Ed. (Elsevier, New York, 1981), pp. 207-267.
3. T. J. Merimee, S. E. Fineberg, W. Hollander, *Diabetes* 22, 813 (1973).
4. K. Lundback and A. B. Hansen, in *Secondary Diabetes*, S. Podolsky and M. Viswanathaw, Eds. (Raven, New York, 1980), pp. 373-390; R. Osterby and K. Seyer-Hansen, *Diabetologia* 15, 487 (1978); J. M. B. Bloodworth, Jr., in *Endocrine Pathology: General and Surgical*, J. M. B. Bloodworth, Ed. (Williams and Wilkins, Baltimore, 1982), pp. 15-21.

5. K. G. Alberti and C. M. Press, in *Complications of Diabetes*, H. Keen and J. Jarrett, Eds. (Arnold, London, 1982), pp. 231-270.
6. V. M. Dilman *et al.*, *Exp. Gerontol.* 14, 217 (1979).
7. P. L. Horn and J. T. Wood, *ibid.* 16, 141 (1981).
8. W. Singer *et al.*, in *Endocrine Aspects of Aging*, S. G. Korenman, Ed. (National Institute of Aging, Endocrine Society, and Veterans Administration, Washington, DC, 1979).
9. E. Kohner, C. T. Dollery, T. R. Frazer, C. J. Bulpitt, *Diabetes* 19, 703 (1970); J. T. Ireland, G. C. Viberti, P. J. Watkins, in *Complications of Diabetes*, H. Keen and J. Jarrett, Eds. (Arnold, London, 1982), pp. 137-178.
10. J. E. Johnson, Jr., and R. G. Cutler, *Mech. Ageing Dev.* 13, 63 (1980).
11. K. R. Lynch *et al.*, *Cell* 28, 185 (1982).

SDI Research Funds

The 7 November article "Mathematicians look to SDI [Strategic Defense Initiative] for research funds" by Gina Kolata (News & Comment, p. 665) distorts my comments at a 7 October briefing held at the National Academy of Sciences.

I did not ask any question at that briefing. Instead I rose to comment on the need to strike a balance in government-sponsored basic research in mathematics between group research and the research of individual mathematicians pursuing their own ideas, especially in view of the past tendency to fund group activity at the expense of individual projects.

In short, new money for mathematics does not contribute to the long-term health of our discipline if, in aggregate, it tends to diminish the incentive of gifted mathematicians to develop their own ideas.

G. D. MOSTOW*
Department of Mathematics,
Yale University,
New Haven, CT 06520

*President-elect, American Mathematical Society.

Net Primary Production: The Tomato Example

Roger Lewin's News & Comment article "A mass extinction without asteroids" (3 Oct., p. 14) quotes Paul Ehrlich as saying that 40% of the net primary production (NPP) on Earth is consumed directly or indirectly by the human population. I find this unbelievable.

As an example, our research at Public Service Electric and Gas Company indicates that, with respect to annual tomato production per acre of greenhouse, yields of almost 300,000 pounds per acre are now being approached. If we assumed a U.S. popula-

tion of approximately 240 million people, and if the only food available to them were tomatoes, with each person consuming 16 pounds of tomatoes a day (1 pound of tomatoes equals 100 calories) for 365 days a year, per capita consumption would be 5840 pounds per year per person; this leads to an estimate that the United States would need $16 \text{ pounds} \times 240 \times 10^6 \text{ people} \times 365 \text{ days a year}$, which equals 1401.6×10^9 or 1,401,600,000,000 pounds of tomatoes a year to feed its population. (Note that this would not be the most balanced diet.)

If the annual tomato consumption in this case (1401.6×10^9 pounds per year) were divided by the acreage production of tomatoes (300,000 pounds per acre per year), only 4,672,000 acres of greenhouses would be required to feed the entire population of the United States. The area of the U.S. land mass is approximately 3,920,000 square miles. If we figure that one-third of the continental land mass is tillable, there are 836,268,800 tillable land acres. If we divide 4,672,000 acres by 836,268,800 acres, only 0.56% of the U.S. tillable land would be required to keep the U.S. population supplied on a mono-diet of tomatoes all year long. The above percentage (0.56%) should be increased slightly, since about 25% of the total light in this particular environment on an annual basis is supplied by artificial, supplemental photosynthetic lighting.

I therefore suspect that the estimate of 40% consumption of NPP by the human population is exaggerated by several orders of magnitude.

BRUCE L. GODFRIAUX
Public Service Electric and Gas Company,
80 Park Plaza,
Newark, NJ 07101

Forest Restoration in Costa Rica

I would like to polish Constance Holden's excellent coverage of the tropical dry forest restoration project in northwestern Costa Rica (News & Comment, 14 Nov., p. 809). Holden reports that "the forested area of Costa Rica has shrunk from 20 to 2% in the past two decades," but in fact it is the *dry* forest area that has so shrunk in area; overall, Costa Rica has an excellent conservation record, with approximately 20% of its area in explicitly conserved national parks and reserves.

Holden says that local wild animal populations "might be jacked up to commercially exploitable levels." Such an action is being taken in some wildlife management projects

in the Neotropics, but is not planned for Guanacaste National Park (GNP). Rather, GNP anticipates that wild animal populations may be sufficiently robust that some individuals may be removed to use for restocking in forest restoration projects elsewhere.

The animal figured is a tapir rather than a peccary.

DANIEL H. JANZEN
Department of Biology,
University of Pennsylvania,
Philadelphia, PA 19104-6018

Rejection and Revolution

After adopting the relatively new policy of swiftly rejecting 60% of all manuscripts, *Science* offers the authors of such manuscripts the comforting thought (Editorial, 18 Jan. 1985, p. 249) that on that particular week they competed with someone of the caliber of Darwin. This flippant attitude assumes that scientists of Darwin's caliber arise about once a week.

The new policy has one disadvantage that totally outweighs any advantages. Those

who make the initial decisions are bound to be recognized members of the scientific establishment, who according to Kuhn (1) are usually the last to recognize the value of a justified revolutionary stand in their scientific field. This is bound to turn totally disastrous what had been almost totally disastrous before, that is, the attitude of *Science* toward the authors of manuscripts that support such a revolutionary stand. Contrary to the advice offered by *Science*, the authors of such manuscripts can, however, find solace in the knowledge that the manuscripts of a scientific revolutionary like Darwin would have certainly been swiftly rejected by *Science* with no right of appeal.

R. ROSIN
126 West 83 Street,
New York, NY 10024

REFERENCES

1. T. Kuhn, *The Structure of Scientific Revolutions* (Univ. of Chicago Press, Chicago, 1970).

Response: In Darwin's day there were so few scientists and so few journals that in most cases issues were printed only when enough manuscripts had been received, not on regular dates. With 90% or more of all

the scientists who have ever lived being alive today, the volume of publications is totally different and the number of young Darwins far greater. Our sending back 60% of submitted manuscripts rapidly is a trade-off for faster decision-making at all levels; this is understood and the procedure has been given a generally favorable reception. We never treat manuscripts flippantly and our staff and reviewers are constantly encouraging publication of innovative research findings. I do not doubt for a moment that we will make an occasional mistake, but so far we do not know of any seminal paper that we have refused nor do we have any indication that well-known scientists are not as receptive to innovative ideas as their less famous colleagues.

—DANIEL E. KOSHLAND, JR.


Erratum: In the Research News article by Gina Kolata "Maleness pinpointed on Y chromosome" (28 Nov., p. 1076), it is said that gametes always induce steroids. In fact, steroids can be produced without gametes. What is constant is that gametes can never be produced without steroids.

Erratum: In the caption for the figure on page 939 with the article "Debate about epilepsy: What initiates seizures?" by Deborah M. Barnes (Research News, 21 Nov., p. 938), the second sentence should have read "Middle trace shows electrical activity recorded from the space outside a neuron and bottom trace shows activity inside a single neuron."

Newport's StableTop desktop isolator offers you...

- More comfort
- More productivity
- Superb vibration protection
- Quick set up in cramped spaces

Send for our full color
brochure about this
exciting new product.



714/963-9811
18235 Mt. Baldy Cir.
Fountain Valley, CA 92708
Europe: Newport GmbH Ph. 06151-26116
U.K.: Newport Ltd. Ph. 05827-69995

**Newport
Corporation**

Circle No. 51 on Readers' Service Card

AUTOMATE MEASUREMENT ON YOUR IBM PC




Photo Courtesy/Edward Jones, M.D.

New digitizing tablet with
Sigma-Scan™ measurement
software. \$1195

Cat #3011 — 12" x 12" system

Resolution of .025 mm, accuracy of at least .25 mm. Comes with state-of-the-art software for area, linear, perimeter, length of curvy line, and angular measurements. X, Y point or stream digitizing. Descriptive statistics. Transfer data to other programs in standard ASCII or DIF format.

Call or write today for more information.

JANDEL SCIENTIFIC
MICROCOMPUTER TOOLS FOR THE SCIENTIST
2656 Bridgeway, Sausalito, CA 94965
800-874-1888

(In Calif. call 415-331-3022)

Circle No. 99 on Readers' Service Card