

an "extreme position," as Miller describes. I also disagree with Miller that the editorial board "was responsible for creating the manuscript."

It should be noted that the Library of Congress has now changed the credits on the volumes such that I am listed, correctly in my view, as the first name. However, I was notified by Lippincott on 15 April that I have been terminated as editor.

Michels is quoted in the 24 April issue of the *Duke Chronicle* as saying in regard to the credit given: "Everyone who has looked at the agreement who is competent, including the entire editorial board, thought it was fine." Since a federal judge and I do not concur, is this a reflection on our competence?

JESSE O. CAVENAR, JR.  
Department of Psychiatry,  
Duke University,  
Durham, NC 27705

### Stereo Viewing

In his letter of 6 February (p. 623) Donn C. Young explained how he viewed stereo. I would like to add my method, which may be easier.

Go to a toy store and buy a pair of binoculars (actually field glasses) for about \$3. Remove both eyepieces and the platform on which they are attached. The objectives on my unit are 35 millimeters in diameter; the overall length of the tubes are 9.5 centimeters, and these dimensions may be somewhat critical. It is good if they also swivel. Now with the objectives closest to your eyes and open end near stereo images, it takes only a moment to find the right distance. Beautiful stereo will pop right out at you. Try it, it really works.

PETER T. B. RUNK  
679 Eastbury Drive,  
Escondido, CA 92027

**Erratum:** In Table 1 (p. 897) of the report "Recombinant interferon enhances monoclonal antibody-targeting of carcinoma lesions in vivo" by John W. Greiner *et al.* (20 Feb., p. 895), the first value under the heading "Plasma Hu-IFN- $\alpha$ A level (antiviral unit/ml)" should have been <30.

**Erratum:** Dalesbred sheep, which were tested by K. M. Kendrick and B. A. Baldwin (Reports, 24 Apr., p. 448), are a horned breed. They do not lack horns, as stated in This Week in Science (24 Apr., p. 371).

**Erratum:** In Eliot Marshall's article "California's debate on carcinogens" (News & Comment, 20 Mar., p. 1459), the last sentence of the third paragraph, which refers to fines to be imposed under Proposition 65, should have read, "Companies found guilty of violating it will be charged \$2500 a day and legal costs." The article gave the incorrect figure of "\$25,000 a day."

# Science

## Posters

The following posters of Science covers are available:

9 February 1979, Locomotive in China;  
27 July 1979, Lightning over St. Louis;  
29 July 1983, Cheetah and cub;  
2 December 1983, Snowshoe hare;  
23 December 1983, Cathedral window/DNA molecule;  
26 September 1986, Neurons in motor cortex;  
28 November 1986, Byrd Glacier, Antarctica  
Combination of space covers in scroll format: 1 June 1979, 23 November 1979, 10 April 1981.

Price is \$5.50 each (prepaid)

Write to AAAS Department POST,  
1333 H Street, NW, Washington,  
D.C. 20005.

## ACCESS TO MANUFACTURERS

Serving the clinical laboratory market. Reach all of them by exhibiting in the "OEM" section of the world's largest clinical laboratory exposition, The American Association for Clinical Chemistry National Meeting, July 21-23, 1987, San Francisco. Over 700 booths, 250 companies, and 200 new product introductions. OEM space is limited. Telephone Scherago/OEM Plus in New York at: 212-730-1050. Telex: 968082.

Circle No. 89 on Readers' Service Card

## LIPOSOMES

REPRODUCIBLE — HOMOGENEOUS —  
EXTREMELY RAPID


NEW

Our new commercial LIPOSOMAT guarantees the reproducible preparation of uniformly sized unilamellar or oligolamellar liposomes of controllable size with very high encapsulation efficiency for both, hydrophilic and/or lipophilic agents as drugs, enzymes etc.

Liposome size is selectable between 25 and 600 nm diameter. Formation of the liposomes is achieved within a few minutes! Lipid loss is 1-2% or

less depending on the nature of the lipid. Sample volume is normally 5-10 ml or up to 200 ml if used in combination with an additional instrument. Smaller volumes are prepared by our smaller liposome instruments, industrial quantities can be produced with our pilot plant.

The LIPOSOMAT is used especially for studying of drug/protein/cell-liposome interactions in vitro and in vivo, for membrane reconstitution experiments, and for the transfer of macromolecules e.g. genetic material into cells in gene technology and biotechnology.



**DIANDERM**  
Geräte  
DIANDERM Geräte  
PO Box 126  
D-8000 MÜNCHEN 65  
FRG

Circle No. 127 on Readers' Service Card