Science

17 February 1989 Volume 243 Number 4893

American Association for the Advancement of Science Science serves its readers as a forum for the presentation and discussion of important issues related to the advancement of science, including the presentation of minority or conflicting points of view, rather than by publishing only material on which a consensus has been reached. Accordingly, all articles published in *Science*—including editorials, news and comment, and book reviews—are signed and reflect the individual views of the authors and not official points of view adopted by the AAAS or the institutions with which the authors are affiliated.

Publisher: Philip H. Abelson, *Acting* **Editor:** Daniel E. Koshland, Jr.

Deputy Editor: John I. Brauman (*Physical Sciences*)

EDITORIAL STAFF

Managing Editor: Patricia A. Morgan Assistant Managing Editor: Nancy J. Hartnagel Senior Editor: Eleanore Butz Associate Editors: Keith W. Brocklehurst, Martha Coleman, R. Brooks Hanson, Barbara Jasny, Katrina L. Kelner, Edith Meyers, Linda J. Miller, Phillip D. Szuromi, David F. Voss Letters Editor: Christine Gilbert Book Reviews: Katherine Livingston, *editor* This Week in Science: Ruth Levy Guyer Contributing Editor: Lawrence I. Grossman Chief Production Editor: Ellen E. Murphy Editing Department: Lois Schmitt, *head*; Mary McDaniel, Patricia L. Moe, Barbara E. Patterson Copy Desk: Joi S. Granger, Jane Hurd, MaryBeth Shartle, Beverly Shields Production Manager: Karen Schools Colson Assistant Production: Holly Bishop, James J. Olivarri, Yolanda M. Rook Covers Editor: Grayce Finger Manuscript Systems Analyst: William Carter

NEWS STAFF

News Editor: Barbara J. Culliton Deputy News Editors: Roger Lewin, Colin Norman News and Comment/Research News: William Booth, Gregory Byrne, Mark H. Crawford, Constance Holden, Richard A. Kerr, Eliot Marshall, Jean L. Marx, Robert Pool, Leslie Roberts, Marjorie Sun, M. Mitchell Waldrop European Correspondent: David Dickson Contributing Writer: John Walsh

BUSINESS STAFF

Circulation Director: John G. Colson Fulfiliment Manager: Ann Ragland Business Staff Manager: Deborah Rivera-Wienhold Classified Advertising Supervisor: Karen Morgenstern Gulde to Biotechnology Products and Instruments: Shauna S. Roberts

ADVERTISING REPRESENTATIVES Director: Earl J. Scherago Traffic Manager: Donna Rivera

Traffic Manager: Donna Rivera Traffic Manager (Recruitment): Gwen Canter Advertising Sales Manager: Richard L. Charles Marketing Manager: Herbert L. Burklund Employment Sales Manager: Edward C. Keller Sales: New York, NY 10036; J. Kevin Henebry, 1515 Broadway (212-730-1050); Scotch Plains, NJ 07076; C. Richard Callis, 12 Unami Lane (201-889-4873); Chicago, IL 60914; Jack Ryan, 525 W. Higgins Rd. (312-885-8675); San Jose, CA 95112; Bob Brindley, 310 S. 16th St. (408-998-4690); Dorset, VT 05251; Fred W. Dieffenbach, Kent Hill Rd. (802-867-5581); Damascus, MD 20872; Rick Sommer, 11318 Kings Valley Dr. (301-972-9270); U.K., Europe: Nick Jones, +44(0647)52918; Telex 42513; FAX (0647) 52053.

Information for contributors appears on page XI of the 23 December 1988 issue. Editorial correspondence, including requests for permission to reprint and reprint orders, should be sent to 1333 H Street, NW, Washington, DC 20005. Telephone: 202-326-6500.

The Arctic: A Key to World Climate

The Arctic is part of a great global heat engine. Changes in the arctic atmosphere, ocean, sea ice, and permafrost are early precursors to climate change elsewhere. In the past, those changes have been drastic. Only 18,000 years ago, virtually all of Canada and some of the United States were covered by a thick layer of ice.

At the recent AAAS Annual Meeting, a symposium brought together some of the leading research scientists active in studies of resources and climate.* In one of the sessions, speakers reviewed information about the evolution of the arctic climate.

A principal impression that could be drawn from the symposium was that the current hypothesis concerning effects of greenhouse gases on arctic behavior is probably simplistic and may be quite wrong. The public has been told repeatedly that a result of increased greenhouse gases would be a substantial rise in sea level due to melting of polar ice. A related statement frequently made is that the increase in polar temperatures would be substantially greater than those of global averages. For example, one estimate is that an average rise of 2° C would be accompanied by a 10° C increase in the Arctic. That estimate may or may not adequately take into account climatic feedback mechanisms. At the symposium this point was raised implicitly by John T. Andrews, who stated that the Greenland Ice Sheet and the Laurentide Ice Sheet advanced during a period of warm water influx into Baffin Bay at the end of the last interglaciation.

The vapor pressure of water is quite sensitive to temperature. Condensing moisture in the form of snow provides a surface cover that highly reflects solar energy leading to a regional cooling. The albedo (reflectivity) of ordinary soil is about 0.1. The albedo of snow is about 0.8. At present, some of the arctic land areas that have averaged annual temperatures of about -14° C receive only 10 centimeters of total H₂O per year. Most of the time the surface is bare and is a good absorber of solar heat. Were more precipitation to occur, the total heat absorbed by the surface would decrease. With greater moisture in the air, there would be more clouds. The net effects of these are controversial. Some say that more clouds would reflect more energy away from the earth. Others point out that added moisture would enhance a greenhouse effect in the Arctic. In any event, the factors controlling arctic climate are complex.

One of the obstacles to confidence in predicting the future of the arctic climate is an imperfect knowledge of the past. We know that 70 million years ago, the climate was mild and the Arctic Ocean was ice-free. Sediments formed about 5 million years ago contained glacially related materials. We know little about what happened in the long interval, and knowledge concerning more recent events is sketchy. No long cores have been obtained from the Arctic Ocean. The impediment is the perennial ice sheet that covers most of that ocean. The thickness is usually about 3 to 4 meters, and the sheet tends to keep moving. At the geographic North Pole, the depth of the ocean is about 3500 meters.

Our best source of evidence concerning the last million years is found in near-shore sediments, and particularly on fossil-bearing terraces. Molluscan fossils are particularly useful. Different molluscan species have different temperature affinities. In addition, they contain partially hydrolyzed proteins. The degree of racemization of isoleucine is a function of times and temperatures. The shells also contain strontium as a trace element which provides a dating potential. The ratio of ⁸⁷Sr to ⁸⁶Sr in seawater has changed monotonically over the last several million years. For events during the past 40,000 years ¹⁴C dating can be applied. Other types of fossils are being studied and additional dating methods employed. Measurements of ratios of ¹⁸O to ¹⁶O in ice are useful in determining temperatures at which atmospheric moisture was converted into ice. Oxygen isotope ratios of shells reveal temperatures present during their formation.

The importance of understanding the past, present, and future of the arctic climate requires that support for such activities have a top priority. Desirable efforts include more studies of fossils, an international program of deep drilling in the Arctic, more weather monitoring, and additional satellite surveillance of the polar region.—PHILIP H. ABELSON

Advertising correspondence should be sent to Tenth Floor, 1515 Broadway, New York, NY 10036. Telephone 212-730-1050 or WU Telex 968082 SCHERAGO, or FAX 212-382-3725.

^{*&}quot;The Arctic: A New Key to World Climate and Resources," organized by P. W. Barnes and K. A. Kvenvold of the U.S. Geological Survey, Menlo Park, CA, and held on 19 January 1989 in San Francisco, CA.