

---

# Identification of the Cystic Fibrosis Gene: Cloning and Characterization of Complementary DNA

JOHN R. RIORDAN, JOHANNA M. ROMMENS, BAT-SHEVA KEREM, NOA ALON,  
RICHARD ROZMAHEL, ZBYSZKO GRZELCZAK, JULIAN ZIELENSKI, SI LOK,  
NATASA PLAVSIC, JIA-LING CHOU, MITCHELL L. DRUMM, MICHAEL C. IANNUZZI,  
FRANCIS S. COLLINS, LAP-CHEE TSUI

---

**Overlapping complementary DNA clones were isolated from epithelial cell libraries with a genomic DNA segment containing a portion of the putative cystic fibrosis (CF) locus, which is on chromosome 7. Transcripts, approximately 6500 nucleotides in size, were detectable in the tissues affected in patients with CF. The predicted protein consists of two similar motifs, each with (i) a domain having properties consistent with membrane association and (ii) a domain believed to be involved in ATP (adenosine triphosphate) binding. A deletion of three base pairs that results in the omission of a phenylalanine residue at the center of the first predicted nucleotide-binding domain was detected in CF patients.**

---

**C**YSTIC FIBROSIS (CF) IS AN AUTOSOMAL RECESSIVE GENETIC disorder affecting a number of organs, including the lung airways, pancreas, and sweat glands (1). Abnormally high electrical potential differences have been detected across the epithelial surfaces of the CF respiratory tract, including the trachea and nasal polyps, as well as across the walls of CF sweat gland secretory coils and reabsorptive ducts (2). The basic defect has been associated with decreased chloride ion conductance across the apical membrane of the epithelial cells (3). That the defect also appeared to persist in cultured cells derived from several epithelial tissues suggested that the CF gene is expressed in these cells (4). More recently, patch clamp studies showed that this defect is probably due to a failure of an outwardly rectifying anion channel to respond to phosphorylation by cyclic AMP-dependent protein kinase (protein kinase A) or protein kinase C (5). Although progress has been made in the

isolation of polypeptide components of an epithelial chloride channel that mediates conductance (6), their relation to the kinase-activated pathway and CF has yet to be established, and the basic biochemical defect in CF remains unknown.

Molecular cloning experiments have permitted the isolation of a large, contiguous segment of DNA spanning at least four transcribed sequences from a region thought to contain the CF locus (7). These sequences were initially identified on the basis of their ability to detect conserved sequences in other animal species by DNA hybridization and were subsequently characterized by RNA hybridization experiments, cDNA isolation, and direct DNA sequence analysis (7). Three of the transcribed regions were excluded from being the CF locus by earlier genetic or DNA sequence analyses (7, 8). The fourth one, as shown by genetic analysis (9) and DNA sequencing analysis presented below, corresponds to a portion of the CF gene locus.

**Isolation of cDNA clones.** Two DNA segments (E4.3 and H1.6) that detected cross-species hybridization signals (7) were used as probes to screen cDNA libraries made from several tissues and cell types (10). After screening seven different libraries, one single clone (10-1) was isolated with H1.6 from a cDNA library made from the cultured epithelial cells of the sweat glands of an unaffected (non-CF) individual (10).

DNA sequencing showed that 10-1 contained an insert of 920 base pairs (bp) in size and one potential, long open reading frame (ORF). Since one end of the sequence shared perfect sequence identity with H1.6, it was concluded that the cDNA clone was probably derived from this region. The DNA sequence in common was, however, only 113 bp long (Figs. 1 and 2). This sequence in fact corresponded to the first exon of the putative CF gene. The short sequence overlap thus explained the weak hybridization signals in library screening and our inability to detect transcripts in RNA gel-blot analysis. In addition, the orientation of the transcription unit was tentatively established on the basis of alignment of the genomic DNA sequence with the presumptive ORF of 10-1.

Since the corresponding transcript was estimated to be about 6500 nucleotides in length by RNA gel-blot hybridization experiments, further cDNA library screening was required in order to clone the remainder of the coding region. As a result of several successive screenings with cDNA libraries generated from the colon carcinoma cell line T84, normal and CF sweat gland cells, pancreas,

---

J. R. Riordan, N. Alon, and Z. Grzelczak are in the Department of Biochemistry, Research Institute, The Hospital for Sick Children, Toronto, Ontario M5G 1X8, Canada. J. M. Rommens, B. Kerem, R. Rozmahel, J. Zielenski, S. Lok, N. Plavsic, J.-L. Chou, and L.-C. Tsui are at the Department of Genetics, Research Institute, The Hospital for Sick Children, Toronto, Ontario M5G 1X8, Canada. M. L. Drumm, M. C. Iannuzzi, and F. S. Collins are at the Howard Hughes Medical Institute and Departments of Internal Medicine and Human Genetics, University of Michigan, Ann Arbor, MI 48109. J. R. Riordan is also a member of the Departments of Biochemistry and Clinical Biochemistry, and L.-C. Tsui is also a member of the Departments of Medical Genetics and Medical Biophysics, University of Toronto, Toronto, Ontario M5S 1A8, Canada.

and adult lungs, 18 additional clones were isolated (Fig. 1). DNA sequence analysis revealed that none of these cDNA clones corresponded to the length of the observed transcript, but it was possible to derive a consensus sequence based on overlapping regions. Further cDNA clones corresponding to the 5' and 3' ends of the transcript were derived from 5' and 3' primer-extension experiments (Fig. 1). Together, these clones span about 6.1 kb and contain an ORF capable of encoding a protein of 1480 amino acids (Fig. 2).

It was unusual that most of the cDNA clones isolated here contained sequence insertions at various locations (Fig. 1). While many of these extra sequences corresponded to intron regions reverse-transcribed during the construction of the cDNA, as revealed on alignment with genomic DNA sequences, the identities of several others were uncertain because they did not align with sequences at the corresponding exon-intron junctions, namely, the sequences at the 5' ends of clones 13a and T16-1 and at the 5' and 3' ends of T11, and the insertions between exons 3 and 4 in 13a and between exons 10 and 11 in T16-4.5 (legend to Fig. 1). More puzzling were the sequences corresponding to the reverse complement of exon 6 at the 5' end of 11a and the insertion of a segment of a bacterial transposon in clone C16-1; none of these could be explained by mRNA processing errors.

In that the number of recombinant cDNA clones for the putative CF gene detected in the library screening was much less than would have been expected from the abundance of transcripts estimated from RNA hybridization experiments, it seemed probable that the clones that contained aberrant structures were preferentially retained while the proper clones were lost during propagation. Consistent with this interpretation, poor growth was observed for most of our recombinant clones isolated, regardless of the vector used.

**RNA analysis.** To visualize the transcript of the putative CF gene, we used RNA gel-blot hybridization with the 10-1 cDNA as


the probe (Fig. 3). The analysis revealed a prominent band, about 6.5 kb in size, in T84 cells. Identical results were obtained with other cDNA clones as probes. Similar, strong hybridization signals were also detected in pancreas and primary cultures of cells from nasal polyps, suggesting that the mature mRNA of the putative CF gene is about 6.5 kb. Minor hybridization signals, probably representing degradation products, were detected at the lower size ranges, but they varied between different experiments. On the basis of the hybridization band intensity and comparison with those detected for other transcripts under identical experimental conditions, it was estimated that the putative CF gene transcripts constituted about 0.01 percent of total mRNA in T84 cells.

Additional tissues were analyzed by RNA gel-blot hybridization in an attempt to correlate the expression pattern of the putative CF gene and the pathology of CF. Transcripts, all of identical size, were found in lung, colon, sweat glands (cultured epithelial cells), placenta, liver, and parotid gland, but the signal in these tissues was generally weaker than that detected in the pancreas and nasal polyps (Fig. 3). Intensity varied among different preparations; for example, hybridization in kidney was not detectable in the preparation shown in Fig. 3 but was clearly discernible subsequently. Transcripts were not detected in the brain or adrenal gland, nor in skin fibroblast and lymphoblast cell lines.

Thus, expression of the putative CF gene appeared to occur in many of the tissues examined, with higher levels in those tissues severely affected in CF. While this epithelial tissue-specific expression pattern is in good agreement with the disease pathology, no significant difference was detected in the amount or size of transcripts from CF and control tissues (Fig. 3), consistent with the assumption that CF mutations are subtle changes at the nucleotide level.

**Characterization of cDNA clones.** As indicated above, a contig-

**Fig. 1.** Overlapping cDNA clones aligned with genomic DNA fragments. The cDNA clones are represented by open boxes with exons indicated. The corresponding genomic Eco RI fragments are schematically presented on the bottom, with lengths in kilobases. The hatched boxes denote intron sequences, and stippled boxes represent other sequences as outlined below. The filled box in the lower left is the position of the clone H1.6, which was used to isolate the first cDNA clone 10-1 from a normal (N) sweat gland library (10). The definitive restriction sites used for the alignment of cDNA and genomic fragments are indicated. Clones T6, T6/20, T11, T16-1, T13-1, T16-4.5, T8-B3, and T12a were isolated sequentially from the T84 cell library (10). Clones isolated from the human lung cDNA library (10) are designated with the prefix CDL. CDPJ5 is derived from a pancreas library (10). The CF sweat gland cDNA clones, C16-1 and C1-1/5, together cover all but exon 1 and a portion of the 3' untranslated region. Both clones revealed a 3-bp deletion in exon 10.



Both PA3-5 and TB2-7 are 5' extension clones generated from pancreas and T84 RNA by the anchored PCR technique (12), respectively. THZ-4 is a 3' extension clone obtained from T84 RNA. Both T12a and THZ-4 contain a polyadenylation signal and a poly(A)<sup>+</sup> tail.

1 AATTGGAAGCAAAATGACATCAGCAGGTCAGAGAAAAGGGTTGAGCGGCAGGCCACCCA  
61 GAGTAGTAGGCTTTGGCATTAGGAGCTTGA<sup>5</sup>CCAGACGGCCCTAGCAGGGACCCCGC  
121 GCCCGAGAGACCATGAGAGGTCGCCTCTGGAAAAGGCCAGCGTGTCTCCAAACCTTTT  
181 F<sup>1</sup>W<sup>1</sup>T<sup>1</sup>R<sup>1</sup>P<sup>1</sup>I<sup>1</sup>L<sup>1</sup>R<sup>1</sup>K<sup>1</sup>G<sup>1</sup>Y<sup>1</sup>R<sup>1</sup>Q<sup>1</sup>R<sup>1</sup>L<sup>1</sup>E<sup>1</sup>L<sup>1</sup>S<sup>1</sup>D<sup>1</sup>  
TTCAC<sup>1</sup>CTGGACCAGACCAATTTGAGGAAAGATACAGACAGCGCCTGGAATGTGACAGC  
241 I<sup>1</sup>Y<sup>1</sup>Q<sup>1</sup>I<sup>1</sup>P<sup>1</sup>S<sup>1</sup>V<sup>1</sup>D<sup>1</sup>S<sup>1</sup>A<sup>1</sup>D<sup>1</sup>N<sup>1</sup>L<sup>1</sup>S<sup>1</sup>E<sup>1</sup>K<sup>1</sup>L<sup>1</sup>E<sup>1</sup>H<sup>1</sup>E<sup>1</sup>  
ATATACCAAAATCCCTTCTGTTGATCTGCTGACAATCTATCTGAAAAATTGGAAGAGAA  
301 W<sup>1</sup>D<sup>1</sup>R<sup>1</sup>E<sup>1</sup>L<sup>1</sup>A<sup>1</sup>S<sup>1</sup>K<sup>1</sup>K<sup>1</sup>N<sup>1</sup>P<sup>1</sup>K<sup>1</sup>L<sup>1</sup>I<sup>1</sup>N<sup>1</sup>A<sup>1</sup>L<sup>1</sup>R<sup>1</sup>R<sup>1</sup>C<sup>1</sup>  
TGGATAGAGAGCTGGCTTCAAGAAAAATCTTAACTCATTAATGCCCTTCGGCGATGT  
361 F<sup>1</sup>F<sup>1</sup>W<sup>1</sup>R<sup>1</sup>F<sup>1</sup>M<sup>1</sup>F<sup>1</sup>Y<sup>1</sup>G<sup>1</sup>I<sup>1</sup>F<sup>1</sup>L<sup>1</sup>Y<sup>1</sup>L<sup>1</sup>G<sup>1</sup>E<sup>1</sup>V<sup>1</sup>T<sup>1</sup>K<sup>1</sup>A<sup>1</sup>  
TTTCTGGAGATTGATGTTCTTGAATCTTTTATATTTAGGGAAGTCCACAAAGCA  
421 V<sup>1</sup>O<sup>1</sup>P<sup>1</sup>L<sup>1</sup>L<sup>1</sup>L<sup>1</sup>G<sup>1</sup>R<sup>1</sup>I<sup>1</sup>A<sup>1</sup>S<sup>1</sup>Y<sup>1</sup>D<sup>1</sup>P<sup>1</sup>D<sup>1</sup>N<sup>1</sup>K<sup>1</sup>E<sup>1</sup>E<sup>1</sup>  
GTACAGCCTCTTACTGGGAAGAAATCATGCTCTTATGACCCGGATTAACAGGAGGAA  
481 R<sup>1</sup>S<sup>1</sup>I<sup>1</sup>A<sup>1</sup>I<sup>1</sup>Y<sup>1</sup>L<sup>1</sup>G<sup>1</sup>I<sup>1</sup>G<sup>1</sup>L<sup>1</sup>C<sup>1</sup>L<sup>1</sup>L<sup>1</sup>F<sup>1</sup>I<sup>1</sup>V<sup>1</sup>R<sup>1</sup>T<sup>1</sup>L<sup>1</sup>  
CGCTCTATCGCATTATAGGATAGGCTTATGCTTCTTATGAGGACTG  
541 L<sup>1</sup>L<sup>1</sup>H<sup>1</sup>P<sup>1</sup>A<sup>1</sup>I<sup>1</sup>F<sup>1</sup>G<sup>1</sup>L<sup>1</sup>H<sup>1</sup>E<sup>1</sup>I<sup>1</sup>G<sup>1</sup>M<sup>1</sup>Q<sup>1</sup>M<sup>1</sup>R<sup>1</sup>I<sup>1</sup>A<sup>1</sup>M<sup>1</sup>  
CTCTACACCCAGCATTGTTGGCCTTCAATCACTGGAATGCAGATGAGAAATGACTATG  
601 F<sup>1</sup>S<sup>1</sup>L<sup>1</sup>I<sup>1</sup>Y<sup>1</sup>K<sup>1</sup>K<sup>1</sup>I<sup>1</sup>T<sup>1</sup>L<sup>1</sup>K<sup>1</sup>L<sup>1</sup>S<sup>1</sup>S<sup>1</sup>R<sup>1</sup>V<sup>1</sup>L<sup>1</sup>D<sup>1</sup>K<sup>1</sup>I<sup>1</sup>S<sup>1</sup>  
TTTGTATGATTTATAAGAGACTTAAAGCTTCAAGCCGCTTCTAGATAAAATAAGT  
661 I<sup>1</sup>G<sup>1</sup>Q<sup>1</sup>L<sup>1</sup>V<sup>1</sup>S<sup>1</sup>L<sup>1</sup>S<sup>1</sup>N<sup>1</sup>N<sup>1</sup>L<sup>1</sup>N<sup>1</sup>K<sup>1</sup>F<sup>1</sup>D<sup>1</sup>E<sup>1</sup>G<sup>1</sup>L<sup>1</sup>A<sup>1</sup>  
ATTGACAACTTGTAGTCTCTTCCCAACAACCTGAAACAAATTTGATGAAAGACTTGA  
721 L<sup>1</sup>A<sup>1</sup>H<sup>1</sup>F<sup>1</sup>V<sup>1</sup>W<sup>1</sup>I<sup>1</sup>A<sup>1</sup>P<sup>1</sup>L<sup>1</sup>O<sup>1</sup>V<sup>1</sup>A<sup>1</sup>L<sup>1</sup>L<sup>1</sup>M<sup>1</sup>G<sup>1</sup>L<sup>1</sup>I<sup>1</sup>W<sup>1</sup>  
TTGGCATTTCGTGGATTCCTTTGCAAGTGGCCTCTTCAATCTG  
781 E<sup>1</sup>L<sup>1</sup>L<sup>1</sup>Q<sup>1</sup>A<sup>1</sup>S<sup>1</sup>A<sup>1</sup>F<sup>1</sup>C<sup>1</sup>G<sup>1</sup>L<sup>1</sup>G<sup>1</sup>F<sup>1</sup>L<sup>1</sup>I<sup>1</sup>V<sup>1</sup>L<sup>1</sup>A<sup>1</sup>L<sup>1</sup>F<sup>1</sup>  
GAGTTGTACAGCGCTGCTTCTGTTGAGCTTGGTTTCTGATAGTCTTGCCTTTT  
841 O<sup>1</sup>A<sup>1</sup>G<sup>1</sup>L<sup>1</sup>G<sup>1</sup>R<sup>1</sup>M<sup>1</sup>M<sup>1</sup>M<sup>1</sup>K<sup>1</sup>Y<sup>1</sup>R<sup>1</sup>D<sup>1</sup>Q<sup>1</sup>R<sup>1</sup>A<sup>1</sup>G<sup>1</sup>K<sup>1</sup>I<sup>1</sup>S<sup>1</sup>  
CAGGCTGGGCTAGGAGAAATGATGAGTACAGAGATCAGAGCTGGGAGATCAGT  
901 E<sup>1</sup>R<sup>1</sup>L<sup>1</sup>V<sup>1</sup>I<sup>1</sup>T<sup>1</sup>S<sup>1</sup>E<sup>1</sup>M<sup>1</sup>I<sup>1</sup>E<sup>1</sup>N<sup>1</sup>I<sup>1</sup>Q<sup>1</sup>S<sup>1</sup>V<sup>1</sup>K<sup>1</sup>A<sup>1</sup>Y<sup>1</sup>C<sup>1</sup>  
GAAAGACTGTGATCTCAGAAATGATGAAAATATCCAATCTGTAAGGCATCTGC  
961 W<sup>1</sup>E<sup>1</sup>E<sup>1</sup>A<sup>1</sup>M<sup>1</sup>E<sup>1</sup>K<sup>1</sup>M<sup>1</sup>I<sup>1</sup>E<sup>1</sup>N<sup>1</sup>L<sup>1</sup>R<sup>1</sup>Q<sup>1</sup>T<sup>1</sup>E<sup>1</sup>L<sup>1</sup>K<sup>1</sup>L<sup>1</sup>T<sup>1</sup>  
TGGGAAGAACAAATGGAAAAATGATGAAAATTAAGACAAACAGAACTGAAACTGACT  
1021 R<sup>1</sup>K<sup>1</sup>A<sup>1</sup>A<sup>1</sup>Y<sup>1</sup>V<sup>1</sup>R<sup>1</sup>Y<sup>1</sup>F<sup>1</sup>N<sup>1</sup>S<sup>1</sup>S<sup>1</sup>A<sup>1</sup>F<sup>1</sup>F<sup>1</sup>F<sup>1</sup>S<sup>1</sup>G<sup>1</sup>G<sup>1</sup>T<sup>1</sup>  
CGAAGGACCCATGTGAGACTTCAATAGCTCAGCCCTCTTCTCAGGTTCTTT  
1081 V<sup>1</sup>V<sup>1</sup>F<sup>1</sup>L<sup>1</sup>S<sup>1</sup>V<sup>1</sup>L<sup>1</sup>P<sup>1</sup>Y<sup>1</sup>A<sup>1</sup>L<sup>1</sup>I<sup>1</sup>K<sup>1</sup>G<sup>1</sup>I<sup>1</sup>L<sup>1</sup>R<sup>1</sup>K<sup>1</sup>I<sup>1</sup>  
GTGGTGTTTTATCTGTGCTTCCCTATGCACTAATCAAGAAATCATCTCCGAAAAATA  
1141 F<sup>1</sup>T<sup>1</sup>C<sup>1</sup>T<sup>1</sup>I<sup>1</sup>S<sup>1</sup>F<sup>1</sup>C<sup>1</sup>I<sup>1</sup>V<sup>1</sup>L<sup>1</sup>R<sup>1</sup>M<sup>1</sup>A<sup>1</sup>V<sup>1</sup>T<sup>1</sup>R<sup>1</sup>Q<sup>1</sup>F<sup>1</sup>W<sup>1</sup>  
TTCACCATTCTCATCTGATGTTCTGCGCAATGGCGTCACTCGGACTTCCCTGG  
1201 A<sup>1</sup>V<sup>1</sup>Q<sup>1</sup>T<sup>1</sup>W<sup>1</sup>Y<sup>1</sup>D<sup>1</sup>S<sup>1</sup>L<sup>1</sup>G<sup>1</sup>A<sup>1</sup>I<sup>1</sup>N<sup>1</sup>K<sup>1</sup>I<sup>1</sup>Q<sup>1</sup>D<sup>1</sup>F<sup>1</sup>L<sup>1</sup>Q<sup>1</sup>  
GCTGTACAAACATGGTATGACTTCTTGGGCAATAAACAATAACAGATTTCTTACAA  
1261 K<sup>1</sup>Q<sup>1</sup>E<sup>1</sup>Y<sup>1</sup>K<sup>1</sup>T<sup>1</sup>L<sup>1</sup>E<sup>1</sup>Y<sup>1</sup>N<sup>1</sup>L<sup>1</sup>T<sup>1</sup>T<sup>1</sup>E<sup>1</sup>V<sup>1</sup>M<sup>1</sup>E<sup>1</sup>N<sup>1</sup>  
AAGCAAGAAATAAAGACTTGAATATAACTTAAGACTACAGAAAGTAGTGGAGAA  
1321 V<sup>1</sup>T<sup>1</sup>A<sup>1</sup>F<sup>1</sup>W<sup>1</sup>E<sup>1</sup>E<sup>1</sup>G<sup>1</sup>F<sup>1</sup>G<sup>1</sup>E<sup>1</sup>L<sup>1</sup>F<sup>1</sup>E<sup>1</sup>K<sup>1</sup>A<sup>1</sup>K<sup>1</sup>Q<sup>1</sup>N<sup>1</sup>N<sup>1</sup>  
GTAACAGCCTTCTGGGAGGAGGATTTGGGAAATTTTGGAGAAACAAAACAAAT  
1381 N<sup>1</sup>N<sup>1</sup>R<sup>1</sup>K<sup>1</sup>T<sup>1</sup>D<sup>1</sup>S<sup>1</sup>L<sup>1</sup>F<sup>1</sup>F<sup>1</sup>S<sup>1</sup>L<sup>1</sup>L<sup>1</sup>  
AACAAATAGAAAATCTTAATGGTATGACAGCCTCTTCTCAGTAATTTCTACTTCT  
1441 G<sup>1</sup>T<sup>1</sup>P<sup>1</sup>V<sup>1</sup>L<sup>1</sup>K<sup>1</sup>D<sup>1</sup>I<sup>1</sup>N<sup>1</sup>F<sup>1</sup>K<sup>1</sup>I<sup>1</sup>E<sup>1</sup>R<sup>1</sup>G<sup>1</sup>O<sup>1</sup>L<sup>1</sup>L<sup>1</sup>A<sup>1</sup>V<sup>1</sup>  
GGTACTCTGCTGAAAGATTAATTTCAAGATAGAAAGAGACAGTGTGGCGGTT  
1501 A<sup>1</sup>G<sup>1</sup>S<sup>1</sup>T<sup>1</sup>G<sup>1</sup>A<sup>1</sup>G<sup>1</sup>K<sup>1</sup>T<sup>1</sup>S<sup>1</sup>L<sup>1</sup>L<sup>1</sup>M<sup>1</sup>M<sup>1</sup>I<sup>1</sup>G<sup>1</sup>E<sup>1</sup>L<sup>1</sup>E<sup>1</sup>  
GCTGGATCCACTGGAGCAGCAAACTTCACTTAAATGATGATTATGGGAGAACTGGAG  
1561 P<sup>1</sup>S<sup>1</sup>E<sup>1</sup>G<sup>1</sup>K<sup>1</sup>I<sup>1</sup>K<sup>1</sup>H<sup>1</sup>S<sup>1</sup>G<sup>1</sup>R<sup>1</sup>I<sup>1</sup>S<sup>1</sup>F<sup>1</sup>C<sup>1</sup>S<sup>1</sup>O<sup>1</sup>F<sup>1</sup>W<sup>1</sup>T<sup>1</sup>  
CCTTCAGAGGTAATAAATAGCAGCAGTGGAAATTTCAATCTGTTCTGATTTCTCGG  
1621 I<sup>1</sup>M<sup>1</sup>P<sup>1</sup>G<sup>1</sup>T<sup>1</sup>I<sup>1</sup>K<sup>1</sup>E<sup>1</sup>N<sup>1</sup>I<sup>1</sup>F<sup>1</sup>G<sup>1</sup>V<sup>1</sup>S<sup>1</sup>Y<sup>1</sup>D<sup>1</sup>E<sup>1</sup>Y<sup>1</sup>R<sup>1</sup>  
ATTATGCTGGCACCATAAAGAAATATCATCTTGGTGTCTTCTATGATGAATATAGA  
1681 Y<sup>1</sup>R<sup>1</sup>S<sup>1</sup>V<sup>1</sup>I<sup>1</sup>K<sup>1</sup>A<sup>1</sup>C<sup>1</sup>O<sup>1</sup>L<sup>1</sup>E<sup>1</sup>D<sup>1</sup>I<sup>1</sup>S<sup>1</sup>K<sup>1</sup>F<sup>1</sup>A<sup>1</sup>E<sup>1</sup>K<sup>1</sup>  
TACAGAAAGCTCATCAAGCATGCCACTAGAAAGGACATCTCCAAGTTTCAGAGAAA  
1741 D<sup>1</sup>N<sup>1</sup>I<sup>1</sup>V<sup>1</sup>L<sup>1</sup>G<sup>1</sup>E<sup>1</sup>G<sup>1</sup>G<sup>1</sup>I<sup>1</sup>T<sup>1</sup>L<sup>1</sup>S<sup>1</sup>G<sup>1</sup>G<sup>1</sup>O<sup>1</sup>R<sup>1</sup>A<sup>1</sup>R<sup>1</sup>I<sup>1</sup>  
GACAAATAGTCTTGGAGAGGTTGAATCAGACTGAGTGGAGGTCACAGCAAGAAAT  
1801 S<sup>1</sup>L<sup>1</sup>A<sup>1</sup>R<sup>1</sup>A<sup>1</sup>V<sup>1</sup>Y<sup>1</sup>K<sup>1</sup>D<sup>1</sup>A<sup>1</sup>D<sup>1</sup>L<sup>1</sup>Y<sup>1</sup>L<sup>1</sup>D<sup>1</sup>S<sup>1</sup>P<sup>1</sup>F<sup>1</sup>G<sup>1</sup>  
TCTTAGCAAGGAGCTATACAAAGATGCTGATTTGATTTAATAGACTCTCTTTTGA  
1861 Y<sup>1</sup>L<sup>1</sup>D<sup>1</sup>V<sup>1</sup>L<sup>1</sup>T<sup>1</sup>E<sup>1</sup>K<sup>1</sup>E<sup>1</sup>I<sup>1</sup>F<sup>1</sup>E<sup>1</sup>S<sup>1</sup>C<sup>1</sup>V<sup>1</sup>C<sup>1</sup>K<sup>1</sup>L<sup>1</sup>M<sup>1</sup>A<sup>1</sup>  
TACCTAGATGTTTAAACAGAAAAGAAATTTGAAAGCTGTGCTGTAAACTGATGGCT  
1921 N<sup>1</sup>K<sup>1</sup>T<sup>1</sup>R<sup>1</sup>I<sup>1</sup>L<sup>1</sup>V<sup>1</sup>T<sup>1</sup>S<sup>1</sup>K<sup>1</sup>M<sup>1</sup>E<sup>1</sup>H<sup>1</sup>L<sup>1</sup>K<sup>1</sup>K<sup>1</sup>A<sup>1</sup>D<sup>1</sup>K<sup>1</sup>I<sup>1</sup>  
AACAAAATAGGATTTGGTCACTTCTAAAATGGAACATTTAAAGAAAGCTGACAAAATA  
1981 L<sup>1</sup>I<sup>1</sup>L<sup>1</sup>N<sup>1</sup>E<sup>1</sup>G<sup>1</sup>S<sup>1</sup>Y<sup>1</sup>F<sup>1</sup>Y<sup>1</sup>G<sup>1</sup>T<sup>1</sup>F<sup>1</sup>S<sup>1</sup>E<sup>1</sup>L<sup>1</sup>O<sup>1</sup>N<sup>1</sup>L<sup>1</sup>  
TAAATTTGAATGAAGGTAGCAGCTATTTTATGGGACATTTTCAGAACTCCAATCTA  
2041 Q<sup>1</sup>P<sup>1</sup>D<sup>1</sup>F<sup>1</sup>S<sup>1</sup>K<sup>1</sup>L<sup>1</sup>M<sup>1</sup>G<sup>1</sup>C<sup>1</sup>D<sup>1</sup>S<sup>1</sup>F<sup>1</sup>D<sup>1</sup>Q<sup>1</sup>F<sup>1</sup>S<sup>1</sup>A<sup>1</sup>E<sup>1</sup>  
CAGCCAGACTTAGCTCAAACTCATGGGATGTATCTTCGACCAATTTAGTGAGAA  
2101 R<sup>1</sup>R<sup>1</sup>S<sup>1</sup>I<sup>1</sup>L<sup>1</sup>T<sup>1</sup>E<sup>1</sup>T<sup>1</sup>L<sup>1</sup>H<sup>1</sup>R<sup>1</sup>F<sup>1</sup>S<sup>1</sup>L<sup>1</sup>E<sup>1</sup>G<sup>1</sup>D<sup>1</sup>A<sup>1</sup>P<sup>1</sup>  
AGAAGAAATCAATCTAATGAGACTTACACCTTTCATAGAAAGGADGCTCTCT  
2161 V<sup>1</sup>S<sup>1</sup>W<sup>1</sup>T<sup>1</sup>E<sup>1</sup>T<sup>1</sup>K<sup>1</sup>K<sup>1</sup>O<sup>1</sup>S<sup>1</sup>F<sup>1</sup>K<sup>1</sup>O<sup>1</sup>T<sup>1</sup>G<sup>1</sup>E<sup>1</sup>F<sup>1</sup>G<sup>1</sup>E<sup>1</sup>K<sup>1</sup>  
GTCCTCGGACAGAAAACAAAACAACTTTTAAACAGACTGGAGTTTGGGAAAAA  
2221 R<sup>1</sup>K<sup>1</sup>N<sup>1</sup>S<sup>1</sup>I<sup>1</sup>L<sup>1</sup>N<sup>1</sup>P<sup>1</sup>I<sup>1</sup>N<sup>1</sup>S<sup>1</sup>I<sup>1</sup>R<sup>1</sup>K<sup>1</sup>F<sup>1</sup>S<sup>1</sup>I<sup>1</sup>V<sup>1</sup>Q<sup>1</sup>K<sup>1</sup>  
AGGAAATTTCTATTCTCAATCCAATCAACTATACGAAAAATTTCCATTGTGCAAAAG

2281 T P L Q M N G I E E D S D E P L E R R L 736  
ACTCCCTTACAAATGAATGGCATCGAAGAGGATCTGATGAGCCTTAGAGAGAGGCGT  
2341 S L V P D S E Q G E A I L P R I S V I S 756  
TCTTAGTACCAGATTCTGAGCAGGAGAGGCGTCTGCTGCATCAGCCTGATCAGC  
2401 T G P T L Q A R R R Q S V L N L M T H S 776  
ACTGGCCCCACGCTTACAGCAGAGAGGAGGAGTCTGCTCAACCTGATGACACACTA  
2461 V N Q G Q N I H R K A T A S T R K V S L 796  
GTTAACCAAGGTCAGAACATTCACCAGAAAGACACAGCATCCACAGAAAAGTGCAGT  
2521 A P Q A N L T E L D I Y S R R L S Q E T 816  
GCCCTCAGGCAAACTGACTGAAGTATATATTAAGAAAGTTATCTCAAGAAACT  
2581 G L E I S E E I N E E D L K E C L F D D 836  
GGCTTGGAAATAAGTGAAGAAATTAACGAAGAAAGACTTAAAGCAGTGCCTTTTGTATGAT  
2641 M E S I P A V T T W N T Y L R Y I T V H 856  
ATGGAGAGCATACCAGCAGTACATGGAACACATACCTTCCATATATTTACTGTCCAC  
2701 K S L I F V I I W C I V I F L A E V A A 876  
AAGAGCTTAATTTTGTGCTAATTTGGTCTTAGTAATTTTCTGGCAGAGGTCGCTCT  
2761 S I L V T L W L L G M T P L Q D K G N S T 896  
TCTTTGGTGTCTGCTGGCTCTTGGAACTCTCTCAAGCAAAAGGGAATAGTACT  
2821 H S R N N S Y A V I I T S T S S Y Y V F 916  
CATAGTAGAAAATACAGCTATGAGTATATCACAGCAGCAGTCTCGATATATGTGTTT  
2881 Y I Y V G V A D T L L A M G F F R G L P 936  
TACATTTACGTTGGAGTAGCCGACCTTGTCTGCTTGAAGGATCTCAGAGGCTTACCA  
2941 L V H T L I T V S K I L H H K M L H S V 956  
CTGGTACTACTTAATCAGTGTGAAAATTTTACACCACAAAATGTTACATCTCTGT  
3001 L Q A P M S T L N T L K A F G I L N R F 976  
CTTCAAGCACCATTCAACCTCAACCGTGAAGCAGTGGGCTTCTTAATAGTTC  
3061 S K D I A I L D D L L P L T F D F I O 996  
TCCAAGATATAGCAATTTGGATGACCTTCTGCTTCTACCATATTTGACTTCAATCCAG  
3121 L L L I F V I G A I A V V A V I O P Y I F 1016  
TTGTTATTAATGTTGAGTGGAGCTATAGCAGTTGCTGCGACTTCAACCCCTCATCTTT  
3181 V Z T V P V I V A F I M L R A Y F Q T 1036  
GTTCCACAGTCCAGTACTAGTGGCTTTTATTGTTGAGAGCAATTTCTCCCAAAC  
3241 S Q L K Q L E S E G R S P I F T H L V 1056  
TCACAGCACTCAAACAATGGAATCTGAAGCAGGAGTCCAATTTCACTACTTCTGTT  
3301 T S L K G L W T L R A F G R Q P Y F E T 1076  
ACAAGCTTAAAGACTATGGACCTTCTGCTCGGACGGCAGCCTTCAATTTGAAACT  
3361 L F H K A L N L H T A N W F L Y L S T L 1096  
CTGTTCCAAAAGCTCTGAATTTACATCTGCAACTGGTCTTGTACTGTCAACTG  
3421 R W F Q M R I E M I F V I F F I A V T F 1116  
CGTGTCCAAATGAGAATAGAAATGATTTTGTCTCTCTCTTCAATGATGCTTCT  
3481 I S I L T T G E G E G N V G I I L T L A 1136  
ATTTCATTTTAAACAAGAGAGAAAGGAGTGTGATATTCTGACTTTAGCC  
3541 M N I M S T L O W A V N S I D V D S L 1156  
ATGAATCATGAGTACATTCAGCTGGGCTGTAACCTCAGCATGATGTTGATGCTTCT  
3601 M R S V S R V F K F I D M P T E G K P T 1176  
ATGCGATCTGTGAGCCGAGTCTTAAAGTTCATGACATGCCACAGAAAGTAAACCTACC  
3661 K S T K P Y K N G Q L S K V M I I E N S 1196  
AAGTCAACCAACATACAGAAATGGCCAACTCTGAAAGTTATGATTTGAGAAATCA  
3721 H V K K D D I W P S G G Q M T V K D L T 1216  
CAGGTGAGAAAGATGACATCTGGCCCTCAGGGGCAAACTGACTGCAAGATCTCACA  
3781 A K Y T E G G N A I L E N I S F S I S P 1236  
GCAAAATACACAGAAAGTGAATGCCATATAGAGAACATTCTCTCAATAAGCTCT  
3841 G O R V G L L G R T G S G K S T L L S A 1256  
GGCCAGAGGTGGCCCTTGGGAGAACTGGATCAGGGAAGACTTGTATTCAGCT  
3901 F L R L N T E G E I O I D G V S W D S 1276  
TTTTGAGACTACTGAACCTGAAGGAGAAATCCAGATCGATGGTGTCTTGGGATCA  
3961 I T L O O W R K A F G V I P O K V F I F 1296  
ATAACTTTGCAACAGTGGAGAAAGCTTGGAGTATACACAGAAAGTATTTATTTT  
4021 S G T F R K N L D P Y E O F F G K I W 1316  
TCTGGAACATTTAGAAAACCTTGGATCCCTATGAACAGTGGAGTATCAAGAAATG  
4081 K V A D E V G C L R S V I E O F P G K L D 1336  
AAAGTGCAGATGAGTTGGCTCAGATCTGTGATAGACAGTTTCTGGGAAGDGTGAC  
4141 F V L V D G G C V L S H G E K Q L M C L 1356  
TTTTCTCTGTGGATGGGGCTGTGCTTAAAGCCTAGCCACAGGAGTGTGATGTGCTG  
4201 A R S V L S K A K I L L L D E P S A H L 1376  
GCTAGATCTGTTCTCAGTAAAGGCAAGTCTTCTGCTTGTATGAACCAAGTCTCATTTG  
4261 D P V T Y Q I I R R T L K Q A F A D C T 1396  
GATCCAGTACATACCAATTAATAGAAAGCTTCAAAAACAGCATTTGCTGATGACACA  
4321 V I L C E H R I E A M L E C C O Q F L V I 1416  
GTAATCTCTGTGAACACAGGATAGAACTGCTGGAATGCCAACAAATTTTGTGTCATA  
4381 E E N K V R Q Y D S I O K L L N E R S L 1436  
GAAGAGACAAAGTGGGAGTACGATCCATCCAGAACTGCTGAACGAGAGGAGCCCTC  
4441 F R Q A I S P S D R V K L F P H R N S 1456  
TTCCGGACAGCCATCAGCCCTCCGACAGGTTGAGCTCTTCCCAACCCGAACTCAAGC  
4501 K C K S K P Q I A A L K E E T E E E V Q 1476  
AAGTGAAGCTAAGCCCAAGATGCTGCTGTAAGAGGAGACAGAAAGAGGAGTCAAA

D T R L =

```

4561 GATACAAGGCTTGAAGAGCAGCATAAATGTTGACATGGGACATTTGCTCATGGAATGGG
4621 AGCTCGTGGGAGCAGTCACCTCATGGAATGGAGCTCGTGGAAACAGTTACCTCTGCCTCAG
4681 AAACAAGGATGAATTAAGTTTTTTTTTAAAGAAACATTGGTAAGGGGAATGAGG
4741 ACACATGATGGGCTTGATAAATGGCTTCCTGGCAATAGTCAAATGTGTGAAAGGATG
4801 TTCAAATCCCTGAAGATTACCACTTGTGTTTGAAGCCGAGATTTTCCTGAAAACCCCTT
4861 CCGATGCTAGTAAATGGAAGGCGAGCTCTAAAATGTCATCAGCCTAGTGTAGTACGCTT
4921 ATTGCTAGTGAATCGTTAATTTGATGTTGAGGAAAGAACTGAAATCATACTCTCTTA
4981 GGGTTATGATTAAGTAAATGATAACTGGAAATTCAGCGGTTTATATAAGCTTGTATTCCT
5041 TTTTCTCTCTCCCATGATGTTTGAAGAACACAACATATATTGTTGCTAAGCATTCCA
5101 ACTATCTCATTCCCAAGCAAGTATAGAATACACAGGAACCAACAGACTGCACATCAAA
5161 ATATGCCCAATCAACATCTAGTGAGCAGTCAGGAAAGAGAACTCCAGATCCTGGAAAT
5221 CAGGGTATGATTTGCCAGGCTACCAAAAACTCAATATTCAGATAATCACAAATACAT
5281 CCCTTACCTGGGAAAGGCTGTTAATCTTTCACAGGGGACAGGATGGTCCCTTGATG
5341 AAGAAGTTGATATGCTTTTCCCAACTCCAGAAAGTGACAGCTCACAGACCTTGAAC
5401 AGAGTTTAGCTGGAAAGATGTTAAGTGCAAAATGTCACAGGACAGCCCTTCTTCCACA
5461 GAAGTCCAGGTAGAGGCTGTGTAAGTAGATAGGCCATGGGACCTGTGGGTAGACACACA
5521 TGAAGTCCCAAGCATTAGATGTATAGGTTGATGSGTGTATGTTTTCAGGCTAGATGTATG
5581 TACTTCATGCTGTCTACACTAAGAGAGAATGAGAGACACTGAAAGACCAACATCATG
5641 AATTAGTTTATATGCTTCTGTTTATAAATTTGTAAGCAAAATTTTTCTTAGGAAA
5701 TATTTATTTAATAATCTTCAACAGTATATACAACTGCTATTTAAAGAAATGATTA
5761 TGAATACAGCTGATAAATAATTTTATATTTGAATATGACTTTTATGGCACTAG
5821 TATTTATGAATATTATGTTAATACTGGGACGGGAGAACCTAGGCTGATTTAAAC
5881 AGGGCCATGAATACCTTTGGTCTGGAGGAAAGCCTGGGGCTGATCGAGTGTGTGCC
5941 CACAGCTATGATTTCCAGCCAGACAGCCCTTTAGATGCACTTGAAGAAGATGGT
6001 ACCACAGTCTGACTGTTTCCATCAAGGCTACACTGCCTTCTCAACTCCAAACTGACTCT
6061 TAAGAAGACTGCATATATTTATTACTGTAAAGAAATATCACTTGTCAATAAAATCCATA
6121 CATTGTGT (A)n


```

**Fig. 2.** Nucleotide sequence of cDNA encoding the CF transmembrane conductance regulator together with the deduced amino acid sequence. DNA sequencing was performed by the dideoxy chain termination method (34) with <sup>32</sup>S-labeled nucleotides or by the Dupont Genesis2000 automatic DNA sequencer. Numbers on the left of columns indicate base positions and numbers on the right amino acid residue positions. The first base position corresponds to the first nucleotide in the 5' extension clone PA3-5, which is one nucleotide longer than TB2-7 (12). The 3' end and the noncoding sequence are shown above [nucleotides 4561 to 6129 plus the poly(A)<sup>+</sup> tail]. Arrows indicate position of transcription initiation site by primer extension analysis (11). Nucleotide 6129 is followed by a poly(A) tract. Positions of exon junctions are indicated by vertical lines. Potential membrane-spanning segments ascertained with the use of the algorithm of Eisenberg *et al.* (35) are enclosed in boxes. Amino acids comprising putative ATP-binding folds are underlined. Possible sites of phosphorylation (21) by protein kinases A or C are indicated by open and closed circles, respectively. The open triangle indicates the position at which 3 bp are deleted in CF. Abbreviations for the amino acid residues are: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr.


uous coding region of the CF locus could be deduced from overlapping cDNA clones. Since most of the cDNA clones were apparently derived from unprocessed transcripts, further studies were performed to ensure the authenticity of the consensus sequence. Each cDNA clone was first tested for chromosome localization by hybridization analysis with a human-hamster somatic cell hybrid containing a single human chromosome 7 and by pulsed field gel electrophoresis (7). The ones that did not map to the correct region on chromosome 7 were not pursued. Fine restriction enzyme mapping was then performed for each clone. While overlapping regions were clearly identified for most of the clones, many contained single copy, additional regions not readily recognizable by restriction enzyme analysis.

The cDNA was further characterized in gel hybridization experiments with genomic DNA. Five to six different restriction fragments could be detected with the 10-1 cDNA in Eco RI- or Hind III-digested total human DNA and a similar number of fragments with several other cDNA clones, suggesting the presence of multiple exons for the putative CF gene. The hybridization studies also identified the cDNA clones with unprocessed intron sequences when they showed preferential hybridization to a smaller subset of genomic DNA fragments with relatively greater intensities. For the confirmed cDNA clones, their corresponding genomic DNA segments were isolated (7) and the exons and exon-intron boundaries were sequenced. In all, 24 exons were identified (Fig. 2). Physical mapping experiments (7) showed that the gene locus spanned about 250 kb.

The 5' terminus of the transcript was determined by primer extension (11). A modified polymerase chain reaction, anchored PCR (12), was also used to facilitate cloning of the 5' end sequences.



**Fig. 3.** RNA gel-blot analysis. Hybridization by the cDNA clone 10-1 to a 6.5-kb transcript is shown in the tissues indicated. RNA samples were prepared from cells and tissue samples obtained from surgical pathology or at autopsy according to the methods described in (10). Total RNA (10 µg) from each tissue and 1 µg of poly(A)<sup>+</sup> RNA from T84 cells were separated on formaldehyde gels and transferred onto nylon membranes (Zetaprobe, Bio-Rad), which were hybridized with DNA probes labeled to high specific activity by the random priming method (36, 37). The positions of the 28S and 18S rRNA bands are indicated.



**Fig. 4.** DNA sequence around the  $\Delta F_{508}$  deletion. The normal sequence from base position 1627 to 1651 (from cDNA T16-1) is shown beside the CF sequence (from cDNA C16-1). The left panel shows the sequences from the coding strands obtained with the B primer (5'-GTTTTCTGGAT-TATGCCTGGGAC-3') and the right panel those from the opposite strand with the D primer (5'-GTTGGCATGCTTTGATGACGCTC-3'). The brackets indicate the three nucleotides in the normal that are absent in CF (arrowheads). Sequencing was performed as described in (34).

Two independent 5' extension clones, one from pancreas and the other from T84 RNA, were characterized by DNA sequencing and differed by only 1 base in length, thus establishing the most probable initiation site for the transcript (Fig. 2). Since the initial cDNA clones did not contain a poly(A)<sup>+</sup> tail indicative of the end of a mRNA, anchored PCR was also applied to the 3' end of the transcript (12). The results derived from the use of several different 3'-extending oligonucleotides were consistent with the interpreta-





not believed to conduct ions, has only two charged residues in all 12 transmembrane domains. Alternatively, CFTR may not be an ion channel but instead it may serve to regulate ion channel activities. In support of the latter possibility, none of the recently purified polypeptides (from trachea and kidney) that are capable of reconstituting chloride channels in lipid membranes (6) appear to be CFTR, judged on the basis of molecular mass.

In any case, the presence of ATP-binding domains in CFTR suggests that ATP hydrolysis is directly involved and required for the transport function. The high density of phosphorylation sites for protein kinases A and C and the clusters of charged residues in the R domain may both serve to regulate this activity. The deletion of Phe<sup>508</sup> in the NBF may prevent proper binding of ATP or the conformational change required for normal CFTR activity, consequently resulting in the observed insensitivity to activation by protein kinase A- or protein kinase C-mediated phosphorylation of the CF apical chloride conductance pathway (5). Since the predicted structure of CFTR contains several conserved domains and belongs to a family of proteins, most of which function as parts of multicomponent molecular systems (15), the CFTR protein may also participate in epithelial cell functions not related to ion transport.

To understand the basic defect in CF, it is necessary to determine the precise role of Phe<sup>508</sup> in the regulation of ion transport and to understand the mechanism that leads to the pathophysiology of the disease. With the CF gene (that is, the cDNA) now isolated, it should be possible to elucidate the control of ion transport pathways in epithelial cells in general. Knowledge gained from study of the CF gene product (CFTR), both the normal and mutant forms, will provide a molecular basis for the development of improved means of treatment of the disease.

#### REFERENCES AND NOTES

1. T. F. Boat, M. J. Welsh, A. L. Beaudet, in *The Metabolic Basis of Inherited Disease*, C. L. Scriver, A. L. Beaudet, W. S. Sly, D. Valle, Eds. (McGraw-Hill, New York, ed. 6, 1989), pp. 2649-2680.
2. M. R. Knowles *et al.*, *Science* **221**, 1067 (1983); M. R. Knowles, J. Gatzky, R. Boucher, *J. Clin. Invest.* **71**, 1410 (1983); P. M. Quinton, *Nature* **301**, 421 (1983); K. Sato, *Am. J. Physiol.* **247**, R646 (1984).
3. J. H. Widdicombe, M. J. Welsh, W. E. Finkbeiner, *Proc. Natl. Acad. Sci. U.S.A.* **82**, 6167 (1985); P. M. Quinton and J. Bijman, *N. Engl. J. Med.* **308**, 1185 (1983).
4. J. R. Yankaskas, C. U. Cotton, M. R. Knowles, J. T. Gatzky, R. C. Boucher, *Am. Rev. Resp. Dis.* **132**, 1281 (1985); P. S. Pedersen, E. H. Larsen, B. Hainau, N. J. Brandt, *Med. Sci. Res.* **15**, 1009 (1987); T. J. Jensen and J. R. Riordan, unpublished results.
5. M. J. Welsh, *Science* **232**, 1648 (1986); R. A. Frizzell, G. Rechkemmer, R. L. Shoemaker, *ibid.* **233**, 558 (1986); M. J. Welsh and C. M. Liedtke, *Nature* **322**, 467 (1986); R. A. Schoumacher *et al.*, *ibid.* **330**, 752 (1987); M. Li *et al.*, *ibid.* **331**, 358 (1988); T.-C. Hwang *et al.*, *Science* **244**, 1351 (1989); M. Li *et al.*, *ibid.*, p. 1353.
6. D. W. Landry *et al.*, *Science* **244**, 1469 (1989).
7. J. M. Rommens *et al.*, *ibid.* **245**, 1059 (1989).
8. M. Farrall *et al.*, *Am. J. Hum. Genet.* **43**, 471 (1988).
9. B. Kerem *et al.*, *Science* **245**, 1073 (1989).
10. The cDNA libraries from cultured epithelial cells were prepared as follows: sweat gland cells derived from a non-CF individual and from a CF patient were grown to first passage as described [G. Collic, M. Buchwald, P. Harper, J. R. Riordan, *In Vitro Cell. Dev. Biol.* **21**, 592 (1985)]. The presence in these cells of an outwardly rectifying Cl<sup>-</sup> channel was confirmed (J. A. Tabcharani, T. J. Jensen, J. R. Riordan, J. W. Hanrahan, *J. Membrane Biol.*, in press), but the CF cells were insensitive to activation by cyclic AMP [T. J. Jensen, J. W. Hanrahan, J. A. Tabcharani, M. Buchwald, J. R. Riordan, *Pediatric Pulmonol. Suppl.* **2**, 100 (1988)]. Polyadenylated RNA was isolated [J. M. Chirgwin, A. E. Przybyla, R. J. Macdonald, W. J. Rutter, *Biochemistry* **18**, 5294 (1979); H. Aviv and P. Leder, *Proc. Natl. Acad. Sci. U.S.A.* **69**, 1408 (1972)] and used as template for the synthesis of cDNA according to U. Gubler and B. Hoffman [*Gene* **25**, 263 (1983)]. After methylation of internal Eco RI sites, ends were made flush with T4 DNA polymerase, and phosphorylated Eco RI linkers were added to the cDNA. After digestion with Eco RI and removal of excess linkers, the cDNA products were ligated into the Eco RI site of  $\lambda$  ZAP (Stratagene, San Diego, CA). The same procedures were used to construct a library from RNA isolated from preconfluent cultures of the T84 colon carcinoma cell line [K. Dharmathaphorn, J. A. McRoberts, K. G. Mandel, L. D. Tisdale, H. Masui, *Am. J. Physiol.* **246**, G204 (1984)]. The numbers of independent recombinants in the three libraries were: 2.0  $\times 10^6$  for the non-CF sweat gland cells, 4.5  $\times 10^6$  for the CF sweat gland cells, and 3.2  $\times 10^6$  from T84 cells. Standard procedures were used for screening [T. Maniatis, E. F. Fritsch, J. Sambrook, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1982)]. Bluescript plasmids were rescued from plaque-purified clones by excision with M13 helper phage (Stratagene). The lung and pancreas libraries were purchased from Clontech Lab Inc. (Catalog Nos. HL1066b and HL1069h, respectively).
11. The start point of the CF gene transcript was derived by primer extension procedures [F. J. Calzone, R. J. Britten, E. H. Davidson, *Methods Enzymol.* **152**, 611 (1987)]. The oligonucleotide primer [positioned 157 nucleotides (nt) from the 5' end of the 10-1 clone] was end-labeled with [ $\gamma$ -<sup>32</sup>P]ATP (Amersham, 5000 Ci/mole) and T4 polynucleotide kinase, purified by gel filtration, and annealed with ~5  $\mu$ g of T84 poly(A)<sup>+</sup> RNA for 2 hours at 60°C. The extension reaction was performed at 41°C for 1 hour with avian myeloblastosis virus (AMV) reverse transcriptase (Life Sciences, Inc.) and terminated by addition of NaOH to 0.4M and EDTA to 20 mM, with subsequent neutralization with ammonium acetate (pH 4.6). The products were treated with phenol, precipitated with ethanol, redissolved in buffer with formamide, and analyzed on a polyacrylamide sequencing gel.
12. The anchored PCR procedure [M. A. Frohman, M. K. Dush, G. R. Martin, *Proc. Natl. Acad. Sci. U.S.A.* **85**, 8998 (1988)] was used to synthesize cDNA corresponding to the 5' and 3' ends of the transcript. For the 5' end clones, poly(A)<sup>+</sup> RNA from pancreas and T84 cells were subjected to reverse-transcription with the use of an exon 2-specific primer (11). The first strand cDNA products were fractionated on an agarose column and the fractions containing large species were identified by gel electrophoresis after the polymerase chain reaction [R. K. Saiki *et al.*, *Science* **230**, 1350 (1985)] with a pair of oligonucleotide primers (145 nt apart within the 10-1 sequence) just 5' of the extension primer. These products were pooled, concentrated, and treated with terminal deoxynucleotidyl transferase (BRL) and dATP, as recommended by the supplier. Second strand synthesis was performed with Taq Polymerase (Cetus, AmpliTaq) and an oligonucleotide containing a linker sequence, 5'-CGAATTCCTCGAGATC(T)<sub>12</sub>-3'. This linker, together with another primer (internal to the extension primer) with an Eco RI restriction site at its 5' end, was then used for PCR. After digestion with Eco RI and Bgl II, products were purified and cloned in Bluescript KS (Stratagene) by standard procedures. All the recovered clones contained inserts of more than 350 nt. The 3' end clones were generated with the use of similar procedures. PCR amplification was carried out with the linker described above and an oligonucleotide with the sequence 5'-ATGAAGTCCAAGGATTTAG-3', which is ~70 nt upstream of the Hind III site at position 5027 (Fig. 2). The products were digested with Hind III and Xho I and cloned in the Bluescript vector. Candidate clones were identified by hybridization with the 3' end of cDNA T16-4.5. All PCR's were performed for 30 cycles as described by the enzyme supplier.
13. M. Kozak, *Nucleic Acids Res.* **12**, 857 (1984); *Cell* **44**, 283 (1986).
14. Other sequence differences were noted between the normal (T16-4.5) and CF (C1-1/5) cDNA clones. At position 2629 (Fig. 2), T16-4.5 contained a C and C1-1/5 a T, resulting in a change of Leu to Phe. At position 4555, the base was G in T16-4.5 but A in C1-1/5 (Val to Met). The differences may be results of cDNA cloning artifacts or may represent sequence polymorphisms. Specific oligonucleotide hybridization analysis of patient or family DNA should distinguish these possibilities. Since these changes are conserved amino acid substitutions, they are unlikely to be causative mutations. Additional nucleotide differences were observed in the 3' untranslated region between different cDNA clones and the corresponding genomic DNA sequence.
15. C. F. Higgins, M. P. Gallagher, M. L. Mimmack, S. R. Pearce, *BioEssays* **8**, 111 (1988); C. F. Higgins *et al.*, *Nature* **323**, 448 (1986).
16. P. Gros, J. Croop, D. E. Housman, *Cell* **47**, 371 (1986); C. Chen *et al.*, *ibid.*, p. 381; J. H. Gerlach *et al.*, *Nature* **324**, 485 (1986); P. Gros, M. Raymond, J. Bell, D. Housman, *Mol. Cell. Biol.* **8**, 2770 (1988).
17. Several large families of integral membrane proteins are known, including: (i) A number of ligand-gated ion channels of which the nicotinic acetylcholine receptor [R. M. Stroud and J. Finer-Moore, *Annu. Rev. Cell Biol.* **1**, 317 (1985)] is the prototype. Receptors for the inhibitory neurotransmitters GABA (33) and glycine are included in this family. (ii) A family of ion channels with a totally different structural motif are the voltage-gated, sodium, calcium, and potassium channels (27). (iii) Involved in the translocation of ions are the structurally related cation pumps such as the Ca<sup>2+</sup>-ATPase [C. J. Brandt, N. M. Green, B. Korczak, D. H. MacLennan, *Cell* **44**, 597 (1986)], the Na<sup>+</sup>,K<sup>+</sup>-ATPase [G. E. Shull and J. B. Lingrel, *Proc. Natl. Acad. Sci. U.S.A.* **84**, 4039 (1987)], and the H<sup>+</sup>,K<sup>+</sup>-ATPase [G. E. Shull and J. B. Lingrel, *J. Biol. Chem.* **261**, 16788 (1986)]. These are but examples.
18. G. B. Wilson, T. L. Jahn, J. R. Fonseca, *Clin. Chim. Acta* **49**, 79 (1973); V. van Heyningen, C. Hayward, J. Fletcher, C. McAuley, *Nature* **315**, 513 (1985).
19. J. Garnier, D. J. Osguthorpe, B. Robson, *J. Mol. Biol.* **120**, 97 (1978); J. Kyte and R. F. Doolittle, *ibid.* **157**, 105 (1982).
20. D. D. Pless and W. J. Lennarz, *Proc. Natl. Acad. Sci. U.S.A.* **74**, 134 (1977).
21. P. J. Blackshear, A. C. Nairn, J. F. Kuo, *FASEB J.* **2**, 2957 (1989).
22. J. Hess, W. Wels, M. Vogel, W. Goebel, *FEMS Microbiol. Lett.* **34**, 1 (1986).
23. K. O'Hare, C. Murphy, R. Levis, G. M. Rubin, *J. Mol. Biol.* **180**, 437 (1984).
24. M. J. Friedrich, L. C. DeVeaux, R. J. Kadner, *J. Bacteriol.* **167**, 928 (1986).
25. K. Ohyama *et al.*, *Nature* **322**, 572 (1986).
26. G. F.-L. Ames, *Annu. Rev. Biochem.* **55**, 397 (1986).
27. M. Noda *et al.*, *Nature* **320**, 188 (1986); T. Tanabe *et al.*, *ibid.* **328**, 313 (1987); A. Baumann *et al.*, *EMBO J.* **6**, 3419 (1987).
28. C.-M. Chen, T. K. Misra, S. Silver, B. P. Rosen, *J. Biol. Chem.* **261**, 15030 (1986).
29. J. Krupinski *et al.*, *Science* **244**, 1558 (1989).
30. In addition to the major NBF homologies, searches of the PIR and SWISSPROT data bases detected shorter stretches of sequence homology with other proteins including the following:

CFTR 122-136	Y L G I G L C L L F I V R T L
	: : : : : : : :
GLNP 198-212	Y L I I T L V L S F I L R R L
CFTR 307-319	S S A F F F S G F F V V F
	: : : : : : : :
COX 89-101	S E V F F F A G F F W A F
CFTR 701-713	I L N P I N S I R K F S I
	: : : : : : : :
NaCh 111-123	I L T P F N P I R K L A I
CFTR 1425-1442	D S I Q K L L N E R S L F R Q A I S
	: : : : : : : :
raf 578-595	D S I K K L R D E R P L F P Q I L S

GLNP, glutamine permease of *E. coli* [T. Nohno, T. Saito, J. Hōng, *Mol. Gen. Genet.* **205**, 260 (1986)]; COX, human cytochrome c oxidase polypeptide III [S. Anderson *et al.*, *Nature* **290**, 457 (1981)]; NaCh, rat brain sodium channel III (32); raf, the serine-threonine kinase proto-oncogene of *Xenopus laevis* (31). The first two sequences are within membrane spanning segments and probably reflect only coincidental arrangements of the hydrophobic residues suited to this function. In contrast, the latter two sequences are both in polar hydrophilic regions of the proteins. The large extent of amino acid conservation (11 of 13 residues) implies some functional relation between these short segments of the primary structure of the Na<sup>+</sup> channel and CFTR. Similarities between sequences at the same relative locations with respect to the COOH-termini of the raf kinase and CFTR suggest that they may also share at least a small facet of their structures and functions.

31. R. LeGuellec, K. LeGuellec, J. Paris, M. Philippe, *Nucleic Acids Res.* **16**, 10357 (1988).

32. M. Noda *et al.*, *Nature* **312**, 121 (1984); L. Salkoff *et al.*, *Science* **237**, 744 (1987).  
 33. P. R. Schofield *et al.*, *Nature* **328**, 221 (1987).  
 34. F. Sanger, S. Nicklen, A. R. Coulson, *Proc. Natl. Acad. Sci. U.S.A.* **74**, 5463 (1977).  
 35. D. Eisenberg, E. Schwarz, M. Komaromy, R. Wall, *J. Mol. Biol.* **179**, 125 (1984).  
 36. A. P. Feinberg and B. Vogelstein, *Anal. Biochem.* **132**, 6 (1983).  
 37. J. Rommens *et al.*, *Am. J. Hum. Genet.* **43**, 645 (1988).  
 38. S. J. Foote *et al.*, *Cell* **57**, 921 (1989).  
 39. J. P. McGrath and A. Varshavsky, *Nature* **340**, 400 (1989).  
 40. B. P. Surin, H. Rosenberg, G. B. Cox, *J. Bacteriol.* **161**, 189 (1985).  
 41. C. F. Higgins *et al.*, *Nature* **298**, 723 (1982).  
 42. E. Gilson, H. Nikaido, M. Hofnung, *Nucleic Acids Res.* **10**, 7449 (1982).  
 43. I. D. Hiles, M. P. Gallagher, D. H. Jamieson, C. F. Higgins, *J. Mol. Biol.* **195**, 125 (1987).  
 44. A. W. Bell *et al.*, *J. Biol. Chem.* **261**, 7652 (1986).  
 45. R. F. Doolittle *et al.*, *Nature* **323**, 451 (1986).  
 46. I. J. Evans and J. A. Downie, *Gene* **43**, 95 (1986).  
 47. D. R. Gill, G. H. Hatfull, G. P. C. Salmond, *Mol. Gen. Genet.* **205**, 134 (1986).  
 48. J. E. Walker, M. Saraste, M. J. Runswick, N. J. Gay, *EMBO J.* **1**, 945 (1982).  
 49. D. C. Fry, S. A. Kuby, A. S. Mildvan, *Proc. Natl. Acad. Sci. U.S.A.* **83**, 907 (1986).  
 50. We thank O. Augustinas for the collection of tissues; T. Jensen and R. Baird for the culturing of epithelial cells; L. Naismith for the isolation of RNA; D. Kennedy and D. Markiewicz for technical assistance and M. Buchwald and M. Dean for discussions. Supported by NIH grants DK39690 (F.S.C.) and DK34944 (L.C.T.), the Cystic Fibrosis Foundation (U.S.A.), the Canadian Cystic Fibrosis Foundation, and the Sellers Fund. J.M.R. holds a postdoctoral fellowship from the Medical Research Council (MRC) of Canada and F.S.C. is an Associate Investigator of the Howard Hughes Medical Institute.

7 August 1989; accepted 18 August 1989

# Identification of the Cystic Fibrosis Gene: Genetic Analysis

BAT-SHEVA KEREM, JOHANNA M. ROMMENS, JANET A. BUCHANAN,  
DANUTA MARKIEWICZ, TARA K. COX, ARAVINDA CHAKRAVARTI,  
MANUEL BUCHWALD, LAP-CHEE TSUI

Approximately 70 percent of the mutations in cystic fibrosis patients correspond to a specific deletion of three base pairs, which results in the loss of a phenylalanine residue at amino acid position 508 of the putative product of the cystic fibrosis gene. Extended haplotype data based on DNA markers closely linked to the putative disease gene locus suggest that the remainder of the cystic fibrosis

mutant gene pool consists of multiple, different mutations. A small set of these latter mutant alleles (about 8 percent) may confer residual pancreatic exocrine function in a subgroup of patients who are pancreatic sufficient. The ability to detect mutations in the cystic fibrosis gene at the DNA level has important implications for genetic diagnosis.

**A**LTHOUGH THE FREQUENCY OF CYSTIC FIBROSIS (CF) IS not uniformly high among all Caucasian populations, a consensus estimate is that it occurs once in 2000 live births (1). On the basis of the autosomal recessive mode of inheritance for this disease, a mutant allele frequency of 0.022 may be derived. Several different mechanisms, including high mutation rate (2),

heterozygote advantage (3), genetic drift (4), multiple loci (5), and reproductive compensation (6), have been proposed in attempts to explain the high incidence and, indirectly, the nature of the CF mutations. Although some of these hypotheses could not be further addressed because of the lack of knowledge about the basic defect in CF, several important observations have been made during the past few years through genetic analysis of the families of affected individuals (7-20).

B. Kerem, J. M. Rommens, J. A. Buchanan, D. Markiewicz, M. Buchwald and L.-C. Tsui are in the Department of Genetics, Research Institute, The Hospital for Sick Children, Toronto, Ontario M5G 1X8, Canada. T. K. Cox and A. Chakravarti are in the Department of Human Genetics, University of Pittsburgh, Pittsburgh, PA 15261. M. Buchwald and L.-C. Tsui are also members of the Departments of Medical Genetics and Medical Biophysics, University of Toronto, Toronto, Ontario M5S 1A8, Canada.

Extensive linkage analysis provides evidence for the existence of a single CF locus on human chromosome 7 (region q31) (7-10, 21). The detection of allelic and haplotype association between the CF