the twig morph directly, or as Faeth and Hammon suggest, through their interaction and binding with proteins, thus lowering the amount of digestible protein. The feed-ing experiments (1, table 2) do not distinguish between these two alternatives.

The main purpose of the paper (1) was to report the existence of this diet-induced developmental polymorphism. Although I suggested that polyphenolic compounds may be important, I had hoped to make it clear that we still do not know what it is that induces the development of the different morphs by stating, "This developmental polymorphism *may* be triggered by the concentration of defensive secondary compounds in the larval diet" (emphasis added) and "a possible mechanism is that receptors respond to tannin levels...." I thank Faeth and Hammon for clarifying these points and for stating more explicitly another hypothesis concerning the mechanism involved.

> ERICK GREENE Department of Avian Sciences, University of California, Davis, CA 95616

REFERENCES

1. E. Greene, Science 243, 643 (1989).

17 May 1989; accepted 19 September 1989

## Does Voltage Affect Excitation-Contraction Coupling in the Heart?

In their report of experiments investigating excitation-contraction coupling in cardiac muscle, Näbauer et al. (1) say they have demonstrated that "it is exclusively Ca<sup>2-</sup> influx through calcium channels ... that regulates Ca<sup>2+</sup> release in cardiac myocytes (emphasis added)." The experiments they present demonstrate that internal Ca<sup>2+</sup> release can occur when the membrane is depolarized under conditions that permit calcium influx and that if Ca2+ influx is prevented, Ca2+ release does not occur. Their experiments therefore confirm previous work demonstrating that membrane depolarization alone cannot trigger release of Ca<sup>2+</sup> from internal stores (2-4). However, they did not examine the effects of varying membrane potential under conditions that allow Ca<sup>2+</sup> influx. Thus their data do not contradict the records of Cannell et al. (3), which show that repolarization can switch off calcium release from the sarcoplasmic reticulum (3, 4), nor do they refute our suggestion (3)that "Ca<sup>2+</sup>release may be mediated by a mechanism that requires a  $Ca^{2+}$  influx but which may be modulated by changes in voltage directly." (We did not suggest that voltage alone can trigger Ca2+ release, as might be inferred from the context of the citation to our work.) Since modulation of calcium release by voltage can explain all experimental data to date, more critical tests of this hypothesis are needed.

M. B. CANNELL

Department of Pharmacology, University of Miami School of Medicine, Miami, FL 33143 J. R. BERLIN Graduate Hospital, Bockus Research Institute, Philadelphia, PA 19146 W. J. LEDERER Department of Physiology, University of Maryland School of Medicine, Baltimore, MD 21201

## REFERENCES

- 1. M. Näbauer, G. Callewaert, L. Cleemann, M. Morad, Science 244, 800 (1989).
- H. A. Fozzard and D. C. Hellam, Nature 218, 688 (1968); W. R. Gibbons and H. A. Fozzard, J. Gen. Physiol. 65, 345 (1975); R. Ochi and W. Trautwein, Pflügers Arch. Eur. J. Physiol. 323, 187 (1971); W. Trautwein, T. F. McDonald, O. Tripathi, ibid. 354, 55 (1975); G. W. Beeler and H. Reuter, J. Physiol. (London) 207, 211 (1970); C. Leoty, ibid. 239, 237 (1974; B. London and J. W. Krueger, J. Gen. Physiol. 88, 475 (1986).
  M. B. Cannell, J. R. Berlin, W. J. Lederer, Science
- M. B. Cannell, J. R. Berlin, W. J. Lederer, Science 238, 1419 (1987)
- 4. D. J. Beukelmann and W. G. Wier, J. Physiol. (London) 405, 233 (1988).

15 June 1989; acccepted 15 September 1989

Response: The main issue addressed in our report (1) is the regulation of  $Ca^{2+}$  release from the sarcoplasmic reticulum in cardiac myocytes. Our experiments showed that there is no  $Ca^{2+}$  release when there is no  $Ca^{2+}$  current through the  $Ca^{2+}$  channel. Nor was there any release if  $Na^+$  or  $Ba^{2+}$  carried the charge through the  $Ca^{2+}$  channel [this finding was independent of holding potentials in the range from -60 mV to -90 mV, which rules out possible inactivation of gating charge (2)]. We feel justified, therefore, in maintaining that it is exclusively the influx of  $Ca^{2+}$  through the  $Ca^{2+}$  channel which regulates  $Ca^{2+}$  release in cardiac cells.

Our findings also suggest that membrane potential plays no direct role in the release process. Previously we showed that the voltage dependence of the intracellular  $Ca^{2+}$ 

transients closely follows that of the inward  $Ca^{2+}$  current (3). It is not correct, therefore, to say that we "did not examine the effects of varying membrane potentials." In addition, others have also found that early estimates of the Ca<sup>2+</sup> transients are well correlated with the inward Ca<sup>2+</sup> current, even when a more slowly developing  $Ca^{2+}$  transient is present (as it may be in  $Na^+$ -loaded cardiac cells) (4). The confusion surrounding the voltage dependence of the Ca<sup>2+</sup> transient may result from comparison of the Ca<sup>2+</sup> current with the much later occurring maximum value of the  $Ca^{2+}$  transient (5). Such a comparison may be flawed, not only because of the presence of other Ca<sup>2+</sup> transport mechanisms, but also because of saturation of the Ca<sup>2+</sup> indicator dye or exhaustion of the releasable Ca<sup>2+</sup> pools.

As for the question of whether membrane potential plays a limited modulatory role, Cannell et al. point to their observation that the rising phase of the  $Ca^{2+}$  transient at 0 mV is interrupted by early repolarization to the holding potential (-40 to -80 mV). We repeated these experiments and found in addition that the rising phase of the Ca<sup>2+</sup> transient could also be interrupted by further depolarization to +80 mV (ECa), where Ca<sup>2+</sup> transients are generally completely suppressed. We interpret this as indicating that the  $Ca^{2+}$  release is stopped when the  $Ca^{2+}$  current is interrupted either by repolarization-induced deactivation or by depolarization to the reversal potential. Repolarizing pulses to -80 mV, in addition, would activate the Na<sup>+</sup>-Ca<sup>+</sup> exchange to extrude Ca<sup>2+</sup> and help suppress the Ca<sup>2+</sup> transients. Therefore, even these experiments support our basic idea that it is not the membrane potential as such which is important for Ca<sup>2+</sup> release; rather it is the degree to which the membrane potential activates the  $Ca^{2+}$  current.

> M. MORAD L. CLEEMANN G. CALLAWAERT Department of Physiology, University of Pennsylvania, Philadelphia, PA 19104

## REFERENCES

- M. Näbauer, G. Callewaert, L. Cleemann, M. Morad, Science 244, 800 (1989).
- 2. G. Brum, E. Rios, E. Stefani, J. Physiol. 398, 441 (1988).
- 3. G. Callewaert, L. Cleemann, M. Morad, Proc. Natl. Acad. Sci. U.S.A. 85, 2009 (1988).
- G. Isenberg et al., in Biology of Isolated Adult Cardiac Myocytes, W. A. Clark, Ed. (Elsevier, New York, 1988), p. 354.
- 1988), p. 354.
  M. B. Cannell, J. R. Berlin, W. J. Lederer, *Science* 238, 1419 (1987).

<sup>3</sup> July 1989; accepted 18 September 1989