

ology can distort comparisons and produce spurious measures of biodiversity.

4. R. J. O'Hara, D. R. Maddison, and P. F. Stevens [*Science* **241**, 275 (1988)] concur.
5. K. S. Thomson [*Am. Scientist* **77**, 264 (1989)] notes the damage that data-centered terminology has inflicted on systematics.

EPA and Asbestos Removal

Environmental Protection Agency (EPA) Administrator William K. Reilly's letter disclaiming EPA responsibility for the removal of asbestos from buildings (1 June, p. 1064) is self-indicting. He clearly states that EPA only requires asbestos removal when building demolition or renovation activities threaten to release significant amounts of asbestos fibers into the air; he also seems to say that it is not the fault of EPA if, in his words, "a number of building owners are removing asbestos from their buildings . . . due to forces (for example, concerns about property devaluation, insurance, and liability) that may be unrelated to health risks."

Over the years there have been numerous EPA press releases pointing to the danger of asbestos and the need to protect public health. Reilly should reread those releases and then reaffirm the extent of EPA's involvement in creating the "forces" that have incited the need for asbestos removal.

EPA alone bears the responsibility for the "killer" image of asbestos. The American public, after spending billions of dollars removing it, now deserves to hear the facts with which the EPA can prove its claims about the danger of asbestos. The disclosure of that proof is long overdue.

JIM BERRY

Post Office Box 1811,
Rancho Santa Fe, CA 92067

British Radiation Study

The conundrum resulting from Martin J. Gardner's recent report (1) of paternal irradiation and childhood leukemia (*News & Comment*, 6 Apr., p. 24) begs for resolution by synthesis rather than refutation of either side. The missing factor may be the dietary habits (particularly the dietary fat intake) of the British and Japanese populations studied.

There is a dramatic difference in the incidence of certain cancers in the Japanese compared with that in Western populations, and a possible explanation involves both the lower intake of total fat and the higher percentage of ω -3 fatty acids in the Japanese diet. Animal studies have demonstrated a marked effect of oil seed ω -6 fatty acids as

tumor promoters when provided in the range similar to that in the current Western diet (2) and a countervailing effect of the long-chain ω -3 fatty acids obtained from cold water fish (3). Differences in dietary fatty acid composition can affect membrane content of highly unsaturated essential fatty acids, their eicosanoid products, and the expression of the *ras* oncogene (3).

At the time of exposure to the radiation from nuclear weapons used at Hiroshima and Nagasaki, the population of those cities were consuming only 10 to 15% of calories as fat, with ratios of ω -6 to ω -3 of about 1 to 1. In recent decades the Western diet has consisted of 30 to 40% fat calories with an ω -6 to ω -3 ratio of more than 10 to 1 (4). These dramatic differences in total fat and the ratios of metabolically distinct fatty acid families could be a factor in the differences in the post-exposure incidence of leukemia in the two populations. The diet of the father or the child may thus amplify or suppress the oncogenic initiating effect of the ionizing radiation.

STEPHEN D. PHINNEY

Division of Clinical Nutrition,
Department of Internal Medicine,
University of California,
Davis, CA 95616

REFERENCES

1. M. J. Gardner *et al.*, *Br. Med. J.* **300**, 423 (1990).
2. C. Ip, C. A. Carter, M. M. Ip, *Cancer Res.* **45**, 1997 (1985).
3. R. A. Karmali, in *Dietary Omega 3 and Omega 6 Fatty Acids; Biological Effects and Nutritional Essentiality*, C. Galli and A. P. Simopoulos, Eds. (Plenum, New York, NY, 1989), pp. 351-359.
4. O. Adams, in *ibid.*, pp. 33-41.

Usefulness of the Human Genome Project

The crisis in funding by the National Institutes of Health (NIH) of research R01 grants has renewed the dialog about the usefulness of the Human Genome Project. As long as there was adequate funding for basic research, this discussion was based on more theoretical grounds (for example, whether the human, mouse, yeast, and so forth, genome should be sequenced and how). However, with the drastic cutbacks in NIH funding, the discussion has become more personal as many investigators are questioning their survival in academic science without NIH grants.

Both the Reagan and Bush administrations have been committed to providing a certain amount of money for biological research. What has become apparent is that basic research funds are being removed from the general allotment of federal funds for

biological research and are being funneled into the Human Genome Project, which results in a contraction of basic science research funding.

With this point in mind I recently reviewed the research in my own lab in the context of what we would have done differently if the mouse genome had been sequenced before the start of our project. Specifically, we have recently completed the cloning and sequencing of the coding sequences for the murine complement receptor *Cr1* and *Cr2* genes. If we had had the 60 to 80 kilobases of mouse genomic sequence that contains these genes before we had started our work, what would we have done differently? Because one cannot look at a piece of genomic DNA and determine coding sequences, cDNAs have to be isolated and sequenced for the analysis of any gene. The most time-consuming and laborious steps in this project were the production and screening of the spleen and liver cDNA libraries and the subsequent sequencing of the cDNA clones. The sequence of the mouse genome would not have aided in this step of the project other than to provide confirmatory sequence information.

The only information that the full genomic sequence would have provided, in the context of our study, would have been the intron-exon border junctions. The *Cr2* gene covers about 50,000 base pairs in the genome and probably contains between 15 to 20 exons. We could easily determine the intron-exon organization of this gene on our own for the price of the oligonucleotides needed to sequence across the junctions. These oligonucleotides would cost about \$500. The mammalian genome contains about 5×10^9 base pairs of DNA, which, on the basis of our \$500-for-50,000-base-pairs estimate, means that we as a scientific community can obtain the pertinent sequences from the human genome for \$50 million, or the equivalent of one-quarter of next year's projected funding for the Human Genome Project.

Obviously the Human Genome Project must be considered as two distinct projects: (i) the mapping of the genome and (ii) determining its DNA sequence. The former project is laudatory and worth the money and time invested. The latter project is not cost-effective because the pertinent coding sequences must be obtained from the analysis of messenger RNA transcripts and cannot be deduced solely from the analysis of the genomic sequence.

JOHN H. WEIS

Division of Cell Biology and Immunology,
Department of Pathology,
University of Utah School of Medicine,
Salt Lake City, UT 84132