

The Pleasant Island sediment core provides a continuous record of late Wisconsin environments in southeastern Alaska but by itself does not demonstrate that the Younger Dryas occurred in this region. If a similar climatic signal can be found in lacustrine sediments from other sites in the area, then the existence of the Younger Dryas event will be established for the Pacific Northwest, and a mechanism will be required for transmitting this well-established North Atlantic climatic perturbation around the Northern Hemisphere, if not the globe.

REFERENCES AND NOTES

- W. A. Watts, in *The Lateglacial of North-West Europe*, J. J. Lowe, J. M. Gray, J. E. Robinson, Eds. (Pergamon, New York, 1980), pp. 1–22; D. G. Sutherland, *Quat. Sci. Rev.* **3**, 157 (1984); U. Siegenthaler and U. Eicher, in *Handbook of Holocene Palaeoecology and Palaeohydrology*, B. E. Berglund, Ed. (Wiley, New York, 1986), pp. 407–422.
- S. Björck, P. Sandgren, B. J. Holmquist, *J. Quat. Sci.* **2**, 133 (1987).
- J. Mangerud, in *Abrupt Climatic Change: Evidence and Implications*, W. H. Berger and L. D. Labeyrie, Eds. (D. Reidel, Dordrecht, 1987), pp. 163–171.
- W. F. Ruddiman and A. McIntyre, *Quat. Res.* **16**, 125 (1981).
- W. Dansgaard, J. W. C. White, S. J. Johnsen, *Nature* **339**, 532 (1989).
- W. S. Broecker et al., *Palaeoceanography* **3**, 1 (1988); W. S. Broecker et al., *Nature* **341**, 318 (1989); W. S. Broecker and G. S. Denton, *Geochim. Cosmochim. Acta* **53**, 2465 (1989).
- H. E. Wright, Jr., *Quat. Sci. Rev.* **8**, 295 (1989).
- R. J. Mott, D. R. Grant, R. Stea, S. Ochietti, *Nature* **123**, 247 (1986).
- D. M. Peteet, in *Abrupt Climatic Change: Evidence and Implications*, W. H. Berger and L. D. Labeyrie, Eds. (D. Reidel, Dordrecht, 1987), pp. 185–193; et al., *Quat. Res.* **33**, 219 (1990).
- L. C. K. Shane, *Boreas* **16**, 1 (1987).
- D. Rind, D. Peteet, W. S. Broecker, A. McIntyre, W. F. Ruddiman, *Clim. Dyn.* **1**, 3 (1986).
- V. Markgraf, *Quat. Res.* **31**, 426 (1989).
- J. Jouzel et al., *Nature* **329**, 403 (1987).
- F. Gasse, R. Téhé, A. Durand, E. Gibert, J. Ch. Fontes, *ibid.* **346**, 141 (1990); F. A. Street-Perrott, R. A. Perrott, *ibid.* **343**, 607 (1990).
- J. T. Overpeck, L. C. Peterson, N. Kipp, J. Imbrie, D. Rind, *ibid.* **338**, 553 (1989).
- Edgecumbe tephra from southeast Alaska represent a period of eruption that lasted 500 to 1000 years and ended about 10,500 years ago. The corrected dates bracketing the tephra deposit from Pleasant Island (11,100 and 11,430 years ago) are consistent with this regional correlation, but could be as much as 400 years too old. J. Riehle, unpublished manuscript.
- L. C. Cwynar and W. A. Watts, *Quat. Res.* **31**, 377 (1989).
- D. R. Engstrom and H. E. Wright, Jr., in *Lake Sediments and Environmental History*, E. Y. Haworth and J. W. G. Lund, Eds. (Univ. Minnesota Press, Minneapolis, 1984), pp. 11–67.
- J. E. Armstrong, *Geol. Surv. Can. Bull.* **322**, 1 (1981).
- J. J. Clague, *Geol. Surv. Can. Pap.* **80-35**, 1 (1981).
- R. E. Ackerman, T. D. Hamilton, R. Stuckenrath, *Can. J. Archeol.* **3**, 195 (1979); D. H. Mann, in *Glaciation in Alaska—The Geologic Record*, T. D. Hamilton, K. M. Reed, R. M. Thorson, Eds. (Alaska Geological Society, Anchorage, 1986), pp. 237–265.
- C. J. Heusser, *Quat. Res.* **30**, 284 (1973); R. W. Mathewes, *Can. J. Bot.* **51**, 2085 (1973); C. J. Heusser, *Geol. Soc. Am. Bull.* **85**, 1547 (1974); C. W. Barnosky, *Quat. Res.* **16**, 221 (1981); R. J. Hebda, *Can. J. Bot.* **61**, 3172 (1983).
- R. W. Mathewes and J. J. Clague, *Can. J. Earth Sci.* **19**, 1185 (1982); R. W. Mathewes, *International Union for Quaternary Research, XII International Congress Abstr.* (National Research Council of Canada, Ottawa, 1987), pp. 221.
- K. Chinzai et al., *Mar. Micropal.* **11**, 273 (1987); N. Kallel et al., *Oceanol. Acta* **11**, 369 (1988).
- L. E. Heusser and J. J. Morley, *Quat. Res.* **34**, 101 (1990).
- We thank B. Coffin, M. Noble, J. Janssens, and R. Carstenson for assistance in the field, D. Mann and

G. Streveler for helpful comments on the manuscript, and D. B. Lawrence, the National Park Service, and National Science Foundation Ecology and Climate Dynamics Programs (BSR-8705371, ATM-85114335, ATM-8714122) for support. Limnological Research Center contribution No. 403.

16 August 1990; accepted 19 September 1990

El Niño—Southern Oscillation Displacements of the Western Equatorial Pacific Warm Pool

M. J. McPHADEN AND J. PICAUT

The western equatorial Pacific warm pool (sea-surface temperatures $>29^{\circ}\text{C}$) was observed to migrate eastward across the date line during the 1986–1987 El Niño–Southern Oscillation event. Direct velocity measurements made in the upper ocean from 1986 to 1988 indicate that this migration was associated with a prolonged reversal in the South Equatorial Current forced by a large-scale relaxation of the trade winds. The data suggest that wind-forced zonal advection plays an important role in the thermodynamics of the western Pacific warm pool on interannual time scales.

THE SOUTHERN OSCILLATION IS A large-scale seesaw in atmospheric pressure between the eastern subtropical Pacific and the maritime land masses of Australia and Indonesia (1, 2). Pressure differences between these two regions drive easterly trade winds, which are the principal forcing function of tropical ocean circulation. The trade winds converge in the western Pacific in regions of deep atmospheric convection and high rainfall rates. A drier air mass then returns eastward aloft before descending in the subtropical high region of the eastern Pacific. Periods when the pressure is unusually high in the west and low in the east are associated with El Niño, an oceanic phenomenon characterized by weakened trade winds in the central and western Pacific and sea-surface temperatures (SSTs) that are warmer than usual from the coast of South America to west of the date line (3). El Niño–Southern Oscillation (ENSO) episodes occur on an irregular cycle of about 2 to 10 years, the most recent of which occurred in 1986 to 1987 (4, 5). Although the origins of ENSO may be traced to the tropics, its manifestations are felt worldwide through disruptions in the atmospheric general circulation and associated weather patterns (1, 6).

The western Pacific warm pool, defined by SSTs $>29^{\circ}\text{C}$, has recently taken on a

special significance in ENSO studies (7, 8). The SSTs in the western equatorial Pacific are the highest in the world ocean. Such high temperatures lead to tremendous variability in deep convection because the ability of the atmosphere to hold water vapor increases nonlinearly with temperature (9). During ENSO, the warm pool and deep convection associated with converging easterlies are observed to migrate eastward across the date line. This migration of the warm pool has generally been attributed to wind-driven zonal current advection, on the basis of model studies (7, 10, 11) and empirical mass-balance studies (12–14). However, until recently there have been no direct current measurements west of the date line suitable for testing this hypothesis. In this report, we present moored and shipboard velocity measurements for 1986 to 1988 and show that the western Pacific warm pool is responding to current variations in a manner consistent with inferences drawn from models, theory, and earlier empirical studies.

To place these measurements in perspective, we review the broad outlines of air-sea interaction on ENSO time scales as discussed in the context of several simple coupled ocean-atmosphere models (15). In these models, warm tropical SST anomalies generate anomalous deep atmospheric convection fed by convergence of surface winds. This surface convergence leads to a significant weakening of the trade winds west of the warm SST anomalies. The ocean responds rapidly to this weakening: surface currents locally accelerate eastward, and

M. J. McPhaden, Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, 7600 Sand Point Way Northeast, Seattle, WA 98115. J. Picaut, Groupe SURTROPAC, Institut Français de Recherche Scientifique pour le Développement en Coopération (ORSTOM), Nouméa, New Caledonia.

large-scale equatorial Kelvin and Rossby waves radiate out from the directly forced region to reinforce and spread the warm SST anomaly across the basin (16). This couples the ocean to the atmosphere in a positive feedback loop, wherein westerly wind anomalies reinforce the eastward currents, and this process in turn leads to further warming. As this feedback continues, the thermocline rises in the west and descends in the east to compensate for the mass of water drained from the western Pacific. Warm SST anomalies continue to grow and persist for 12 to 18 months, until easterly trade winds return to their full strength and the event terminates.

In 1986, maximum SSTs ($>29.5^{\circ}\text{C}$) were centered near 165°E (Fig. 1) during boreal summer (June, July, and August), and the winds within about 10° of the equator were

near normal. One year later, during the height of the 1986–1987 ENSO, the warmest waters in the equatorial Pacific were found between 160°W and 180°W , and over this warm-water pool there was a strong anomalous convergence of surface winds. This convergence was associated with anomalous deep atmospheric convection over the same range of longitudes (4) and weakened trade winds between 160°E and 160°W within 5° of the equator (17). In boreal summer 1988, the warmest surface waters were west of 160°E between the equator and 10°N , and a well-developed cold tongue with minimum temperatures on the equator formed all the way west to 165°E . The cold tongue is usually confined to east of about 160°W (18), and its presence in the western Pacific in 1988 is consistent with the occurrence of stronger than normal easterly trade winds west of 160°W .

The zonal migration of the warmest surface waters can be related to the direction and magnitude of flow across 165°E , as determined from shipboard and moored velocity measurements made during 1986 to 1988 (19, 20). In particular, meridional sections of zonal velocity between 10°N and 10°S (Fig. 2) show a dramatic interannual reversal of flow in the upper 100 m of the water column. Before the ENSO, in June 1986, flow was westward between 8°S and 3°N in the South Equatorial Current at speeds up to 0.6 m s^{-1} and eastward between 3°N and 7°N in the North Equatorial Countercurrent at speeds up to 0.6 m s^{-1} (21). However, 1 year later, flow was eastward between about 7°S and 6°N at speeds

up to 0.8 m s^{-1} . The volume transport associated with currents in the upper 100 m between 5°N and 5°S was 11 Sv to the west in 1986 and 47 Sv to the east in 1987 ($1\text{ Sv} = 10^6\text{ m}^3\text{ s}^{-1}$). This is a difference of 58 Sv, which is larger than the annual mean transports of all the major zonal currents in the western and central equatorial Pacific (20, 22). The large anomalous eastward mass flux in 1987 drained the warm surface layer in the western Pacific and thus elevated the upper thermocline by 20 to 40 m (Fig. 2, D and E) and depressed sea level by 20 to 30 cm (4, 5). After the ENSO, in boreal summer 1988, the westward South Equatorial Current returned to its pre-ENSO strength and displaced the warm surface layer to the west across 165°E at a rate of 7 Sv. A depression of the upper thermocline and an increase in the volume of water having a temperature $>29^{\circ}\text{C}$ relative to 1987 values was associated with this westward volume transport (Fig. 2, E and F).

These transport estimates are based on only three meridional sections, each separated by approximately 1 year. The high variability of the near-surface equatorial ocean on time scales shorter than this could affect the interpretation of our measurements in terms of interannual variability. However, our transport estimates are the same magnitude as indirect estimates of yearly averaged volume transports from the western to the eastern Pacific inferred from island sea-level data during the 1976 ENSO [27 Sv (13)] and the 1982 to 1983 ENSO [40 Sv (14, 23)]. The equatorial velocity profiles are also consistent with averaged time-series




Fig. 1. Contours of SST for (A) June to August 1986, (B) June to August 1987, and (C) June to August 1988. Contour interval is 1°C for temperatures $\leq 29^{\circ}\text{C}$. The 29.5°C isotherm is dashed and water warmer than 29.5°C is shaded. Also plotted are wind pseudostress anomaly vectors (defined as $|U|U$, where U is vector wind speed) for each period. The SST analyses are from the Climate Analysis Center of the National Meteorological Center; pseudostress data are from (28). The square indicates the location of a current meter mooring at 0° , 165°E ; the vertical line between 10°N and 10°S , 165°E indicates the section along which measurements shown in Fig. 2 were made.





Fig. 2. Meridional sections of zonal velocity (A, B, and C) and temperature (D, E, and F) for June 1986, July 1987, and June 1988 along 165°E . Westward flow and temperatures $>29^{\circ}\text{C}$ are shaded.

measurements collected from a current meter mooring at 0°, 165°E during boreal summers from 1986 to 1988. The mooring data (Fig. 3A) have the same character as the shipboard velocity profiles at the equator and show that flow was eastward in the surface layer during 1987 but westward before and after the ENSO event. The differences are largest in the upper 100 m (for example, 0.6 m s^{-1} between 1987 and 1988 at 50 m) and diminish into the thermocline (24). The mooring data, like the shipboard data, indicate that the thermocline shoaled in 1987 (Fig. 3C), as is consistent with the notion that a large eastward volume flux drained the western Pacific of its warm surface layer.

Further evidence supporting this advective hypothesis is given by estimates of the horizontal distance over which water parcels would be displaced. These estimates were obtained by integrating the velocity measurements from the 165°E mooring in time. The resultant time series of positions differs from actual fluid-parcel trajectories because of inhomogeneities in the horizontal flow field and because of the neglect of vertical velocities. However, the method provides a useful indicator of lateral displacements of water mass in lieu of true Lagrangian measurements. The longest continuous moored velocity record available for this purpose at this station is from 13 December 1986 to 31 December 1988 at a depth of 50 m. Meridional velocities are relatively small (Fig. 3B) so that the largest displacements are in the zonal direction. These displacements, superimposed on contours of SST in Fig. 4, are comparable in magnitude and duration to displacements derived from a model of the ENSO (10) and indicate that water parcels could reach the central Pacific by October 1987 before returning westward.

The 29°C isotherm at the eastern edge of the warm pool was displaced to 150°W during July and August 1987 before retreating to near 160°E in mid-1988 (Fig. 4). Displacements of this isotherm and those inferred from the mooring data would closely correspond to one another if (i) zonal advection were the primary determinant in the surface heat balance; and (ii) the velocity field were uniform over distances separating the 29°C isotherm from the mooring at 165°E. In support of the zonal-advection hypothesis, excursions of the 29°C isotherm resemble those inferred from the mooring velocity data collected when the 29°C isotherm was near 165°E in 1988. Conversely, when the 29°C isotherm was located east of the date line in 1987, its excursions tended to be smaller in amplitude than those inferred from the mooring data, and they tended to lead excursions inferred from the

Fig. 3. Profiles of (A) zonal velocity; (B) meridional velocity; and (C) temperature averaged during 1 June to 2 July 1986, 1 June to 31 August 1987, and 1 June to 31 August 1988. Instrument depths are indicated on the left axes for 1986 and on the right axes for 1987 and 1988.



mooring data by about 2 months. The lead-lag relation, if interpreted in terms of water-mass displacements, implies a westward phase propagation of about 0.8 m s^{-1} , comparable to that expected for large-scale equatorial Rossby waves. The smaller zonal displacements of the 29°C isotherm relative to those inferred from the mooring data can be interpreted as a damping of SST anomalies by loss of latent heat to the atmosphere (25). In addition, one would expect that zonal displacements on the equator (as computed from the mooring data) would have been larger than the average displacement between 5°N and 5°S (the interval over which SST was averaged in Fig. 4) because of the tendency for zonal currents off the equator to be deflected meridionally by Coriolis forces.

Earlier analyses have shown that the zonal-velocity variations at 0°, 165°E are significantly correlated with local wind variations (5). Variations in local winds do not account for all of the observed zonal velocity variance, however, and the foregoing discussion suggests that velocities associated with remotely generated Rossby waves may also be important. Direct evidence for equatorial Rossby waves has been obtained from basin-wide satellite sea-level measurements in the Pacific during 1986 and 1987 (26). Moreover, wind-forced westward-propagating Rossby waves are an important dynamical component in nearly all ocean models of ENSO (27). In these models, westerly wind anomalies (like those shown in Fig. 1b) generate Rossby waves that advect the surface layer eastward and elevate the thermocline in the western Pacific in a manner consistent with our moored and shipboard velocity and temperature data.

In summary, we attribute zonal displacements of the western equatorial Pacific warm pool observed from 1986 to 1988 to wind-forced variations in current velocity and transport in the upper ocean. In addition, our data can be interpreted to imply that remotely forced equatorial Rossby waves contribute to these current variations,




Fig. 4. Averaged SST from 5°N to 5°S as a function of longitude in the Pacific from 1986 to 1988. Superimposed as a dashed line are estimated water-mass displacements from the current meter mooring at 0°, 165°E.

consistent with our present understanding of ocean dynamics. However, inferences about the role of Rossby waves must be considered tentative on the basis of the data used in this study, because our velocity measurements derive from only a single meridian (165°E). More complete oceanic data sets than are currently available are needed to understand fully the dynamics of current variations like those we have observed.

REFERENCES AND NOTES

1. E. M. Rasmusson and J. M. Wallace, *Science* **222**, 1195 (1983).
2. S. G. H. Philander, *Nature* **302**, 295 (1983).
3. E. M. Rasmusson and T. H. Carpenter, *Mon. Weather Rev.* **110**, 354 (1982); M. A. Cane, *Science* **222**, 1189 (1983).
4. V. E. Kousky and A. Leetmaa, *J. Climate* **2**, 254 (1989).
5. M. J. McPhaden et al., *J. Phys. Oceanogr.* **20**, 190 (1990).
6. C. F. Ropelewski and M. S. Alpert, *Mon. Weather Rev.* **114**, 2352 (1986); *ibid.* **115**, 1606 (1987); G. N. Kiladis and H. F. Diaz, *J. Climate* **2**, 1069 (1989). R. T. Barber and F. P. Chavez [*Science* **222**, 1203 (1983)] also discussed the biological consequences of El Niño on the oceanic ecosystems of the eastern equatorial Pacific.
7. A. E. Gill and E. M. Rasmusson, *Nature* **306**, 229 (1983).
8. T. N. Palmer and D. A. Mansfield, *ibid.* **310**, 483 (1984); N. Nicholls, *Mon. Weather Rev.* **112**, 424 (1984).
9. K. M. Lau and P. H. Chan, *J. Atmos. Sci.* **45**, 506 (1986).
10. A. E. Gill, *J. Phys. Oceanogr.* **13**, 586 (1983).
11. R. Seager, *ibid.* **19**, 419 (1989).
12. K. Wyrtki, *ibid.* **5**, 572 (1975).

13. ———, *ibid.* 9, 1223 (1979).
14. ———, *J. Geophys. Res.* 90, 7129 (1985).
15. S. G. H. Philander *et al.*, *J. Atmos. Sci.* 41, 604 (1984); P. S. Schopf and M. J. Suarez, *ibid.* 45, 550 (1988); D. Battisti, *ibid.*, p. 2889; S. E. Zebiak and M. A. Cane, *Mon. Weather Rev.* 115, 2262 (1987).
16. These waves have zonal scales of $\sim 10^3$ km and meridional scales of $\sim 10^2$ km. The fastest Kelvin waves propagate eastward at speeds of 2 to 3 m s⁻¹, whereas the fastest Rossby waves propagate westward at one-third the Kelvin wave speed. It takes about 2 months for a Kelvin wave to traverse the zonal extent of the Pacific basin.
17. Wind variations within about 5° to 7° of the equator are the most important in the evolution of ENSO events on the basis of several modeling studies: J. P. McCreary, *J. Phys. Oceanogr.* 6, 632 (1976); M. A. Cane, *ibid.* 14, 586 (1984); D. E. Harrison, *ibid.* 19, 691 (1989).
18. K. Wyrtki, *ibid.* 11, 1205 (1981).
19. Two cruises per year along 165°E have been made both by the French SURTROPAC program since 1984 (20) and by the United States/People's Republic of China bilateral air-sea interaction program since 1986. A current meter mooring has been maintained at 0°, 165°E since January 1986 as part of the US/PRC program (5).
20. T. Delcroix *et al.*, *J. Phys. Oceanogr.* 17, 2248 (1987).
21. These meridional sections of velocity were obtained with an Aanderaa-Tareq type profiling current meter that freely hangs on a cable under a drifting buoy. Currents are calculated relative to flow at 600 m, which is assumed to be negligible. The errors inherent in this type of measurement are approximately 0.15 to 0.20 m s⁻¹, on the basis of a comparison of seven profiles taken within 2 nautical miles of the current meter mooring in June 1986 and July 1988. We expect that the depth-coherent part of this error will lead to uncertainties in transport estimates in the upper 100 m of approximately 15 Sv between 5°N and 5°S.
22. K. Wyrtki and B. Kilonsky, *J. Phys. Oceanogr.* 14, 242 (1984).
23. G. Meyers and J.-R. Donguy [*Nature* 312, 258 (1984)] used expendable bathythermograph data and the geostrophic approximation to estimate that about 10 Sv of anomalous eastward transport occurred in the North Equatorial Countercurrent between 3°N and 9°N during the 1982–1983 ENSO.
24. The Equatorial Undercurrent at depths of 150 to 200 m was weak during the 1986–1987 ENSO. It eventually disappeared in October to November 1987 because of prolonged anomalous westerly wind forcing that led to a reversal of the zonal pressure gradient in the thermocline (5).
25. C. S. Ramage *et al.*, *Univ. Hawaii Tech. Rep. UHMET 80-03* (1980); R. K. Reed, *Nature* 322, 6078 (1986).
26. T. Delcroix *et al.*, *J. Geophys. Res.*, in press.
27. See, for example, J. P. McCreary, Jr., and D. L. T. Anderson, *Mon. Weather Rev.* 112, 934 (1984); A. Busalacchi *et al.*, *J. Geophys. Res.* 88, 7551 (1983); S. E. Zebiak, *J. Phys. Oceanogr.* 19, 475 (1989); N. E. Graham and W. B. White, *Science* 240, 1293 (1988).
28. D. M. Legler and J. J. O'Brien, *Intergov. Oceanogr. Comm. Tech. Ser.* 4, 11 (1988).

5 June 1990; accepted 7 September 1990

Preceramic Occupations in the Orinoco River Valley

WILLIAM P. BARSE

Two sites in the Orinoco Valley containing preceramic remains from excavated contexts are described. Radiocarbon dating and stylistic comparisons indicate that the northern tropical lowlands were inhabited at the onset of the Holocene, suggesting a time depth of 9000 years before the present for tropical forest-savanna adaptations in northern South America.

PRECERAMIC SITES HAVE LONG BEEN assumed to exist in the northern tropical lowland interior of South America, although evidence has been restricted to surface manifestations in Guyana (1) and Venezuela (2). Recent excavations conducted along the Orinoco River in the vicinity of Puerto Ayacucho, Territorio Federal Amazonas, Venezuela, revealed preceramic occupations in well-stratified contexts at two sites, Culebra and Provincial (Fig. 1). These two sites were located during the course of a project aimed at developing a regional cultural sequence for the Orinoco between the Atures Rapids and Meta River (3). Although both contained ceramic components, only the preceramic occupations will be discussed. These early Holocene sites demonstrate that tropical forest-savanna adaptations in the interior lowlands of northern South America date to at least 9000 years B.P. The preceramic phases are followed by a lengthy ceramic period. Excavations at these two sites and three others in the area documented an early series of Barrancoid phases beginning at ~ 1000 B.C., followed in time by Saladoid (A.D. 200 to 300), Ronquin (A.D. 500 to 700), and

Araucuin (A.D. 1200 to 1400) related phases (3).

The Culebra site is located near the confluence of the Cataniapo River with the Orinoco, 12 km south of Puerto Ayacucho. The Cataniapo enters into the Orinoco River midway through the Atures Rapids, the upper limit of large boat traffic on the river. The site is situated on an alluvial terrace bordered by sporadic granite outcrops. It is in a woody savanna setting surrounded by tropical gallery forest. Two preceramic components, Atures I and Atures II, were encountered during the test excavations at the Culebra site.

The earliest component, Atures I, was encountered in a 1 by 2 m excavation unit in the northern portion of the site between 85 and 100 cm below the surface. Contextually, it was in a sandy clay loam paleosol, the occupation coinciding with the upper boundary of the soil horizon. Forty centimeters of sterile sandy loam separated Atures I from the earlier of the two ceramic components in the unit (Fig. 2A). The Atures I assemblage is characterized by two flake scrapers, a core fragment, and waste flaking debris (Table 1A). The predominant lithic material used was transparent crystalline quartz, although several quartzite flakes were also present. The scrapers show re-

touch or use wear along either the lateral margin, distal end, or both (Fig. 3, A and B). The quartz was obtained locally from cobbles in the bed load of the Orinoco River. Many flakes, one flake scraper, and the core fragment exhibit cobble cortex.

Atures II, the second preceramic component, was encountered in two separate excavations: a 2 by 2 m unit in the southern part of the site and a 1 by 2 m unit along its eastern edge. It was in a buried A horizon just below the earlier of the two ceramic components at the site, stratified above the paleosol containing the Atures I component (Fig. 2A). Atures II contained crystal quartz




Fig. 1. Study area and its location within Venezuela.