An Ancient Molecular Mechanism for Establishing Embryonic Polarity?

Judith Kimble

Worms, butterflies, and chimpanzees all have the same body axes—head and tail, front and back, and left and right sides. How are these axes established during development? Is there a single molecular map used by most metazoan embryos or have similar coordinates been achieved during evolution by diverse routes? A comparison of the mechanisms that establish body axes in distantly related organisms can begin to answer this fundamental question.

Progress has been spectacular in dissecting the molecular mechanisms that control

axis formation in one organism, the fruit fly Drosophila (1), but much less is known about these mechanisms in other organisms. Recently, similarities have become apparent between mechanisms that govern early embryonic polarity in the nematode Caenorhabditis elegans and in Drosophila (2). In both species, translational repression at the posterior pole establishes asymmetry along the anterior-posterior axis (2-4). Nematodes and insects diverged at least 600 million vears ago-when metazoans first made their appearance in the fossil record (3, 4)—and

so such localized translational repression may be an ancient molecular mechanism for specification of one body axis, the anterior-posterior axis that runs from head to tail. In Drosophila, asymmetries that are established in the oocyte generate localized protein gradients and the overall body plan during early embryogenesis (1). Formation of the anterior-posterior axis relies on localized maternal mRNAs-bicoid at the anterior and nanos at the posterior pole. These localized mRNAs are translated soon after fertilization to create gradients of their protein products in the early embryo. Bicoid encodes a transcriptional activator, which turns on zygotic expression of another gene, hunchback, in the anterior region; nanos, by contrast, represses translation of maternal hunchback mRNA in the posterior region

and leads to its destruction. The gradient of hunchback therefore is formed by transcriptional activation at the anterior pole and translational repression at the posterior pole. The hunchback gradient, in turn, triggers a cascade of transcription factors that specify patterning along the anteriorposterior axis.

Early embryogenesis in *Drosophila* takes place within a large cytoplasm that is shared by many nuclei, the syncytial blastoderm. By contrast, most animal embryos begin life with a series of cleavage divisions



Establishment of embryonic polarity during metazoan development.

that separate the zygote's cytoplasm into distinct cellular domains. Nematode embryos, for example, divide unequally at the first cleavage division, producing a large anterior daughter and a small posterior daughter. The anterior daughter generates mostly nerve and epidermis, whereas the posterior daughter forms muscle, gut, and germ cells (5). Amphibian embryos possess animal and vegetal poles at fertilization, which subsequently give rise to cells with distinct fates. These dramatic differences in embryonic morphology are commonly thought to reflect fundamental differences in the mechanisms that control embryonic patterning. One mechanism for generating asymmetry along the anterior-posterior axis appears to be shared by worms and flies. The glp-1 protein encoded by glp-1, a membrane receptor required for inductive cell interactions in the early C. elegans embryo (6), is first observed in the anterior, but not the posterior, blastomere of late two-cell

SCIENCE • VOL. 266 • 28 OCTOBER 1994

embryos; this glp-1 asymmetry is maintained in 4- to 28-cell embryos (2). The anterior-posterior asymmetry of glp-1 protein may be crucial for the establishment of asymmetrical patterns of cell-cell interaction, which in turn specify dorsal-ventral and left-right axes of the C. elegans embryo (7-10). In contrast to glp-1 protein, glp-1 maternal mRNA is distributed uniformly. which eliminates two models for control of glp-1 protein asymmetry-localization of glp-1 maternal RNA and localized control of glp-1 RNA stability (2). In fact, the glp-1 pattern is controlled translationally, by the glp-1 3' untranslated region (3'UTR), the site of translational regulatory elements in many maternal mRNAs (11). Reporter RNAs carrying a naïve 3'UTR or no 3'UTR are translated in an unregulated fashion, whereas reporter RNA with the glp-1 3'UTR is translated in a pattern that is virtually the same as that of endogenous glp-1 protein (2). Specific translational

PERSPECTIVES

regulatory elements are found within regions of the glp-1 3'UTR. Removal of 125 nucleotides from the 3' end of the glp-1 3'UTR permits translation during oogenesis. whereas removal of 61 nucleotides from the center of the glp-1 3'UTR permits translation in both anterior and posterior blastomeres, starting at the four-cell stage. Therefore, one type of cisacting regulatory element appears to be located at the 3' end of the 3'UTR for repression of translation until the two- to four-cell stage; another type of element appears to be located in the central re-

gion for repression in posterior cells.

Regulation of glp-1 asymmetry in the early C. elegans embryo bears a remarkable resemblance to the establishment of hunchback asymmetry in the early Drosophila embryo (see figure). Repression in the posterior region restricts translation of both hunchback and glp-1 maternal transcripts to the anterior region of their respective embryos. In addition, similar cis-acting regulatory elements may be involved. In Drosophila, translational control depends on a short bipartite sequence motif, called the nanos response element (NRE), which resides in the hunchback 3'UTR (12). A trans-acting regulator that functions through the NREs has not been identified biochemically, but an excellent candidate is the product of the nanos gene (13). In C. elegans, a bipartite sequence that is strikingly similar to the Drosophila NRE occurs within that central region of the glp-1 3'UTR required for translational repression in the posterior blastomere (2).

The author is in the Laboratory of Molecular Biology, Howard Hughes Medical Institute, University of Wisconsin, Madison, WI 53706, USA.

The similarity between Drosophila and C. elegans early embryos extends beyond the parallels between hunchback and glp-1 translational regulation. Both embryos contain cytoplasmic "granules" in the posterior region of the zygote that are segregated ultimately to germ cell precursors during embryogenesis-the polar granules of Drosophila (14) and the P granules of C. elegans (15) (see figure). In Drosophila, maternal nanos RNA is associated with polar granules; perhaps in C. elegans, a homolog of nanos is associated with P granules.

What about vertebrates? Does translational repression in the posterior cytoplasm establish embryonic polarity in these "higher" animals? A hint that this mechanism may indeed function in vertebrates comes from the identification of a maternal transcript that encodes a nanos-like protein called Xcat-2 in Xenopus embryos (16). Although the function of Xcat-2 is unknown, its location at the vegetal pole suggests a role in early pattern formation. Furthermore, a "germ plasm" exists in the vegetal cytoplasm of amphibian embryos that may be analogous to P granules and polar granules of worms and flies (17). Over the past decade, a handful of molecular mechanisms have been implicated in the patterning of Drosophila, C. elegans, and Xenopus embryos (1, 18, 19). On the basis of the diversity of these mechanisms, the prevailing view has been that each embryo has differentially employed a handful of common molecular mechanisms to create its own coordinate system. For example, localized transcriptional activators are utilized for patterning of both C. elegans and Drosophila early embryos (20-23), but the mechanisms for localization, types of DNA binding protein, and specified fates are not obviously similar.

By contrast, the molecular parallels between hunchback and glp-1 regulation suggest the existence of an ancient mechanism for creating asymmetric patterns of gene expression in early embryos (see figure). This mechanism is predicted to depend on a trans-acting regulator similar to nanos and to act through cis-acting sequences similar to NREs in the 3'UTRs of maternal transcripts. If this molecular machinery regulates polarity in embryos as diverse as worms, flies, and frogs, it becomes plausible that it influences axis formation in all animal embryos, including mammals. "Molecular tinkering" (24) may then come into play to reinforce this primitive patterning control and to derive other axes from it.

Research in Drosophila has pioneered our understanding of the molecular mechanisms that can establish the body axes in an early embryo. Now, phylogenetic comparisons will tell us which mechanisms are primitive and which have evolved to reinforce, modify, or extend the underlying map. Are the controls that localize translational repression conserved? Are polar granules the ancient seat of pattern governance? What links the early controls of axis formation to the later controls of homeobox genes, a highly conserved system that specifies individual regions along the anterior-posterior axis of all known metazoa (25)? The hunchback protein is a transcriptional regulator that resides at the top of a cascade of transcriptional regulators, whereas the glp-1 protein is a membrane receptor that directs inductive interactions. Clearly, these distinct modes of regulation must converge to control expression of homeobox genes. How similar are the mechanisms of convergence? Answers to these questions, among the most fundamental of all developmental biology, may be waiting around the corner.

References

- 1. D. St. Johnston and C. Nüsslein-Volhard, Cell 68, 201 (1992)
- 2. T. C. Evans et al., ibid. 77, 183 (1994).

- 3. S. C. Morris, Nature 361, 219 (1993).
- 4. A. Sidow and W. K. Thomas, Curr. Biol. 4, 596 (1994).
- 5. J. E. Sulston et al., Dev. Biol. 100, 64 (1983).
- 6. J. R. Priess, H. Schnabel, R. Schnabel, Cell 51, 601 (1987).
- 7. C. C. Mello, B. W. Draper, J. R. Priess, *ibid.* 77, 95 (1994).
- 8. H. Hutter and R. Schnabel, Development 120, 2051 (1994).
- 9. S. E. Mango et al., ibid., p. 2305.
- I. P. G. Moskowitz, S. B. Gendreau, J. H. 10. Rothman, ibid., p. 3325.
- 11. M. Wickens, Ed., Semin. Dev. Biol. 3 (1992)
 - 12. R. P. Wharton and G. Struhl, Cell 67, 955 (1991).
- 13. C. Wang and R. Lehmann, ibid. 66, 637 (1991). 14. A. P. Mahowald, J. Exp. Zool. 176, 329 (1971).
- 15. S. Strome and W. B. Wood, Cell 35, 15 (1983)
- 16. L. Mosquera et al., Development 117, 377 (1993).
- 17. A. W. Blackler, Dev. Biol. 14, 330 (1966).
- 18. W. B. Wood and L. G. Edgar, Trends Genet. 10,
- 49 (1994).
- 19. D. A. Melton, Science 252, 234 (1991).
- W. Driever and C. Nüsslein-Volhard, Cell 54, 83 20 (1988).
- 21. _____, Nature 337, 138 (1989).
 22. B. Bowerman *et al.*, Cell 74, 443 (1993).
- 23. B. Bowerman, B. A. Eaton, J. R. Priess, ibid. 68. 1061 (1992).
- F. Jacob, The Possible and the Actual. The Jessie 24 and John Danz Lectures (Pantheon, New York, 1982)
- 25. C. Kenyon, Cell 78, 175 (1994).

The Embryonic Vertebrate Forebrain: The Prosomeric Model

John L. R. Rubenstein, Salvador Martinez, Kenji Shimamura, Luis Puelles

The mammalian forebrain—the cerebral cortex, basal ganglia, hypothalamus, and thalamus-is the seat of higher cognitive functions. How much of forebrain development and structure is controlled by a genetic program? Although at the later stages of development incoming synaptic information from the thalamus has been shown to influence patterning in the neocortex (1), at early embryological stages a specific set of newly discovered genes pattern the brain into a highly organized structure-before synaptic influences are present. Furthermore, the primordia of major structural elements, such as the thalamus, are segregated by cellular boundaries that are aligned parallel to the topologically transverse and longitudinal axes of the neural tube. Specific combinations of genes that are ex-

SCIENCE • VOL. 266 • 28 OCTOBER 1994

pressed in these domains direct the unique development of each region. Finally, the organization of the forebrain indicates that it is a segmental structure.

The basic morphogenetic unit of embryonic insects is a transverse domain, or segment (2). The identity of each segment is determined by its position along the anterior-posterior axis and is controlled by the expression of the homeobox segment identity genes (3). These genes encode transcriptional regulators of specific sets of target genes, which define the unique developmental pathway of each individual segment.

It is widely held that this paradigm may apply to the organization of the somitic mesoderm (the vertebral column), the rhombencephalon (hindbrain), and the branchial arches of vertebrates. This view is based on the existence of homologs of the homeobox segment identity genes in vertebrates (the Hox genes) (3) as well as the metameric (segmental) morphological and histological features of these structures. This hypothesis has been confirmed in part by the use of genetic manipulations that al-

J. L. R. Rubenstein and K. Shimamura are in the Nina Ireland Laboratory of Developmental Neurobiology, Center for Psychiatry and Neurobiology, Programs in Neuroscience and Developmental Biology, 401 Parnassus Avenue, University of California, San Francisco, CA 94143-0984, USA. S. Martinez and L. Puelles are in the Department of Morphological Sciences, Faculty of Medicine, University of Murcia, Murcia, Spain 30100.