

heavy exposures 20 to 70 years ago. The geological distribution of tremolite within the Quebec chrysotile ore body may well vary in time and place and, at present levels of environmental control, any mesothelioma risk from exposure in either area A or B would be far below the limits of epidemiological detection.

**J. Corbett McDonald
Alison D. McDonald**

National Heart and Lung Institute,
London University,
Royal Brompton Hospital,
London SW3 6LY, United Kingdom

References

1. J. C. McDonald, F. D. K. Liddell, A. Dufresne, A. D. McDonald, *Br. J. Ind. Med.* **50**, 1073 (1993).
2. P. S. Sébastien, J. C. McDonald, A. D. McDonald, B. Case, R. Harley, *ibid.* **46**, 180 (1989).


Corrections and Clarifications

In the report "GDNF: A potent survival factor for motoneurons present in peripheral nerve and muscle" by C. E. Henderson *et al.* (11 Nov., p. 1062), the seventh author's name should have been "Laura Simmons." The affiliation for authors H. S. Phillips, M. Armanini, and Aron Rosenthal should have been "Department of Neuroscience, Genentech,

South San Francisco, CA 94080-4990, USA." The affiliation for B. Moffet and R. A. Vandlen should have been "Department of Protein Chemistry, Genentech," and that for Laura Simmons should have been "Department of Cell Genetics, Genentech." Alun M. Davies is at the University of St. Andrews, St. Andrews, Fife, Scotland KY16 9AJ, United Kingdom, and Vassilis E. Koliatsos is in the Departments of Pathology, Neurology, and Neuroscience and the Neuropathology Laboratory, Johns Hopkins University School of Medicine, Baltimore, MD 21208, USA. Beginning on line 19 of the first column on page 1064, the text should have read, "The average volume of the surviving motoneurons in the facial nuclei that received GDNF was $5186 \pm 308 \mu^3$, close to that in the control nuclei ($6081 \pm 105 \mu^3$) ($n = 3$). In contrast, although BDNF and NT-4/5 both enhanced motoneuron survival to a similar degree as did GDNF (2, 3), neither of them prevented the axotomy-induced shrinkage of facial motoneurons. In BDNF and NT-4/5-treated nuclei, the volume of the surviving motoneurons was only $65.5 \pm 4\%$ and $63.3 \pm 2.6\%$ of the control nuclei, respectively ($n = 3$) (14)."

Throughout the Research Article "Autoproteolysis in hedgehog protein biogenesis" by John J. Lee *et al.* (2 Dec., p. 1528) the word "disc" was incorrectly spelled "disk." In figure 4C (p. 1531), lane 9 should have been labeled "flu227," not "flu408."

We'll Help You Build A New Lab





Let the free Science product information service put you in touch with the vendors whose products you will need.

Simply write us a letter stating the specifics about your proposed lab and the instruments and supplies you need. We will do the rest. Write to:

**SCIENCE Magazine
New Lab Service Department
1333 H Street, N.W., Rm 830
Washington, DC 20005**

The Global Weekly of Research
SCIENCE

NEW! EndNote Plus 2.0



Bibliographies made easy!

More than 75,000 researchers use EndNote to create bibliographies and organize references. Now version 2 makes this even easier with great new features such as term lists, new search options, and 240 journal styles! And to make it even easier, EndLink lets

you import references from more than 100 sources without typing a word!

Niles & Associates, Inc.
800 Jones Street
Berkeley, CA 94710

Voice: 800-554-3049
Fax: 510-559-8683
email: endnote@aol.com

Mail to Niles & Associates, Inc., 800 Jones St., Berkeley, CA 94710

☐ Yes! send me information about EndNote for...

☐ Mac

Version 2.0

☐ Windows

Under Development

☐ DOS

Version 1.3.3
2.0 not available

Name _____

Address _____

City/State/Zip _____