National Cancer Institute, Bethesda, MD 20892, USA E-mail: rss@nchgr.nih.gov

Richard Peters

Howard Hughes Medical Institute Fellow, Harvard Medical School, and Department of Medicine, Massachusetts General Hospital, Boston, MA 02114, USA E-mail: rpeters@vp3.med.harvard.edu

Cockroach Immunity

Although I was pleased to see our work on the American cockroach cited in the article by Virginia Morell about female immune responses (News, 11 Aug., p. 773), I feel I must come to the defense of the male cockroach, which was characterized as less than able to withstand toxic challenges of honeybee venom. In fact, 80% of male roaches will survive a lethal dose of the venom delivered 2 weeks after immunization with a toxoid preparation (1). However, as Morell correctly implies, female roaches do indeed have superior immune responses compared with those of males. In fact, we have found that the female roach is so much more adept at generating a protective response that it requires a second immunization of males to even begin to match the response levels found in females receiving only a single immunization.

Richard D. Karp

Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45221, USA

References

L. E. Duwel-Eby et al., in Immunology of Insects and Other Arthropods, A. P. Gupta, Ed. (Academic Press, New York, 1991), pp. 385–402.

Corrections and Clarifications

The Random Sample item "Diagnostic imagery for Florida" (8 Sept., p. 1343) should have stated that the accompanying map shows the Caloosahatchee River flowing out of Lake Okeechobee (to the Gulf of Mexico), not into

In the response by O. Steinbock and K. Showalter to a technical comment by D. Gareth Williams ("Minimal path algorithms," 21 July, p. 418), the word "his" was inadvertently inserted by Science before the word "co-workers" in the second sentence. The authors originally wrote, "We wish to emphasize that path finding from reaction-diffusion waves, which was suggested by Babloyantz and co-workers in 1991 (2) [J. A. Sepulchre, A. Babloyantz, L. Steels, in Proceedings of the International Conference on Artificial Neural Networks, T. Kohonen, K. Makisara, O. Simula, J. Kangas, Eds. (Elsevier, Amsterdam, 1991), pp. 1265-1268; J. A. Sepulchre and A. Babloyantz, in Chemical Waves and Patterns, R. Kapral and K. Showalter, Eds. (Kluwer, Dordrecht, 1995), pp. 191-217], represents a mechanism by which physical and biological systems might optimize transit times and distances." Science regrets the error.

References related to citation 2 include J. A. Sepulchre and A. Babloyantz, Phys. Rev. E 48, 187 (1993); A. Babloyantz and J. A. Sepulchre, Physica D 49, 52 (1991); and J. A. Sepulchre and A. Babloyantz, Phys. Rev. Lett. 66, 1314 (1991).

Letters to the Editor

Letters may be submitted by e-mail (at science_letters@aaas.org), fax (202-289-7562), or regular mail (Science, 1333 H Street, NW, Washington, DC 20005). Letters will not be routinely acknowledged. Full addresses, signatures, and daytime phone numbers should be included. Letters should be brief (300 words or less) and may be edited for reasons of clarity or space. Beginning in October 1995, our previous policy of consulting with all letter authors before publication will be discontinued.

