

than females.) This assumption seemed confirmed when a host of additional—but larger—specimens were found at Hadar. Johanson, now at the Institute of Human Origins in Berkeley, California, and Tim White of the University of California, Berkeley, concluded that all the material represented a single, sexually dimorphic species: The small individuals like Lucy were females, and the large specimens were males. *A. afarensis*, they further claimed, was the single root giving rise to all subsequent hominids, including our own genus, *Homo*.

Other investigators protested that the disparity in size was too great and implied two separate species, with only the larger one ancestral to *Homo*. The debate sizzled for over a decade, until most experts were persuaded that Johanson and White had been right (*Science*, 1 April 1994, p. 34).

Häusler and Schmid are not persuaded. They compared two different reconstructions of Lucy's pelvis with that of another from the South African site of Sterkfontain, usually attributed to the species *Australopithecus africanus*. They looked at a standard suite of traits used to ascertain sex in modern humans, such as a ridge on the pubic bones

called the ventral arc, found in 95% of modern females pelvises, and the promontorium, a protrusion at the rear of the pelvis that juts forward in males, giving the pelvic inlet a heart shape. The Sterkfontain pelvis appears to be female, while Lucy, with a ridgeless, heart-shaped pelvis, seems to be a male.

Häusler and Schmid concede that modern male features on Lucy's decidedly unmodern pelvis are only partly convincing; they may have had nothing to do with gender in australopithecines. But could such a pelvis give birth to an australopithecine infant? Using standard regression equations, they scaled down estimates for the average size of an adult australopithecine skull to neonatal proportions. They did this calculation two ways: by including the larger specimens from Hadar (which should be done if *A. afarensis* is indeed one species), and by excluding them (as if they belonged to another species). In the first case, birth through Lucy's pelvis was impossible; their calculations indicated that there was simply not enough room to allow the infant's head to pass through. In the second instance, starting with smaller fossils and scaling down to a smaller neonate head, Häusler and Schmid

conclude that birth would have been possible, but with great difficulty. So Lucy could be female—but only if the larger fossils were another species. "I cannot say for certain that Lucy was male," says Häusler. "What I can say is that she did not belong to a species with great sexual dimorphism in body size."

Lovejoy, Johanson, and others strongly disagree. Lovejoy points out that the Swiss study depends on estimates of neonatal head sizes in australopithecines that are themselves based on controversial estimates of adult brain sizes—all to determine whether a hypothetical infant of a vanished species could fit through a pelvis that was itself recovered in a badly crushed condition.

Given these stacked assumptions, says Karen Rosenberg of the University of Delaware, Lucy's small stature may still be the most telling feature. "Lucy is not absolutely the smallest specimen at Hadar in every feature, but she's pretty close," says Rosenberg. "By that standard, it's hard to imagine how she could be male."

—James Shreeve

James Shreeve is a science writer in Takoma Park, Maryland.

PHYSICS


LEPing Up to Higher Energies

The unwritten rule of particle physics has always been build a better accelerator, and you'll discover something new. That expectation has been seriously dampened by the success of the "Standard Model" of particles and forces, which predicts no new particles beyond those already discovered. It has not been quenched entirely, however. Proposed "supersymmetric" theories—efforts to transcend the Standard Model by linking all of the fundamental forces except gravity in a single framework—suggest that an abundance of new particles is waiting to be discovered. And physicists have always had an optimistic streak, in any case.

That optimism is amply evident these days at the European Laboratory for Particle Physics (CERN), where physicists are enjoying a new look at some of the highest energies ever. On 31 October, CERN's Large Electron-Positron Collider (LEP) began a month-long run at 130 billion electron volts (GeV), nearly twice its previous energy and the highest ever reached in an electron-positron machine. With physicists at four detectors analyzing particle collisions online, the result, says LEP physics coordinator Tiziano Camporesi, "is incredible hysteria. ... People are manning shifts, 24 hours a day, on all four experiments."

Since the run began, operators have cranked up the energy another notch, to 140 GeV. That still falls well short of the energy

of the Tevatron Accelerator at the Fermi National Accelerator Laboratory, which collides protons at a trillion electron volts. But because each proton is made of three quarks and many gluons, proton-proton collisions



Minute alchemy. A 131-GeV electron-positron collision at LEP spawns a photon (green) and a massive Z boson, which promptly annihilates in showers of other particles.

are the microscopic equivalent of slamming together bags of marbles. Electrons and positrons, on the other hand, are point particles, which makes LEP "a much cleaner environment" for detecting new particles that might materialize at high energies, says University of Geneva physicist Maurice

Bourquin, a LEP experimentalist.

LEP owes its new potency to 16 new superconducting radio-frequency cavities, which CERN engineers added to the 44 superconducting RF cavities already accelerating particles in the 6-year-old machine. The upgrade, completed last month, is the first in a series that will eventually take LEP to 196 GeV by the summer of 1998. "This run was a so-called pilot run at high energy," says Camporesi, "but it turns out we already have some discovery power in the supersymmetric energy range." Among other things, he says, the beam's luminosity—its density of particles—is unexpectedly high. Now CERN physicists are hoping the run will nail down the lightest predicted supersymmetric particle, the so-called chargino.

For those who want to follow the quest, results from the new LEP run can be seen on the home pages of the various LEP experiments. That of L3, an experiment led by Nobel laureate Sam Ting, can be found at <http://hpl3sn02.cern.ch/130GeV.html>. The page offers images of collisions and their debris. So far, none of the events violates the Standard Model. But the experimenters are undaunted. The page concludes: "The largest excitement from this higher energy run comes from the search of events due to new physics processes." Click on "events," and at press time the display read, "We are very sorry but there are no candidate New Physics events (YET) ..."

—Gary Taubes