It is perhaps not accidental that the view that excellence does not require focused self-determination arises simultaneously with the view that excellent research can be predicted and scheduled, that an effort of 5% on the part of a principal investigator on a project suffices, and that the prolonged adolescence of apprenticeship should continue into the postdoc and perhaps "captive assistant professorship." For committed graduate students wondering whether it was always this way, the answer is "no." For student or professor looking for a little insight into the requirements for excellence, I recommend Gertrude Stein's essay "What are masterpieces and why are there so few of them?'

Douglas E. Brash

School of Medicine, Yale University, New Haven, CT 06520-8040, USA

Resubmission of NIH Grant Proposals

I read with dismay the latest proposal by the National Institutes of Health to streamline the grant process by disallowing more than one resubmission of a grant (ScienceScope, 6 Oct., p. 19). This is ironic since it is admitted that this policy would have a small impact on reducing paperwork. As a current member of several granting agency study sections, I have observed that the review committees frequently do not fund a potentially interesting and important grant submission in part because an element has not been fully expanded or proposed. This points to the larger reason as to why grants are being repetitively submitted-there are insufficient resources to fund many fine grant proposals. In fact, there is often nothing intrinsically wrong with a proposal; rather it is the policy of support for research in this country that "limits enthusiasm." At the risk of sounding descriptive and not proposing any mechanism for change, I believe the researcher's time in preparing resubmissions is every bit as valuable as that of the reviewer and administrative apparatus. What is the alternative to resubmission, assuming that there is a relatively favorable impression of the proposal?

Ellis R. Levin

Department of Medicine, California College of Medicine, University of California, Irvine, CA 92717, USA

Quantal Fluctuations: Correction

We have discovered misstatements in our report (1) "Intrinsic quantal variability due to stochastic properties of receptor-transmitter interactions" (27 Nov. 1992, p. 1494). First, the description of the Monte Carlo model incorrectly states that the diffusion constant used in the simulations was in the range of 0.5 to 1.0×10^{-6} cm² s⁻¹, when it actually was one order of magnitude greater $(0.5 \text{ to } 1 \times 10^{-5})$. Second, the time step was not 1 µs, as stated; the computer program we used automatically adjusted this variable to avoid unacceptably large transition probabilities such as would have occurred with the kinetic parameters used in those simulations. Typically, this resulted in a time step one order of magnitude smaller than what was reported. In addition, the inhibitory postsynaptic potentials illustrated in figure 3, B to D, of the report (p. 1496) were composite responses and not quantal events in the gold-



Fig. 1. Increased variance during the falling phase of quantal inhibitory responses. Intracellular recordings from the M-cell soma of adult goldfish, in the presence of TTX. (A) Superimposed miniature inhibitory postsynaptic potentials (mIPSPs, n =47), demonstrating increased fluctuation in the waveforms during their decay phase. mIPSPs are depolarizing because the recordings were obtained with a KCI containing microelectrode. (B) The mean and standard deviation (σ) waveforms from the same data set. As responses are aligned at onset, σ at that time point is less than that of the background (dashed line), but exceeds it during the decay of the mean mIPSP. (C) TTX blocked medullary impulses, as demonstrated by comparing the antidromic action potential and the subsequent collateral IPSP recorded in the control condition with the same responses obtained at the time the mIPSPs were collected. Noise data were digitized with a 25 kHz sampling rate and filtered at 1 kHz with a second-order Bessel filter.

fish. The quantal responses are quite smaller. Further analyses of quanta isolated in tetrodotoxin (TTX) demonstrates they have a similar relationship between the mean and variance waveforms (Fig. 1). Thus, although the analysis illustrated in figure 3, B to D, is for larger responses, the results remain valid. **Donald S. Faber**

Medical College of Pennsylvania and Hannemann University, Philadelphia, PA 19129, USA **William S. Young** Mellon Institute, Pittsburgh, PA 15213, USA **Pascal Legendre Henri Korn** Institut Pasteur, INSERM U261, Paris, France

References

Authorship Disputes

Jocelyn Kaiser ("Commission proposes new definition of misconduct," News, 29 Sept., p. 1811) says the Commission on Research Integrity may recommend that ORI (the Office of Research Integrity) and institutions "no longer dismiss plagiarism cases involving collaborators as mere authorship disputes." The Commission's intent is better than its logic. There are no "mere" authorship disputes.

Authorship disputes concern stealing. A complaint is made that an author has presented someone else's work as his or her own, an act of intellectual theft. Complaints may be minor—little may have been stolen—or unfounded, but with the term "mere authorship disputes," the illusion is created that no theft is involved. The Commission should not have repeated this canard.

> Charles W. McCutchen 5213 Acacia Avenue, Bethesda, MD 20814, USA

Letters to the Editor

Letters may be submitted by e-mail (at science_letters@aaas.org), fax (202-289-7562), or regular mail (*Science*, 1333 H Street, NW, Washington, DC 20005). Letters are not routinely acknowledged. Full addresses, signatures, and daytime phone numbers should be included. Letters should be brief (300 words or less) and may be edited for reasons of clarity or space. In October 1995, our previous policy of consulting with all letter authors before publication was discontinued.

D. S. Faber, W. S. Young, P. Legendre, H. Korn, Science 258, 1494 (1992).