## Long-Term Effects of Acid Rain: Response and Recovery of a Forest Ecosystem

G. E. Likens,\* C. T. Driscoll, D. C. Buso

Long-term data from the Hubbard Brook Experimental Forest, New Hampshire, suggest that although changes in stream pH have been relatively small, large quantities of calcium and magnesium have been lost from the soil complex and exported by drainage water because of inputs of acid rain and declines in atmospheric deposition of base cations. As a result, the recovery of soil and streamwater chemistry in response to any decreases in acid deposition will be delayed significantly.

**R**esearch on the effects of acid rain (1) in the United States and Europe has focused primarily on the biogeochemistry of sulfur, and to a lesser extent on that of nitrogen (2). This emphasis was because sulfuric acid is the dominant acid in precipitation throughout the eastern United States and Europe where acid rain is a serious environmental problem (3). Records since 1963 show that rain and snow in northeastern United States have had an average annual pH of 4.05 to 4.3 and that sulfuric acid contributed 55 to 75% of this measurable acidity (4). However, this focus on sulfur obscured the role of other elements in the regulation of the long-term, acid-base status of precipitation, soil, and surface waters, as well as linkages between element cycles.

After passage of the Clean Air Act in 1970, emissions of  $SO_2$  and particulates were reduced, and concentrations of sulfate in surface waters in eastern North America declined (Fig. 1, A and B) (5-7). Despite such changes, surface waters throughout the region that are low in acid neutralizing capacity (ANC) (8) have not shown a commensurate recovery in pH and ANC. Additional reductions in emissions and concomitant decreases in the deposition of strong acids are expected as a result of the 1990 amendments to the Clean Air Act. However, critical questions remain: will forest and aquatic ecosystems recover from the effects of acid rain, and if so, at what rate?

The chemistry of bulk precipitation and stream water has been measured continuously at Hubbard Brook Experimental Forest (HBEF) since 1963. These data show that annual, volume-weighted concentrations of  $SO_4^{2-}$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ , and  $Na^+$  declined in both precipitation and stream water since 1963; K<sup>+</sup> declined in precipitation. These declines in atmospheric deposition at HBEF followed declines in atmospheric emissions since 1970 (Fig. 1) (4, 5, 7, 9, 10). On the basis of mass balance calculations, between 77% and 85% of the decline in stream base cations  $C_{\beta}$  (11) can be explained by decreases in the atmospheric deposition of  $C_{\beta}$  (7). The magnitude of decline of  $C_{\beta}$  in stream water was similar to that in bulk precipitation (30 to 40 µeq liter<sup>-1</sup>) after 1963 (Fig. 1D).

Before 1970 (12), concentrations of  $C_{\beta}$  generally were increasing in stream water in





response to high or increasing concentrations of strong acid anions ( $SO_4^{2-}$  +  $NO_3^{-}$ ) (13). Concentrations peaked in 1969 to 1970 (Fig. 1, B and D), coincident with elevated concentrations of  $NO_3^{-}$  in stream water (4), and then declined (Fig. 1B). Significantly, the time path of the relation between  $C_B$  and strong acid anions in stream water shows a hysteresis pattern (Fig. 2).

It is thought that acid rain began around 1950 to 1955 in the northeastern United States (14). Although they are the longest record in North America, the data from HBEF do not include this period. There are data, however, for Ca<sup>2+</sup>, Na<sup>+</sup>, and K<sup>+</sup> in precipitation for 1955–1956 (15). Using the relation of  $Mg^{2+}$  to  $Ca^{2+}$  during 1963 to 1970, we estimated the Mg<sup>2+</sup> concentration for 1955–1956 and thus a  $C_{B}$  concentration of 53.5  $\mu$ eq liter<sup>-1</sup> (Fig. 1D), which is about five times the average value for 1984 to 1993 (Fig. 1D) (16). We also estimated the concentration of  $C_{\beta}$ , strong acid anions, and ANC in stream water in 1940 (Fig. 2) (17).

Calcium is the major cation in stream water at HBEF (4) and is indicative of  $C_{\beta}$ . We apportioned the ecosystem sources and sinks of Ca<sup>2+</sup> (Fig. 3) from 1940 to 1963 using estimated values, and from 1963 to



Fig. 2. Relation between annual, volume-weighted average concentration of the sum of base cations  $(C_{\alpha})$  and the sum of  $SO_{4}^{2-} + NO_{3}^{-}$  for stream water from Watershed 6 of the Hubbard Brook Experimental Forest from 1964-1965 through 1993-1994. The linear regression for data from 1965-1966 through 1969-1970 (O) is Y = 0.83a + 17.4 ( $r^2 = 0.98$ ), and for 1970–1971 through 1993–1994 (**•**) is Y = 0.63a + 29.6 ( $r^2 =$ 0.91), all in microequivalents per liter; the 95% confidence intervals for these two regressions do not overlap. PIR, pre-Industrial Revolution (around 1800). Data for  ${\rm SO_4^{2-}}$  and  ${\rm NO_3^{-}}$  in 1940 and 1963–1964 and  $C_{\rm B}$  in 1940 are estimated [for 1940, see (17);  $C_{B}$  was 143 µeq liter<sup>-1</sup> in 1963, but anions were not measured]. Values for SO42and NO<sub>3</sub><sup>-</sup> were estimated from a 1:1 linear extension of the relation at PIR, assuming an ANC of 20  $\mu$ eq liter<sup>-1</sup>.

G. E. Likens and D. C. Buso, Institute of Ecosystem Studies, Millbrook, NY 12545, USA.

C. T. Driscoll, Department of Civil and Environmental Engineering, Syracuse University, Syracuse, NY 13244, USA.

<sup>\*</sup>To whom correspondence should be addressed.

present using our measurements (18). Concentrations of strong acid anions and Ca<sup>2-</sup> (and  $C_{\beta}$ ) in stream water likely increased after 1940 because of increasing atmospheric inputs of strong acids. Apparently, acid rain directly or indirectly (19) enhanced the release of  $\mathrm{Ca}^{2+}$  (and  $\mathrm{C}_{\mathrm{B}})$  from forest soil from the mid-1950s until the early 1970s (Figs. 1D, 2, and 3). As the more labile pool of  $C_{\beta}$  in soil was depleted, the concentration of  $C_{\beta}$  in stream water decreased (1970 to 1994) (Figs. 1D, 2, and 3). Our initial measurements in 1963 probably captured the end of the first phase of rapidly increasing acidification of soil and drainage water and the marked depletion of  $Ca^{2+}$  (and  $C_{0}$ ) in soil (Figs. 2 and 3).

Increase in streamwater concentration of  $C_{\beta}$  during the 1960s corresponds with high but decreasing inputs of Ca<sup>2+</sup> (and  $C_{\beta}$ ) from atmospheric deposition and large net losses from the soil; declining concentrations of Ca<sup>2+</sup> (and  $C_{\beta}$ ) in stream water thereafter coincide with smaller atmospheric inputs and decreases in net biomass storage and net soil loss (Fig. 3). Other studies suggest that more than 50% of the Ca<sup>2+</sup> supply to surface organic soil exchange and vegetation cation pools of forest soils in the northeastern United States may be derived from atmospheric deposition (20).

We estimate that annual bulk precipitation input of  $Ca^{2+}$  from 1940 to 1955 was seven times greater than that from 1976 to 1993, and average annual loss from soil from 1955 to 1975 was 1.9 times greater than that from 1976 to 1993 (21). Before the mid-1950s, annual  $Ca^{2+}$  depletion from the soil exchange complex was about equal to net biomass storage, and atmospheric deposition plus weathering release nearly



**Fig. 3.** Annual ecosystem fluxes of calcium for Watershed 6 of the Hubbard Brook Experimental Forest from 1940 to 1963 (estimated) and 1964 to 1994 (measured). Fluxes: P is bulk precipitation input, W is weathering release, S is streamwater loss, B is net biomass storage, and D is net release from the soil exchange complex. Data for P during 1955–1956 are from (*15*).

balanced streamwater loss. After the mid-1950s, soil depletion of  $Ca^{2+}$  was greater (1.4 times from 1955 to 1975 and 2 times from 1976 to 1993) than net biomass storage, and streamwater loss was larger (2.2 times from 1955 to 1975 and 2.7 times from 1976 to 1993) than bulk precipitation plus weathering release (Fig. 3). Bulk precipitation provided some 29% of the  $Ca^{2+}$  required in the ecosystem to balance net biomass storage plus streamwater loss before the mid-1950s; however, from 1955 to 1975, bulk precipitation input decreased to meet only 8% of the total demand, while net soil release increased to 77% (Table 1).

Marked declines in the total  $Ca^{2+}$  and  $Mg^{2+}$  concentrations of surface organic soil at HBEF have been measured since 1970 (22), and declines have been calculated or anticipated in soil elsewhere in the northeastern United States (19, 20, 23). Appreciable acidification of forest soils by acid rain during the past 50 to 60 years also has been observed in Sweden (24).

Similarly, the Ca<sup>2+</sup> content of tree rings increased in the 1960s at several locations in the eastern United States. These changes have been attributed to the increased mobilization of  $C_{\beta}$  in surface organic soils by acid rain (25). Stemwood calcium has decreased since the 1960s as the pool of exchangeable  $C_{\beta}$  in the soil has become depleted (Fig. 1E).

Annual forest biomass accumulation at HBEF has declined unexpectedly to a small rate since 1987 (9), perhaps because available  $Ca^{2+}$  became limiting to forest growth as  $C_{\beta}$  was depleted from the soil (23). Pinpointing the cause should become a major area of research. Calcium-to-aluminum ratios for soil waters in high-elevation forest stands at HBEF are in a range that may have adverse effects on the growth or nutrition of forest vegetation (26).

After 1963, the average streamwater pH increased slightly at HBEF (pH 4.85 in 1963 to 5.01 in 1993) (Fig. 1C), but the ANC has been consistently negative (currently about  $-6 \ \mu eq$  liter<sup>-1</sup>). Before the mid-1950s, the ANC in stream water probably was positive. Extrapolating the current

**Table 1.** Percent of ecosystem demand (streamwater loss plus biomass storage) for calcium provided by bulk precipitation (P), weathering release (W), and net release from the soil exchange complex (D).

Period	P (%)	W (%)	D (%)	Total demand (eq ha <sup>-1</sup> )
1940–1955	29	9	63	19,610
1955–1975	8	15	77	25,770
1976–1993	12	8	79	12,300
1940–1993	17	10	73	57,680

rate of decreases in concentrations of strong acid anions and the corresponding changes in  $C_{\beta}$  in stream water at HBEF, we predict that ANC in stream water will become positive again around autumn 1999 (Fig. 2). Nevertheless, even with major reductions in emissions anticipated from the 1990 amendments to the Clean Air Act (27), it is unlikely that the acid-base status of stream water will return to pre–Industrial Revolution levels in the foreseeable future (Fig. 2). Moreover, the large depletion of  $C_{\beta}$  from the soil will retard the recovery of the forest ecosystem in response to any decreases in emissions of SO<sub>2</sub>.

Our analysis suggests that forest ecosystems are much more susceptible to continuing atmospheric inputs of strong acids than expected on the basis of trends in sulfur biogeochemistry alone. The long-term hysteresis pattern in streamwater chemistry suggests that ecosystem acidification may be reversible, but recovery is very slow (Fig. 2). A major policy implication of this analysis is that the 1990 amendments to the Clean Air Act will not be adequate to protect surface waters and forest soils of the northeastern United States against further anthropogenic acidification.

## **REFERENCES AND NOTES**

- The popular term acid rain as used here refers to the atmospheric deposition of all wet and dry substances that cause acidification.
- S. Odén, Acidification of Air and Precipitation and its Consequences on the Natural Environment (Swedish National Science Research Council, Bulletin 1, Stockholm, 1968); J. H. Gibson, Acid Deposition: Long-Term Trends (National Academy Press, Washington, DC, 1986).
- G. E. Likens, R. F. Wright, J. N. Galloway, T. J. Butler, Sci. Am. 241, 43 (October 1979).
- G. E. Likens and F. H. Bormann, *Biogeochemistry of* a Forested Ecosystem (Springer-Verlag, New York, ed. 2, 1995).
- G. E. Likens, *The Ecosystem Approach: Its Use and Abuse* (Ecology Institute, Oldendorf-Luhe, Germany, 1992).
- P. J. Dillon, R. A. Reid, E. de Grosbois, *Nature* **329**, 45 (1987); M. D. Morgan, *J. Environ. Qual.* **19**, 296 (1990); C. T. Driscoll and R. van Dreason, *Water Air Soil Pollut.* **67**, 319 (1993).
- C. T. Driscoll, G. E. Likens, L. O. Hedin, J. S. Eaton, F. H. Bormann, *Environ. Sci. Technol.* 23, 137 (1989).
- ANC is 2[Ca<sup>2+</sup>] + 2[Mg<sup>2+</sup>] + [NA<sup>+</sup>] + [NH<sub>4</sub><sup>+</sup>] + 3[Al<sub>n</sub><sup>+</sup>] - 2[SO<sub>4</sub><sup>2-</sup>] - [NO<sub>3</sub><sup>-</sup>] - [Cl<sup>-</sup>] - [F<sup>-</sup>], where the brackets denote concentration in moles per liter; Al<sub>n</sub> is the nonhydroxide complexes of aluminum.
- 9. G. E. Likens et al., Biogeochemistry 25, 61 (1994).
- T. J. Butler and G. E. Likens, Atmos. Environ. A 25, 305 (1991); L. O. Hedin et al., Nature 367, 351 (1994). Bulk precipitation samplers underestimate the dry portion of total atmospheric deposition, but no temporal dry deposition data are available at HBEF.
- 11. The sum of base cations  $C_{\beta}$  is  $Ca^{2+} + Mg^{2+} + Na^{+} + K^{+}$  in microequivalents per liter.
- On the basis of hydrologic considerations, annual values for HBEF are calculated for a water year, 1 June through 31 May (4).
- Chloride is a small contribution to long-term bulk precipitation (9.0%) and stream water (9.1%) fluxes of strong acid anions at HBEF.
- See, for example, C. V. Cogbill and G. E. Likens, Water Resour. Res. 10, 1133 (1974).

- C. E. Junge and R. T. Werby, *J. Meteorol.* **15**, 417 (1958). Samples were collected from July 1955 through June 1956.
- Estimated Mg<sup>2+</sup> concentration represents 32% of the 1955–1956 C<sub>β</sub> value. If the Mg<sup>2+</sup> concentration were zero, which is unrealistic, the C<sub>β</sub> value would be 36.2 µeq liter<sup>-1</sup>, or about three times the current value.
- 17. Sulfate in pre-Industrial Revolution (PIR) (around 1800) stream water was estimated at 20 µeq liter [T. J. Sullivan et al., Nature 345, 54 (1990); T. J. Sullivan, in Acidic Deposition and Aquatic Ecosystems, D. F. Charles, Ed. (Springer-Verlag, New York, 1991)], and NO<sub>3</sub><sup>-</sup> was estimated at 5  $\mu$ eq liter<sup>-1</sup>, equaling 25  $\mu$ eq liter<sup>-1</sup> for SO<sub>4</sub><sup>2-</sup> + NO<sub>3</sub><sup>-</sup>. PIR values of ANC were probably about 20 µeq liter-1 resulting in a  $C_{\rm B}$  estimate of about 45 µeq liter-Initially, in response to increases in emissions of SO2 and consequent atmospheric deposition of SO42  $C_{\rm B}$  concentrations in drainage water probably increased stoichiometrically with increases in SO<sub>4</sub><sup>2</sup> Empirical relations between SO2 emissions and stream SO<sub>4</sub><sup>2-</sup> at HBEF suggest stream concentra-tions of SO<sub>4</sub><sup>2-</sup> + NO<sub>3</sub><sup>-</sup> of 96  $\mu$ eq liter<sup>-1</sup> in 1940. If  $C_{\rm B}$  increased stoichiometrically with increases in  $SO_4^{-2-}$ ,  $C_\beta$  and ANC would have been 116 and 20  $\mu$ eq liter<sup>-1</sup>, respectively, in 1940.
- 18. The weathering rate of Ca2+ was estimated from Ca:Na (molar ratio) in bedrock, assuming that all streamwater Na+ was derived from rock weathering. Calcium release from weathering before 1963 was assumed to be the same as the average from 1963 to 1993, 105 eq ha-1 year-1. Before 1973, net biomass storage was assumed to be constant (718 eq ha<sup>-1</sup> year<sup>-1</sup>) [on the basis of F. H. Bormann and G. E. Likens, Pattern and Process in a Forested Ecosystem (Springer-Verlag, New York, 1979)], and after 1973, its value came from a logistic curve fitted to data from 1965 to 1992. Although a logistic curve can be fitted to bulk precipitation input from 1956 to 1994 ( $r^2 = 0.87$ ), we used linear trends and projected the 1955-1956 value (350 eq ha-1 year-1) to 1940. Estimated streamwater Ca2+ output was based on a relation between SO<sub>2</sub> emissions and concurrent SO<sub>4</sub><sup>2-</sup> + 5  $\mu$ eq liter<sup>-1</sup> of NO<sub>3</sub><sup>-</sup>. Before 1963, the amount of precipitation and streamflow was taken as the average for 1963 to 1993
- 19. G. B. Lawrence, M. B. David, W. C. Shortle, *Nature* **378**, 162 (1995).
- E. K. Miller, J. D. Blum, A. J. Friedland, *ibid.* 362, 438 (1993). Strontium was measured as a surrogate for calcium.
- 21. The total depletion of Ca<sup>2+</sup> from labile soil pools during 1940 to 1995 was estimated at 42,200 eq ha<sup>-1</sup> (Fig. 3). If Ca<sup>2+</sup> bulk precipitation concentrations were low from 1940 to 1995, such as currently observed (3.6 μeq liter<sup>-1</sup>), depletion of soil Ca<sup>2+</sup> would have been 50,400 eq ha<sup>-1</sup>. In contrast, if Ca<sup>2+</sup> precipitation concentrations were high from 1940 to 1995, such as observed in 1955–1956 (25 μeq liter<sup>-1</sup>), soil depletion of Ca<sup>2+</sup> would have been 34,400 eq ha<sup>-1</sup>. As a result, reductions in atmospheric deposition of Ca<sup>2+</sup> associated with particulate emission control (5) contributed some 20% of the apparent depletion of soil Ca<sup>2+</sup> at HBEF.
- 22. Total Ca<sup>2+</sup> and Mg<sup>2+</sup> concentrations in Oie and Oa soil horizons for Watershed 6 of HBEF declined by 30 to 40% between 1970 and 1987 [R. D. Yanai, T. G. Siccama, M. A. Arthur, *Bull. Ecol. Soc. Am.* **74**, 496 (1993)]. Elsewhere in the region, Ca<sup>2+</sup> in forest floors declined by 22% on average during 1979 to 1994, and Mg<sup>2+</sup>, by 14% [R. D. Yanai, M. A. Arthur, T. G. Siccama, C. A. Federer, *Bull. Ecol. Soc. Am.* **76**, 290 (1995); (19)].
- 23. C. A. Federer *et al.*, *Environ. Manage.* **13**, 593 (1989).
- C. O. Tamm and L. Hällbacken, *Water Air Soil Pollut.* 31, 337 (1986); *Ambio* 17, 56 (1988).
- E. A. Bondietti, N. Monroshima, W. C. Shortle, K. T. Smith, *Can. J. For. Res.* **20**, 1850 (1990); W. C. Shortle and E. A. Bondietti, *Water Air Soil Pollut.* **61**, 253 (1992).
- C. S. Cronan and D. F. Grigal, *J. Environ. Qual.* 24, 209 (1995). Mean Ca:Al molar ratios for spruce, fir, and white birch stands at HBEF (elevation, 750 m) were 1.27 in Oa horizon leachate, decreasing to 0.40

in Bh and 0.27 in Bs horizon solutions. In high-elevation (730 m) deciduous stands, Ca:Al decreased from 1.21 in Oa to 0.59 in Bh and 0.39 in Bs horizon soil water.

- 27. We used current  $NO_3^-$  plus 50% of the 1980  $SO_4^{2-}$  values to estimate the minimum trajectory of strong acid anion concentrations after 1999 in Fig. 2 (5).
- "National Air Pollutant Emission Estimates 1940– 1990," EPA-450/4-91-026 (U.S. Environmental Protection Agency, Research Triangle Park, NC,

November 1991)

29. Financial support was provided by the National Science Foundation and the A. W. Mellon Foundation. We thank the various contributions of M. A. Arthur, C. Eagar, C. E. Johnson, C. W. Martin, T. G. Siccama, and R. D. Yanai. The HBEF is operated by the U.S. Department of Agriculture Forest Service, Radnor, Pennsylvania.

13 November 1995; accepted 18 January 1996

## History of Ancient Copper Smelting Pollution During Roman and Medieval Times Recorded in Greenland Ice

Sungmin Hong, Jean-Pierre Candelone, Clair C. Patterson,\* Claude F. Boutron†

Determination of copper concentrations in Greenland ice dated from seven millennia ago to the present showed values exceeding natural levels, beginning about 2500 years ago. This early large-scale pollution of the atmosphere of the Northern Hemisphere is attributed to emissions from the crude, highly polluting smelting technologies used for copper production during Roman and medieval times, especially in Europe and China. This study opens the way to a quantitative assessment of the history of early metal production, which was instrumental in the development of human cultures during ancient eras.

**M**ost emissions of pollutants to the global atmosphere of our planet occurred after the Industrial Revolution, driven by the combination of unprecedented growth in the population of the Earth and massive technological and economic development. The only exception reported so far is lead emission. Time series recently obtained from Greenland ice (1) and Swedish lake sediments (2) indicate that there was early hemispheric-scale atmospheric pollution from lead during Greek and Roman times. The importance of this early lead pollution remained relatively limited, however; cumulative lead deposition to the Greenland ice cap during the eight centuries of the flourishing of the Greek and Roman civilizations ( $\sim$ 400 metric tons) was about an order of magnitude lower than lead deposi-

C. C. Patterson, Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California 91125, USA.

C. F. Boutron, Laboratoire de Glaciologie et Géophysique de l'Environnement du CNRS, 54 rue Molière, Domaine Universitaire, Boite Postale 96, 38402 Saint Martin d'Hères, France, and Unité de Farmation et de Recherche de Mécanique, Université Joseph Fourier de Grenoble (Institut Universitaire de France), Domaine Universitaire, Boite Postale 68, 38041 Grenoble, France.

\*This paper is dedicated to the memory of C. C. Patterson. †To whom correspondence should be addressed. tion from post–Industrial Revolution fallout [~3200 metric tons (3)]. Here we report evidence that cumulative early atmospheric pollution from copper before the Industrial Revolution was one order of magnitude greater than that from the Industrial Revolution to the present.

We analyzed copper in 23 sections of the 3028.8-m Greenland Ice Core Project (GRIP) deep ice core (4), drilled in central Greenland at Summit (72°34′N, 37°37′W; elevation 3238 m above sea level). Twenty-one sections were selected from ice dated from 2960 to 470 years ago (from depths of 619.3 to 129.3 m) in order to cover the Greek and Roman civilizations, barbaric and medieval times, and the Renaissance. Two sections were chosen from ice dated from 7260 and 7760 years ago (depths of 1230.4 and 1286.5 m, respectively) to assess natural Holocene copper concentrations during times before copper production. We mechanically decontaminated each core section using ultraclean procedures by chiseling successive veneers of ice from the outside to the central part as described in detail elsewhere (5). Copper concentrations were then determined with a precision of  $\pm 10\%$  by graphite furnace atomic absorption spectrometry (GFAAS) after preconcentration. Each value was blank-corrected (5). The accuracy of the data depends on the efficiency of the decontamination procedure. This was systematically checked for all sections by investigation of radial changes in copper concentrations from the outside to the center of each individual core section (6). Copper concentrations measured in the central parts of the

S. Hong, Laboratoire de Glaciologie et Géophysique de l'Environnement du CNRS, 54 rue Molière, Domaine Universitaire, Boite Postale 96, 38402 Saint Martin d'Hères, France

J.-P. Candelone, Department of Applied Physics, Curtin University of Technology, General Post Office Box U 1987, Perth, WA 6001, Australia, and Laboratoire de Glaciologie et Géophysique de l'Environnement du CNRS, 54 rue Molière, Domaine Universitaire, Boite Postale 96, 38402 Saint Martin d'Hères, France.