

15. Overflow of Owens Lake occurs when wetness exceeds ~2.4 times the historical mean [H. S. Gale, *U.S. Geol. Surv. Bull.* **580-L**, 251 (1914)].
16. Discontinuities in the depth-age distribution of OL90-2 indicate a sediment hiatus between ~15,500 and 13,700  $^{14}\text{C}$  years B.P. Sediments at the base of the hiatus are characterized by features that indicate desiccation, including a 1- to 3-mm-thick lag deposit of frosted quartz grains. Removal of sediments by wave reworking or deflation implies that the 15,500-year age represents a maximum estimate of the initiation of desiccation.
17. S. P. Lund, unpublished data.
18. The presence of greigite and the absence of magnetite in high- $\chi$  intervals were inferred from thermomagnetic measurements of magnetic mineral blocking temperatures.
19. F. M. Phillips *et al.*, *Science* **274**, 749 (1996).
20. R. I. Dorn *et al.*, *Quat. Res.* **28**, 38 (1987); M. I. Bursik and A. R. Gillespie, *ibid.* **39**, 24 (1993); R. J. Poreda *et al.*, *Eos* **76** (fall suppl.), 685 (1995).
21. F. M. Phillips *et al.*, *Science* **248**, 1529 (1990).
22. E. Blackwelder, *Geol. Soc. Am. Bull.* **42**, 865 (1931).
23. Values of  $\delta^{18}\text{O}$  from 40,000 to 30,000 years B.P. are generally low, indicating a period in which precipitation fell as snow. The lack of pronounced glaciation during this interval is probably the result of relatively high warm-season (May through August) insolation [A. Berger and M. F. Loutre, *Quat. Sci. Rev.* **10**, 297 (1991)].
24. The introduction of glacially derived clay increased lake turbidity, decreasing photosynthetic production of organic carbon. Seasonal ice cover and decreased water temperatures also decreased productivity.
25. Radiocarbon age control for OL90-2 and V23-81 is no better than 500 to 1000 years. Thus, the time between maxima and minima in both records is of the same order as the composite  $^{14}\text{C}$  error, implying that North Atlantic lithic events cannot be uniquely matched with advances or retreats of Sierran glaciers.
26. The lack of correlation between high-frequency events in the lithic and  $\delta^{18}\text{O}$  records may derive from the fact that North Atlantic climate variability mainly depended on changes in air temperature (Dansgaard-Oeschger cycles), whereas the hydrologic balance of Owens Lake depended not only on air temperature but also on cloud cover, humidity, and precipitation [L. V. Benson, *U.S. Geol. Surv. Water Resour. Invest. Rep.* **86-4148** (1986)]. The poor correlation may also result from the timing of carbonate precipitation. Precipitation is favored by warm water temperature and high concentrations of  $\text{Ca}^{2+}$  and  $\text{CO}_3^{2-}$ . These conditions occur in the late autumn when overflow is at a minimum and  $\delta^{18}\text{O}$  values are high and not necessarily representative of average overflow conditions.
27. There are about 19 glacial cycles between 52,500 and 23,500 years B.P.; however, some of the older Tahoe advances (for example,  $A_2$ ) may consist of more than one glacial cycle.
28. We thank J. Andrews, G. Bond, and S. Lehman for their excellent reviews and suggestions. W. C. McClung of Lake Minerals Corporation allowed us access to the coring sites. Support was provided by the U.S. Geological Survey Global Change Program, NSF, and a University of Southern California Faculty Research Innovation Fund grant. The work was performed in part under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract W-7405-ENG-48.

25 April 1996; accepted 12 August 1996

## Chronology for Fluctuations in Late Pleistocene Sierra Nevada Glaciers and Lakes

Fred M. Phillips, Marek G. Zreda,\* Larry V. Benson, Mitchell A. Plummer, David Elmore, Pankaj Sharma

Mountain glaciers, because of their small size, are usually close to equilibrium with the local climate and thus should provide a test of whether temperature oscillations in Greenland late in the last glacial period are part of global-scale climate variability or are restricted to the North Atlantic region. Correlation of cosmogenic chlorine-36 dates on Sierra Nevada moraines with a continuous radiocarbon-dated sediment record from nearby Owens Lake shows that Sierra Nevada glacial advances were associated with Heinrich events 5, 3, 2, and 1.

During the last glacial period, the climate in the North Atlantic region was characterized by a sequence of quasi-cyclical fluctuations (1). Combined ice core and marine sediment core evidence indicates that during periods ranging in duration from about 500 to 2000 years the climate became progressively colder. The maxima of these Dansgaard-Oeschger cycles were often marked by the expulsion of large numbers of icebergs from the ice caps surrounding the North Atlantic (Heinrich events) (2). The iceberg expulsions were rapidly followed by abrupt warming. The cold episodes culminating in Heinrich events have been postulated to be the cause of mountain glacier advances in western North America (3) and elsewhere (4).

F. M. Phillips, M. G. Zreda, M. A. Plummer, Department of Earth and Environmental Science, New Mexico Institute of Mining and Technology, Socorro, NM 87801, USA.  
L. V. Benson, U.S. Geological Survey, 3215 Marine Street, Boulder, CO 80303, USA.  
D. Elmore and P. Sharma, Physics Department, Purdue University, West Lafayette, IN 47907, USA.

\*Present address: Department of Hydrology and Water Resources, University of Arizona, Tucson, AZ 85721, USA.


This hypothesis has proved difficult to test, in large part because of the difficulties in dating moraines by  $^{14}\text{C}$  and other conventional approaches. Cosmogenic nuclide methods (5) can be used to directly date moraines (6, 7), but various uncertainties (8, 9) render tenuous direct chronological comparisons with millennial-scale events such as iceberg discharges.

An alternative approach that circumvents these difficulties is to investigate continuous and datable sedimentary records in environments associated with mountain glaciers. Although the sediment-based approach provides a nearly continuous record, it must use indirect proxies for glacial extent. Here we test glacial proxies in a sediment record from Owens Lake, California (10), by comparing the  $^{14}\text{C}$  chronology of the proxies with direct  $^{36}\text{Cl}$  ages of Sierra Nevada moraines.

The Owens River drains the eastern flank of the Sierra Nevada (Fig. 1). All of the major valleys originating from the Sierra Nevada contain late Pleistocene moraine complexes showing that the altitude of the equilibrium line was ~1000 m lower

(11). The characteristics of sedimentation in Owens Lake should therefore have been sensitive to changes in the magnitude of discharge and type of sediment load produced by glaciation, particularly the release of large amounts of rock flour by glacial meltwater. Benson *et al.* (10) used increases in magnetic susceptibility and decreases in inorganic carbon, organic carbon, and carbonate  $\delta^{18}\text{O}$  as indicators of glacial advance.

We have used cosmogenic  $^{36}\text{Cl}$  buildup (12) to date late Pleistocene moraines in four drainages (Fig. 1). Two of the drainages, Bishop Creek and Little McGee Creek, are tributary to the Owens River. Bloody Canyon drains into Mono Lake and is about 20 km north of the headwa-



**Fig. 1.** Location of Owens River drainage basin and valleys where moraines were dated with the use of cosmogenic  $^{36}\text{Cl}$ . CH = Chiatovich Creek, BC = Bloody Canyon, LMC = Little McGee Creek, and BpCr = Bishop Creek.

ters of the Owens River. Chiatovich Creek is on the eastern side of the White Mountains, the range immediately east of the Sierra Nevada. All four drainages contain complex sequences of late Pleistocene moraines (Fig. 2).

The largest boulders available on the crest of each moraine (Fig. 2) were sampled by hammer and chisel close to the center of their top surfaces. The samples were processed and analyzed for  $^{36}\text{Cl}$  by standard methods (13). Ages were calculated for rock-surface erosion rates of 0 and 5 mm/ka (Fig. 2 and Table 1) (14).

If all of the boulders sampled on a single moraine had the same history, except for variable rock erosion rates, then all samples should show mutual overlap within some part of the age range indicated by the erosion calculations. Instead, boulders from a single moraine usually show scatter toward young ages because soil erosion progressively exposes boulders after moraine deposition (8). We have therefore interpreted the overlapping range of ages that cluster at the maximum end of the range for each moraine as the depositional age (15).


The data (Fig. 2) show that the younger

moraines in the Sierra Nevada yield  $^{36}\text{Cl}$  ages for major advances at  $49 \pm 2$ ,  $31 \pm 1$ ,  $25 \pm 1$ ,  $19 \pm 1$ , and  $16 \pm 1$  ka. Comparison of this chronology, after conversion to the radiocarbon time scale (16), with the magnetic susceptibility and organic carbon content curves from Owens Lake demonstrates that the proxy indicators for glaciation accurately record glacial advances (17). Furthermore, the advances for which moraines are preserved are those having the strongest signals in the sedimentary record. The Tioga 2 advance at  $22 \pm 1$   $^{14}\text{C}$  kyr has the lowest equilibrium line altitude (ELA) of any Tioga advance and correlates with the maximum magnetic susceptibility ( $\chi$ ) signal in Owens Lake. The advance having the next lowest ELA, Tioga 3, corresponds to the second largest peak in  $\chi$ . The moraine position shows that the Tioga 1 advance was relatively minor and was probably preserved at Little McGee Creek only because of faulting. The core evidence from Owens Lake is consistent with a relatively minor advance. This correspondence supports the inference from the Owens core data that there were a series of minor glacial advances between 40 and 25  $^{14}\text{C}$  kyr that are not

recorded by moraines, presumably because the moraines were overridden by the later, more extensive advances.

These glacial events in the Sierra Nevada appear to be temporally related to climatic cycles in the North Atlantic and elsewhere. The earliest advance, at  $49 \pm 2$  ka, was close to the time of Heinrich event 5 (H5) [43 ka on the GRIP time scale (18) and about 47 ka on the GISP2 scale (19)], although uncertainties in both  $^{36}\text{Cl}$  and ice core chronologies at this age range do not permit any close correlations. No moraines correlative to H4 were dated. The Owens Lake core (Fig. 2) indicates that this was an extended interval of limited glacial extent in the Sierra Nevada, and any moraines associated with H4 were probably destroyed by subsequent, more extensive advances. The Tioga 1 advance at  $28 \pm 1.5$   $^{14}\text{C}$  kyr appears to have slightly preceded H3, and the Tioga 2 advance seems to have slightly preceded H2, although the  $^{14}\text{C}$  chronologies of neither the marine nor the Owens Lake cores are sufficiently precise to permit unambiguous distinction of phase relations. The Tioga 2 advance appears to have been a critical transition point in the climate system. After this advance, the climate did not return to the previous oscillatory pattern but rather locked into a full glacial mode. This pattern of rapid increase in glacial extent beginning at about 23  $^{14}\text{C}$  kyr strongly resembles that inferred for global ice volume from Huon Peninsula terraces (20) and  $^{18}\text{O}$  in tropical ocean cores (21). The Tioga 3 advance clearly immediately preceded H1, and the Tioga 4 advance at  $\sim 14.5$  to  $13$   $^{14}\text{C}$  kyr probably included at least the latter half of H1. The extent of both lakes and glaciers was apparently large, but not maximum, during the time preceding H1. At close to the initiation of H1 (15 to 14.5  $^{14}\text{C}$  kyr), the glaciers retreated suddenly, and Owens Lake shrank to near desiccation [less than half of the modern (22) surface area], but by the end of H1, the glaciers had readvanced and the Owens River lacustrine system rapidly refilled to peak late Pleistocene levels (23) (more than five times the modern surface area). The peak in both glacial and lacustrine extent was apparently brief.

The above summary indicates that although expansions of Sierra Nevada glaciers were apparently related to Heinrich events, the relation was probably not one of simple synchronicity. Rather, Sierra Nevada glacial advances, in the case of H1 at least and possibly the other events, slightly preceded the Heinrich events, and during the events, the water balance may have oscillated strongly. The Owens Lake record (10) may also indicate a similar pattern for the smaller iceberg discharge events (1) between H5 and



**Fig. 2.** Chlorine-36 age distributions compared with proxies for glacial extent from Owens Lake sediments from (9). Peaks in the proxy records are interpreted as indicating extensive glaciation. Moraine maps and corresponding  $^{36}\text{Cl}$  ages are color coded. Bloody Canyon (BC) map modified from (6), Bishop Creek (BpCr) from (26), and Chiatovich (CH) from (27). Dots indicate  $^{36}\text{Cl}$  ages assuming zero rock erosion rate, and bars indicate the ages assuming a 5-mm/ka erosion rate.

**Table 1.** Chlorine-36 ages for boulders on Sierra Nevada moraines, for assumed rock erosion rates varying from 0 to 5 mm/ka.

| Sample                           | <sup>36</sup> Cl age (ka) |         | Sample                        | <sup>36</sup> Cl age (ka) |         |
|----------------------------------|---------------------------|---------|-------------------------------|---------------------------|---------|
|                                  | 0 mm/ka                   | 5 mm/ka |                               | 0 mm/ka                   | 5 mm/ka |
| <i>Bishop Creek: Tioga 3</i>     |                           |         | <i>Little McGee: Tioga 4</i>  |                           |         |
| 91-11                            | 20.3                      | 19.8    | LMC88-6                       | 16.7                      | 16.8    |
| 90-22                            | 18.1                      | 19.1    | LMC88-7                       | 16.7                      | 16.9    |
| 90-23                            | 10.4                      | 10.5    | LMC86-1                       | 13.0                      | 12.1    |
| 90-24                            | 17.7                      | 16.0    | LMC86-2                       | 15.8                      | 15.3    |
| 90-25                            | 18.1                      | 18.9    | LMC86-3                       | 15.6                      | 15.0    |
| 90-26                            | 18.8                      | 18.9    | LMC86-4                       | 12.9                      | 13.0    |
| 91-1                             | 17.2                      | 16.4    | LMC86-5                       | 16.9                      | 16.6    |
| 91-2                             | 6.9                       | 6.4     | <i>Little McGee: Tioga 3</i>  |                           |         |
| 91-3                             | 18.6                      | 17.5    | LMC90-12                      | 11.9                      | 11.0    |
| 91-4                             | 17.6                      | 16.6    | LMC90-13                      | 12.6                      | 12.9    |
| 90-73                            | 17.1                      | 16.2    | LMC90-14                      | 14.1                      | 12.8    |
| 90-74                            | 17.9                      | 16.2    | LMC90-15                      | 10.4                      | 10.5    |
| 90-75                            | 19.3                      | 18.4    | LMC90-16                      | 15.0                      | 15.3    |
| 91-5                             | 14.9                      | 14.2    | LMC90-17                      | 12.4                      | 11.0    |
| 91-6                             | 19.8                      | 18.2    | LMC90-19                      | 15.0                      | 13.8    |
| 91-7                             | 29.7                      | 24.6    | LMC90-21                      | 39.3                      | 44.3    |
| 91-8                             | 22.4                      | 24.3    | LMC90-23                      | 11.7                      | 12.1    |
| 90-19                            | 19.7                      | 20.6    | LMC90-24                      | 15.0                      | 15.6    |
| 90-20                            | 18.8                      | 19.6    | LMC90-25                      | 16.5                      | 17.1    |
| 90-21                            | 15.9                      | 16.6    | LMC90-26                      | 18.1                      | 19.3    |
| <i>Chiatovich Creek: Tioga 4</i> |                           |         | LMC90-27                      | 18.9                      | 20.1    |
| C87-1                            | 14.1                      | 12.5    | <i>Little McGee: Tioga 2</i>  |                           |         |
| C87-2                            | 13.3                      | 12.2    | LMC91-3                       | 15.1                      | 15.6    |
| C87-3                            | 6.7                       | 6.4     | LMC91-4                       | 20.0                      | 21.2    |
| C87-4                            | 6.0                       | 5.6     | LMC91-5c                      | 23.1                      | 24.7    |
| C287-5                           | 11.6                      | 11.0    | LMC91-6                       | 13.5                      | 11.9    |
| C287-6                           | 14.3                      | 12.2    | LMC91-7                       | 24.1                      | 25.9    |
| C387-1                           | 14.3                      | 13.1    | LMC91-8                       | 16.4                      | 14.3    |
| C387-2                           | 12.8                      | 12.1    | <i>Little McGee: Tioga 1</i>  |                           |         |
| C387-3                           | 16.6                      | 14.9    | LMC86-6                       | 30.6                      | 31.4    |
| <i>Chiatovich Creek: Tioga 3</i> |                           |         | LMC86-7                       | 3.2                       | 3.2     |
| C587-1                           | 18.8                      | 16.3    | LMC86-8                       | 32.0                      | 30.3    |
| C587-2                           | 20.1                      | 17.1    | LMC86-9                       | 20.2                      | 20.6    |
| C587-3                           | 16.2                      | 12.7    | LMC86-10                      | 31.3                      | 31.8    |
| C787-1                           | 12.9                      | 11.7    | <i>Bloody Canyon: Tioga 4</i> |                           |         |
| C787-2                           | 20.1                      | 17.3    | BC86-1                        | 17.2                      | 17.6    |
| C787-3                           | 4.0                       | 3.9     | BC86-2                        | 10.5                      | 9.9     |
| C7b87-1                          | 20.6                      | 18.5    | BC87-3                        | 16.6                      | 15.4    |
| CH89-P12                         | 15.2                      | 14.8    | BC86-5                        | 18.1                      | 16.2    |
| CH89-P12-1                       | 12.9                      | 12.8    | BC90-5                        | 13.3                      | 13.6    |
| CH89-P22-1                       | 2.3                       | 2.2     | <i>Bloody Canyon: Tioga 3</i> |                           |         |
| CH89-P22-2                       | 5.7                       | 5.6     | BC86-6                        | 19.3                      | 19.2    |
| CH89-P22-3                       | 10.4                      | 10.0    | BC86-7                        | 4.3                       | 4.2     |
| CH89-P23-1                       | 19.3                      | 17.9    | BC86-8                        | 18.9                      | 17.6    |
| CH89-P23-2                       | 22.5                      | 20.3    | BC86-9                        | 19.8                      | 18.8    |
| CH89-P23-3                       | 12.8                      | 12.4    | BC86-10                       | 5.6                       | 5.6     |
| C1187-1                          | 19.3                      | 16.9    | BC86-11                       | 18.9                      | 19.4    |
| C1187-2                          | 12.9                      | 11.3    | BC90-1                        | 16.6                      | 16.5    |
| C1187-3AW                        | 16.1                      | 14.0    | BC90-2                        | 19.4                      | 18.6    |
| C1287-1AW                        | 15.6                      | 14.2    | BC90-3                        | 17.7                      | 17.5    |
| C1287-2                          | 9.0                       | 8.2     | <i>Younger Tahoe</i>          |                           |         |
| C1287-3                          | 6.2                       | 6.0     | BC87-1                        | 43.1                      | 40.8    |
| <i>Chiatovich Creek: Tioga 2</i> |                           |         | BC87-2                        | 44.7                      | 49.0    |
| CH89-P8-1                        | 25.1                      | 25.1    | BC87-3                        | 32.2                      | 30.3    |
| CH89-P8-2                        | 22.0                      | 21.3    | BC87-4                        | 48.9                      | 43.3    |
| CH89-P8-3                        | 22.0                      | 20.4    | BC87-5                        | 53.8                      | 49.5    |
|                                  |                           |         | BC88-1                        | 4.1                       | 4.0     |
|                                  |                           |         | BC88-5                        | 50.5                      | 49.0    |

H3. Although the nature of the climatic linkages remains to be explained, the close association of both glacial and lacustrine fluctuations in western North America with these events supports the hypothesis that they were of global scale.

## REFERENCES AND NOTES

- G. Bond *et al.*, *Nature* **365**, 143 (1993).
- G. Bond *et al.*, *ibid.* **360**, 245 (1992); W. S. Broecker, *ibid.* **372**, 421 (1994).
- P. U. Clark and P. J. Bartlein, *Geology* **23**, 483 (1995).
- T. V. Lowell *et al.*, *Science* **269**, 1541 (1995).

- T. E. Cerling and H. Craig, *Annu. Rev. Earth Planet. Sci.* **22**, 273 (1994).
- F. M. Phillips *et al.*, *Science* **248**, 1529 (1990).
- J. C. Gosse, E. B. Evenson, J. Klein, B. Lawn, R. Middleton, *Geology* **23**, 877 (1995).
- P. R. Bierman, *J. Geophys. Res.* **99**, 13885 (1994).
- A. B. Gibbons, J. D. Megeath, K. L. Pierce, *Geology* **12**, 327 (1984).
- L. V. Benson *et al.*, *Science* **274**, 746 (1996).
- D. W. Burbank, *Quat. Res.* **36**, 294 (1991).
- F. M. Phillips, B. D. Leavy, N. O. Jannik, D. Elmore, P. W. Kubik, *Science* **231**, 41 (1986).
- The samples were leached in deionized water to remove meteoric <sup>36</sup>Cl and were dissolved in hydrofluoric acid; the Cl precipitated as AgCl. Detailed methods and sample location maps are given in (24). The rocks were also analyzed by standard methods for major elements: Cl, Gd, and B, and in some cases U and Th. Complete data are given in (24). Data for the Bloody Canyon moraines were published previously (6). Chlorine-36 was analyzed by accelerator mass spectrometry [D. Elmore *et al.*, *Nature* **277**, 22 (1979)] at the Purdue Rare Isotope Measurement Laboratory, Purdue University. Exposure ages were calculated with <sup>36</sup>Cl production constants determined by F. M. Phillips, M. G. Zreda, M. R. Flinsch, D. Elmore, and P. Sharma [*Geophys. Res. Lett.* **23**, 949 (1996)]. Production rates were assumed to be constant with time. Production by thermal neutrons was calculated using the approach of B. Liu, F. M. Phillips, J. T. Fabryka-Martin, M. M. Fowler, and W. D. Stone [*Water Resour. Res.* **30**, 3115 (1994)], and epithermal neutron production was also incorporated with an analogous approach. Background <sup>36</sup>Cl production by neutrons from U and Th in the rock was calculated using the approach of J. T. Fabryka-Martin [thesis, University of Arizona (1988)].
- Here we use the convention of "ka" to refer to ages on the <sup>36</sup>Cl and ice core time scales (1 ka = 1000 years) and "<sup>14</sup>C kyr" to refer to the <sup>14</sup>C time scale (1 <sup>14</sup>C kyr = 1000 carbon-14 years).
- Modeling [M. G. Zreda, F. M. Phillips, D. Elmore, *Water Resour. Res.* **30**, 3127 (1994)] has shown that, in the age range considered here, effects of thermal neutron production will not cause significant overestimation of maximum ages. Hallet and Putkonen (25) have argued that moraine and rock-surface erosion may produce anomalously young moraine ages. Their analysis is not applicable for the samples reported here because (i) they did not account for production of <sup>36</sup>Cl by thermal neutrons, which counters age bias due to rock weathering; (ii) the average size of boulders we sampled (1 to 4 m) was much larger than they assumed (0.5 m); and (iii) even if their analysis were correct, the effects would be small in the age range considered here [figure 2B of (25)].
- E. Bard, B. Hamelin, R. G. Fairbanks, A. Zindler, *Nature* **345**, 405 (1990).
- These ages also agree well with cosmogenic <sup>3</sup>He ages of 27 and 20.5 ka obtained at June Lake [R. J. Poreda *et al.*, *Eos* **76** (fall suppl.), 685 (1995)].
- W. Dansgaard *et al.*, *Nature* **364**, 218 (1993).
- P. A. Mayewski *et al.*, *Science* **263**, 1747 (1994).
- J. Chappell and N. J. Shackleton, *Nature* **324**, 137 (1986).
- B. K. Linsley, *ibid.* **380**, 234 (1996).
- By "modern" surface area of Owens Lake we mean before the major diversion of Owens River in 1913.
- F. M. Phillips *et al.*, *New Mexico Water Resour. Res. Inst.* **269** (1992); L. V. Benson *et al.*, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **78**, 241 (1990).
- M. G. Zreda, thesis, New Mexico Institute of Mining and Technology (1994).
- B. Hallet and J. Putkonen, *Science* **265**, 937 (1994).
- P. C. Bateman, *U.S. Geol. Surv. Prof. Pap.* **470** (1965).
- T. Swanson, D. L. Elliott-Fisk, R. L. Southard, *Quat. Res.* **39**, 186 (1993).
- We thank R. Dorn, D. Elliott-Fisk, S. S. Smith, and M. Flinsch. Support was provided by the Geography, Geoscience, and Physics divisions of NSF.

29 April 1996; accepted 12 August 1996