that morphological evolution sometimes occurs episodically."

Jerry A. Coyne Brian Charlesworth Department of Ecology and Evolution, University of Chicago, 1101 East 57 Street, Chicago, IL 60637, USA

References

- N. Eldredge and S. J. Gould, in *Models in Paleobiology*, T. J. M. Schopf, Ed. (Freeman, Cooper, San Francisco, 1973), pp. 82–115.
- 2. J. A. Coyne and B. Charlesworth, *Science* **274**, 1748 (1996).
- S. J. Gould and N. Eldredge, *Paleobiology* 3, 115 (1977).
- 4. S. J. Gould, ibid. 6, 119 (1980).
- R. Lande, *ibid.*, p. 233; B. Charlesworth, M. Slatkin, R. Lande, *Evolution* **36**, 474 (1982); J. S. Levinton, *Genetics, Paleontology and Macroevolution* (Cambridge Univ. Press, Cambridge, 1988); B. Charlesworth in *Paleobiology: A Synthesis*, D. E. G. Briggs and P. R. Crowther, Eds. (Blackwell, Oxford, 1990), pp. 100–106.
- 6. S. J. Gould, Science 216, 380 (1982).
- 7. N. Eldredge, *Macroevolutionary Dynamics* (McGraw-Hill, New York, 1989).
- 8. J. Maynard Smith, Ann. Rev. Genet. 17, 11 (1983).
- 9. S. Wright, Proc. 6th Int. Congr. Genet. 1, 356 (1932).
- R. A. Fisher, Ann. Eugen. 7, 355 (1937).
 _____, The Genetical Theory of Natural Selection
- (Oxford Univ. Press, Oxford, 1930).
- 12. K. E. Weber, Genetics 144, 205 (1996).
- 13. S. J. Gould and N. Eldredge, *Nature* **366**, 226 (1993).
- 14. S. J. Gould, New York Times, 25 June 1996, p. B5.

Molecular Chirality Control and Amplification by CPL: Correction

In our report "Dynamic control and amplification of molecular chirality by circular polarized light [CPL]" (20 Sept., p. 1686) (1), a sentence in column 2 on page 1687 could lead to misinterpretation. The sentence reads, "A large pitch ($p = 580 \ \mu m$ based on helical twisting power 0.1) is seen in the cholesteric phase as a result of low resolution of 1 by CPL irradiation." This might be read as if a pitch of 580 micrometers had actually been determined.

Such a large pitch cannot be measured directly, so the following procedure was used. The cholesteric phase with a large pitch was based on the observed texture and control experiments where an increase in pitch of the cholesteric texture with decreasing enantiomeric excess of the dopant was seen. We then calculated what the pitch of the cholesteric phase must be after irradiation by CPL on the basis of the helical twisting power and the enantiomeric excess determined by circular dichroism. We are grateful to G. B. Schuster of the Georgia Institute of Technology for bringing this matter to our attention.

Nina P. M. Huck Wolter F. Jager Ben de Lange Ben L. Feringa Department of Organic and Molecular Inorganic Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, Netherlands


LETTERS

References

1. N. P. M. Huck, W. F. Jager, B. de Lange, B. L. Feringa, *Science* 273, 1686 (1996).


Corrections and Clarifications

- In the 14 March News & Comment article "Court invalidates expert panel report" by Jocelyn Kaiser (p. 1560), the judge's name should have been Paul L. Friedman and the date of the preliminary injunction 5 March.
- In figure 3, A and D, on page 383 of the report "Receptor and $\beta\gamma$ binding sites in the α subunit of the retinal G protein transducin" by R. Onrust *et al.* (17 Jan., p. 381), the left edge was inadvertently cropped. The correct figure 3A appears below. In figure 3D, the label "GDP" should have appeared at the top left end of the line leading to guanosine diphosphate, which is highlighted in pale yellow.



Letters to the Editor

Letters may be submitted by e-mail (at science_letters@aaas.org), fax (202-789-4669), or regular mail (*Science*, 1200 New York Avenue, NW, Washington, DC 20005, USA). Letters are not routinely acknowledged. Full addresses, signatures, and daytime phone numbers should be included. Letters should be brief (300 words or less) and may be edited for reasons of clarity or space. They may appear in print and/or on the World Wide Web. Letter writers are not consulted before publication.



THE NEW, MINI-PREP 24

- High Purity— sufficient for both automated fluorescent and manual sequencing
- Easy Operation—begin prep with direct loading of crude bacteria culture; *no* centrifugation step
- Fast-up to 24 Mini-Preps per hr.
- Consistent Results—up to 5 μg of plasmid per ml.

Call 1-800-466-7949 now to learn how the new, improved Mini-Prep 24 can automate your plasmid DNA prep. Case closed.



11339 Sorrento Valley Rd. San Diego, CA 92121 (619) 452.2603

1-800-466-7949 Circle No. 2 on Readers' Service Card