

sate, is currently in clinical trials in the United States, and several pharmaceutical companies are actively developing other medications to be placed in clinical trials for addictive disorders. There are still many hurdles to overcome, but there is progress.

Charles P. O'Brien

*University of Pennsylvania/
Veterans Affairs Medical Center,
Philadelphia, PA 19104-6178, USA*

References

1. C. E. Fulco, *The Development of Medications for the Treatment of Opiate and Cocaine Addictions* (National Academy Press, Washington, DC, 1995).



Ancient Sharks and Rays

The item "Shark and ray extinctions" by Ann Simon Moffat (Research News, 31 Oct., p. 801) featured a discussion on a portion of our work on Late Cretaceous (Maastrichtian) vertebrates found in the Fox Hills Formation and Paleocene Cannonball Formation of North Dakota, as reported at the recent Society of Vertebrate Paleontology meeting (1). Moffat discussed the Cretaceous-Tertiary (K-T) boundary extinctions as demonstrated by cartilaginous fish faunas of the Western Interior Seaway, and we would like to clarify some aspects of our research that she mentioned.

We have studied more than 40 sites in the Fox Hills Formation and 70 sites in the Cannonball Formation. There is little disagreement that the 22 species of sharks and rays we found in the Fox Hills were extinct by the Paleocene; rather, the issue may be the rapidity of that extinction. As suggested by the comments attributed to J. David Archibald, the K-T boundary section is not complete because units of terrestrial Hell Creek and Ludlow Formations intervene in North Dakota, where we are working. The K-T boundary occurs within this interval of regression (2), making precise correlation between the marine cartilaginous fish faunas and the terrestrial vertebrate faunas difficult.

These facies, although complex, are well understood paleoenvironmentally. Similar depositional conditions created both the Fox Hills and Cannonball Formations. There is little likelihood the faunal changes we document resulted from environmental changes, as suggested by Archibald, for we have sampled similar suites of nearshore marine facies on each side of the K-T boundary. Habitat-sensitive molluscan faunas confirm similar origins of most Fox Hills and Cannonball facies (3). Furthermore, given the similarity

of habitats sampled, there is little likelihood that our data result from habitat displacement rather than biological extinction. Significant species-level change in cartilaginous fish faunas occurred across the K-T boundary in the Williston Basin, and apparently globally, at a rate not yet determined.

J. Mark Erickson

*Department of Geology,
St. Lawrence University,
Canton, NY 13617, USA
E-Mail: meri@music.stlawu.edu*

John W. Hoganson

*North Dakota Geological Survey,
Bismarck, ND 58505, USA*

Alan M. Cvancara

*4920 South Oak Street,
Casper, WY 82601, USA*

F. D. Holland Jr.*

*University of North Dakota
Grand Forks, ND 58202, USA*

*Retired

References

1. J. W. Hoganson, J. M. Erickson, A. M. Cvancara, F. D. Holland Jr., *J. Vertebr. Paleontol.* **17**, 53A (1997).
2. E. C. Murphy, D. J. Nichols, J. W. Hoganson, N. F. Forsman, *N.D. Geol. Survey Rept. Invest.* **98**, 1 (1995).
3. A. M. Cvancara, *Contrib. Mus. Paleontol. Univ. Mich.* **20**, 1 (1966); I. Speden, *Yale Peabody Mus. Bull.* **33**, 1 (1970); J. M. Erickson, *Bull. Am. Paleontol.* **284**, 131 (1974).

Corrections and Clarifications

■ In the "Association Affairs" essay "Conversation with the community: AAAS at the millennium" (19 Dec., p. 2066), two co-authors, William T. Golden and Richard S. Nicholson, were inadvertently omitted. The authors should have been listed as follows: "Sheila Jasanoff, Rita Colwell, Mildred S. Dresselhaus, William T. Golden, Robert D. Goldman, M. R. C. Greenwood, Alice S. Huang, William Lester, Simon A. Levin, Marcia C. Linn, Jane Lubchenco, Richard S. Nicholson, Michael J. Novacek, Anna C. Roosevelt, Jean E. Taylor, Nancy Wexler."

Letters to the Editors

Letters may be submitted by e-mail (at science_letters@aaas.org), fax (202-789-4669), or regular mail (Science, 1200 New York Avenue, NW, Washington, DC 20005, USA). Letters are not routinely acknowledged. Full addresses, signatures, and daytime phone numbers should be included. Letters should be brief (300 words or less) and may be edited for reasons of clarity or space. They may appear in print and/or on the World Wide Web. Letter writers are not consulted before publication.



**THE NEW
MINI-PREP 24
FOR AUTOMATED
PLASMID MINI-PREPS**

- **Fast**—up to 24 preps per hr, saving you valuable time.
- **High Purity**—sufficient for automated fluorescent and manual sequencing.
- **Easy Operation**—begin prep with direct loading of bacterial culture. No centrifugation step saves you time.
- **Consistent Results**—up to 6 µg of plasmid per ml. Quality DNA ... time and time again.

**Call now to learn how the
New Improved Mini-Prep 24 can give
you quality DNA and save time by
automating your Plasmid DNA preps!**

1-800-466-7949

**MacCONNELL
RESEARCH**

11339 Sorrento Valley Road
San Diego, CA 92121
(619) 452-2603 • Fax (619) 452-6753
www.macconnell.com

Circle No. 5 on Readers' Service Card