

NET TIPS E-MAIL SECURITY

Mailbox: www.sciencemag.org/dmail.cgi?53844

E-mail Trojan Horses

E-mail is the most common Internet application used at home and at work. Literally millions of users rely on this way to communicate on a daily basis (1). Just imagine a week without the technology of e-mail to see how essential it is to today's scientist.

E-mail has become so ubiquitous that you can now get free e-mail accounts from numerous Web sites, and Web browsers have bundled in free programs as add-on components. Recently, however, some bad news has hit many popular e-mail applications. Security flaws were discovered in late July in the mail programs bundled in Microsoft Explorer and Netscape Communicator. One week later, another security flaw was discovered in Eudora Pro. With literally millions of copies of these flawed programs out there, the potential for disaster is great, and users need to be aware of the flaws as well as the fixes that are available. Additional information about these security flaws in MIME (multipurpose Internet mail extension) buffers are further described in the external bulletins on the Web site of the Australian Computer Emergency Response Team (AusCERT; www.auscert.org.au).

The security issue discovered in Internet Explorer and Netscape Communicator is caused by improper handling of file attachments with very long file names (that is, 200 characters or more). This vulnerability was identified first in July by Finnish testers at the University of Oulu. If you receive an e-mail attachment with a very long file name and read your e-mail using the e-mail reader bundled in these browsers, the long file name will cause the application to shut down unexpectedly. A sophisticated hacker could use this flaw to run malicious code that might wreak havoc in your system. Basically, a buffer overrun could occur, and this vulnerability can be exploited to force programs to execute arbitrary commands with the privileges of the user running the program. Also, attempting to open the corrupted file within the browser might lead to the execution of harmful code.

Tech.Sight is published in the third issue of each month. Contributing editors: Robert Sikorski and Richard Peters, Medsite Communications, Boston, MA, USA. Send your comments by e-mail to tech-sight@aaas.org or via the Web to the mailbox URL with each item.

For Microsoft products, the flaw affects Outlook 98 and Outlook Express that were shipped with Microsoft Internet Explorer 4.0 or 4.01 on Windows 98, Windows 95, Windows NT 4.0, and Windows NT for DEC Alpha, Macintosh, or UNIX. Users are advised to download a security patch that is available at the company's Web site (www.microsoft.com/ie/security).

For Netscape products, the mail and news components of Netscape Communicator versions 4.0 through 4.05 and Netscape Communicator 4.5 Preview Release 1 on the Windows 3.1, 95, 98, and NT platforms could be compromised. This vulnerability does not affect the Macintosh or Unix versions of Communicator. Communicator 4.06, as well as a new version of Communicator 4.5 Preview Release 1, contains a fix for this bug, so users are advised to download upgrades for these browsers from the company's site (<http://home.netscape.com/products/security/resources/bugs/longfile.html>).

For Eudora Pro, the security flaw is different. Hackers could use the ability of Eudora to render hypertext links within an e-mail message. By linking to hostile applets or scripts in an e-mail message, an executable program could be launched by merely clicking on what seems to be an innocent hypertext link. The versions of the software affected include Eudora Pro 4.0 for Windows and Eudora Pro e-mail 4.0.1 for Windows. The problem does not affect Eudora Pro 4.0 for Macintosh. A patch is available from the company's Web site (<http://eudora.qualcomm.com/security.html>).

eudora.qualcomm.com/security.html).

Without a central way to correct the flaws (because millions of copies of these programs are already in use), users need to take charge. Anyone using the versions affected should download and install the fixes immediately. Before doing this, however, make sure you back up your system—something you should do regularly anyway and definitely before installing any new software.

—RICHARD PETERS AND ROBERT SIKORSKI

Reference

1. R. Peters and R. Sikorski, *J. Am. Med. Assoc.* **277**, 1258 (1997).

TECHSIGHTING ENZYMOLGY TECHNIQUE

Mailbox: www.sciencemag.org/dmail.cgi?53842a

Two-Hybridizyme

The two-hybrid system, a technique developed in the yeast research community, is widely used. Armed with a couple of plasmids, yeast strains, and toothpicks, the cardiologist, neurobiologist, and immunologist can use the two-hybrid technique to screen clone libraries for novel proteins that bind to any given target. This straightforward technique often leads researchers into very fertile frontiers.

The two-hybrid system has steadily evolved over the past few years to allow formation of diverse types of complexes (one-hybrid, three-hybrid, and so forth). It has also adapted to the ever-increasing knowledge of protein structure, so that the required fusion protein constructs can be made more intelligently. Nevertheless, enzymologists seem to have avoided the technique, because enzyme-substrate complexes have not been byproducts of two-hybrid studies, until now.

The Tsugimoto group from Japan (1) has recently succeeded in tweaking the two-hybrid system so that it can be used by researchers looking for candidate protein substrates for a specific enzyme. The authors set out to find the substrates for the enzyme caspase-3, a complex of two polypeptide subunits of 10 and 20 kilodaltons (kD). Caspase-3 is a protease that, in many cell types, cleaves target proteins in a cascade that ultimately leads to cell death by apoptosis. There is great interest in these downstream targets for general study and for possible therapeutic strategies.

Applying the two-hybrid method to the problem of caspase-3 binding partners required careful design of the enzyme fusion proteins and modification of the active site of the enzyme. The crystallographic structure of

SITEFINDER SECURITY ADVISORIES

www.auscert.org.au/Information/advisories.html

For anyone interested in keeping up with security flaws in software and hardware, the Australian Computer Emergency Response Team (AusCERT) Web site publishes advisories and alerts in the field of computer security. These bulletins are organized by date of publication and report on security flaws in operating systems, applications, or hardware. Each bulletin consists of a description of the flaw, its impact, recommended solutions, and "workarounds."

www.mednav.com/zone/science

If computer security is important to you, it is well worth your time to check out this site at least once a month to find out what type of new flaws have been reported. In addition, we have collected a list of security resources on our Web site.

TECHSIGHTING
TUMOR GROWTH ASSAY

Mailbox:www.sciencemag.org/dmail.cgi?53842b

caspase-3 showed it to be a tetramer of two 10-kD and two 20-kD subunits. With this knowledge, the investigators wisely chose to fuse the DNA binding domain of the LexA protein to the relatively exposed NH₂-terminus of the p10 subunit. The GAL4 activating domain was fused to the NH₂-terminus of the cDNA library clones. In addition, the p20 subunit was made as a native protein. All were expressed at high concentrations by the yeast ADH1 gene.

Next, the authors chose to modify the active site of the caspase-3 protease to create an inactive complex. The idea was to use a defective enzyme to trap the enzyme substrate at the precleavage step. As do many proteases, caspase-3 has an active-site cysteine that is essential for its enzymatic activity. They substituted a serine residue for the cysteine.

The test experiment consisted of transforming the fusion and native components of caspase-3 into yeast and assaying its binding to a known substrate, CrmA. By a "readout" β -galactosidase assay, the authors showed that only a yeast cell with all components of the system was functional. Next, they moved on to screen a library (mouse embryo cDNA). They isolated 69 clones that could be divided into 13 different clone groups. In vitro assays of the cloned potential targets confirmed that 10 were bona fide substrates for purified caspase-3 enzyme. Sequencing them showed that three clones were parts of the same gene that encodes gelsolin. Additional screens of a mouse thymus library picked out gelsolin clones as well.

Further studies in this report were focused on fleshing out the functional significance of the gelsolin connection to apoptosis. In one series of experiments, they transfected a gelsolin-overexpressing clone into Jurkat cells that are sensitive to apoptotic death. In this cell type, apoptosis can be triggered by adding an antibody to the membrane protein Fas. Amazingly, clones that overexpressed gelsolin were largely immune to the deadly effects of the antibody to Fas. Furthermore, biochemical changes in the gelsolin-overexpressing clones were reflected in a blunting of the apoptotic cascade.

The "two-hybridzyme" approach to finding enzyme substrates might have a broad applicability to other polypeptide-modifying enzymes, such as acetylases, isomerases, and carboxylases. The combination of good enzymatic data, crystallographic coordinates, and yeast genetics provides a new tool for molecular enzymologists.

—ROBERT SIKORSKI AND RICHARD PETERS

Reference

1. S. Kamada et al., *Proc. Natl. Acad. Sci. U.S.A.* **95**, 8532 (1998).

Metastasis in Eggs

Studies of the molecular mechanisms of cancer depend on the use of appropriate model systems. Systems are chosen to emphasize different aspects of the cancer "life cycle." Cell culture systems are used to assay the transformation ability of an oncogene or the blocking effect of a tumor suppressor. Transgenic mice are often used to study multistage genetic changes in a tumor, but how do you study one of the earliest phases of tumor formation, such as intravasation—when a neoplastic cell gains the ability to enter the blood stream and metastasize?

For a simple, in vivo intravasation system, one can inject nude mice with tumor cells and observe the appearance of distant site metastases in the lungs or liver; however, the "readout" of the mouse system is not very sensitive, and the expense of the assay is prohibitive for large-scale drug screening efforts. For an in vitro assay, one could use a cell culture with multi-chamber assay wells that employ a "digestible" membrane. Cells that migrate through the membrane can be measured with high sensitivity, but the in vitro membranes are no match for the complexity found in natural blood vessels and basement membranes.

A system that uses real tissue barriers and affords high sensitivity measurements of intravasation has just been developed by investigators at the Mount Sinai School of Medicine in New York (1). They combined chicken eggs and the polymerase chain reaction (PCR) to uncover new details about the process of cancer cell metastasis.

The underlying anatomy of the chicken egg is key to how the new assay works. The egg's hard shell is lined by a membrane called the chorioallantoic membrane (CAM). The embryo itself lies wrapped in a sac created by the CAM. Separating the CAM and the embryo proper are layers of connective tissue and blood vessels. Because of the asymmetric nature of the CAM, experiments can be designed that involve the upper and lower CAM (polar ends of the egg).

The authors began their investigations by creating an artificial air sac to separate the upper CAM from the egg shell. They then injected human tumor cells or tissues into this air sac and monitored the movement of these neoplastic cells to the lower CAM. Because the only way that cells can get to the lower CAM is by way of blood vessels, the assay measures the foreign cells' intravasation.

To assay metastasized cells in the lower CAM, the authors designed a PCR strategy to amplify human Alu sequences. Alu DNA sequences are plentiful in even one cell, so there are many target copies. In mock mixing experiments, they were able to detect as few as one human tumor cell in 2×10^6 normal chicken cells.

Armed with this powerful assay system, they performed a series of experiments to test the time course of intravasation in the egg and the effect of genetic and chemical modifications on the process. They also looked at different tumor types as well.

For example, they found that human tumor cells (Hep3) were detectable in the lower CAM about 32 hours after seeding the upper CAM. This delay argues for an active entry process and against a simple entry of the cells into leaky blood vessels. Tumor cells detected in the lower CAM soon after the CAM has been seeded would more likely indicate the latter process.

To examine specific molecules, they attempted to block a class of proteases, known as metalloproteinases (MMPs), suspected of playing an essential role in the intravasation process. They added increasing amounts of an MMP inhibitor, marimastat, and showed a dose-dependent block in intravasation potential. In addition, they manipulated the intracellular levels of the receptor urokinase plasminogen activator (uPA) with antisense constructs. By decreasing the level of uPA as much as 70%, they could almost completely block intravasation in the chick system.

These studies are clearly only a hint of what will likely be a wealth of information about the metastatic process that will come out of work with the chicken egg system. The list of obvious future experiments is long and includes cataloging the intravasation potential of existing cell lines, determining the inhibitory effect of potential drugs, and examining the effects of genetic alterations in oncogenes and tumor suppressors.

—ROBERT SIKORSKI AND RICHARD PETERS

Reference

1. J. Kim, W. Yu, K. Kovalski, L. Ossowski, *Cell* **353**, (1998).




PHOTO:KE/NI