

- Genetics* **137**, 175 (1994); M. Palopoli and C.-I. Wu, *ibid.* **138**, 329 (1994); J. R. True, B. S. Weir, C. C. Laurie, *ibid.* **142**, 819 (1996); H. Hollocher and C.-I. Wu, *ibid.* **143**, 1243 (1996).
5. D. E. Perez, C.-I. Wu, N. A. Johnson, M.-L. Wu, *ibid.* **134**, 261 (1993).
 6. D. P. Perez and C.-I. Wu, *ibid.* **140**, 201 (1995).
 7. D. E. Perez, thesis, University of Chicago (1995).
 8. The RT-PCR procedure has been done on all three species of *Drosophila* indicated in Fig. 4. For technical details, refer to the supplementary material at www.sciencemag.org/feature/data/983600.
 9. D. M. Miller *et al.*, *Nature* **355**, 841 (1992) (GenBank accession number X64904); A. Mansouri *et al.*, *Dev. Dynamics* **210**, 53 (1997) (GenBank accession number Z96107); T. Saito, L. Lo, D. J. Anderson, K. Miko-shiba, *Dev. Biol.* **180**, 143 (1996) (GenBank accession number D87748); A. M. Munoz-Marmol *et al.*, GenBank accession number Y10299.
 10. F. Leméunier, J. R. David, L. Tsacas, M. Ashburner, in *The Genetics and Biology of Drosophila*, M. Ashburner, H. L. Carson, J. N. Thompson, Eds. (Academic Press, New York, 1986), vol. 3e, pp. 148–256.
 11. J. Hey and R. M. Kliman, *Mol. Biol. Evol.* **10**, 804 (1993).
 12. F. J. Ayala and D. L. Hartl, *ibid.*, p. 1030.
 13. A. Caccione, E. N. Moriyama, J. M. Gleason, L. Nigro, J. R. Powell, *ibid.* **13**, 1224 (1996).
 14. S. Kumar and S. B. Hedges, *Nature* **392**, 917 (1998).
 15. J. Zhang, H. Rosenberg, M. Nei, *Proc. Natl. Acad. Sci. U.S.A.* **95**, 3708 (1998).
 16. K. Tabuchi, S. Yoshikawa, M. Okabe, K. Sawamoto, H. Okano, paper presented at the 39th Annual *Drosophila* Research Conference, Washington, DC, 25 to 29 March 1998, p. 545C (abstr.).
 17. W. G. Eberhard, *Sexual Selection and Animal Genitalia* (Harvard University Press, Cambridge, MA, 1985); S. Maiti *et al.*, *Genomics* **34**, 304 (1996). C.-I. Wu, N. A. Johnson, M. F. Palopoli, *Trends Ecol. Evol.* **11**, 281 (1996); S. C. Tsaur and C.-I. Wu, *Mol. Biol. Evol.* **14**, 544 (1997); S. C. Tsaur, C.-T. Ting, C.-I. Wu, *ibid.* **15**, 1040 (1998); A. Civetta and R. Singh, *ibid.*, p. 901.
 18. Y.-H. Lee, T. Ota, V. D. Vacquier, *Mol. Biol. Evol.* **12**, 231 (1995); E. C. Metz and S. R. Palumbi, *ibid.* **13**, 397 (1996). W. J. Swanson and V. D. Vacquier, *Science* **281**, 710 (1998); C.-I. Wu and M. F. Palopoli, *Annu. Rev. Genet.* **28**, 283 (1994).
 19. S. B. Carroll, *Development Suppl.*, 217 (1994). M. Averof and N. Patel, *Nature* **388**, 682 (1997).
 20. W. M. Fitch, *Syst. Zool.* **20**, 406 (1971).
 21. T. H. Jukes and C. R. Cantor, in *Mammalian Protein Metabolism*, H. N. Munro, Ed. (Academic Press, New York, 1969).
 22. Supported by National Science Foundation and National Institutes of Health grants to C.-I.W. We are indebted to M. Itoh for collaborating with us in construction of the *D. simulans* and *D. mauritiana* genomic libraries and to T. Hazelrigg for sending us the testis cDNA library. We also thank C. H. Langley, N. Patel, M. F. Polopalli, I. Boussey, C. M. Bergman, P. Andolfatto, M. Long, and S. M. Rollmann for technical consultations and helpful discussions.

6 July 1998; accepted 7 October 1998

Local GABA Circuit Control of Experience-Dependent Plasticity in Developing Visual Cortex

Takao K. Hensch,* Michela Fagiolini, Nobuko Mataga, Michael P. Stryker, Steinunn Baekkeskov, Shera F. Kash

Sensory experience in early life shapes the mammalian brain. An impairment in the activity-dependent refinement of functional connections within developing visual cortex was identified here in a mouse model. Gene-targeted disruption of one isoform of glutamic acid decarboxylase prevented the competitive loss of responsiveness to an eye briefly deprived of vision, without affecting cooperative mechanisms of synapse modification *in vitro*. Selective, use-dependent enhancement of fast intracortical inhibitory transmission with benzodiazepines restored plasticity *in vivo*, rescuing the genetic defect. Specific networks of inhibitory interneurons intrinsic to visual cortex may detect perturbations in sensory input to drive experience-dependent plasticity during development.

After even a brief period of monocular occlusion in early life, input to visual cortex from the closed eye is functionally weakened, then anatomically reduced in size (1). How perturbed levels of neuronal activity are detected to produce changes in connectivity within cortex remains unknown. Inhibitory interactions between inputs may play a role (2), as supported by the profound effects on cortical activity of pharmacologically manipulating γ -aminobutyric acid (GABA) receptors during a period of monocular deprivation. GABA_A agonists induce a robust reverse shift in favor of the

deprived eye (3), whereas antagonists provide mixed results, disrupting plasticity in some cases but yielding no effect under other conditions sufficient to produce continuous epileptiform activity (4). Moreover, these gross drug treatments that shut down or hyperexcite the cortex offer little insight into the normal function of intrinsic networks of inhibitory interneurons during visual cortical plasticity.

Mouse models deficient in the enzymes that produce GABA provide an opportunity to address the role of endogenous inhibitory transmission in cortical plasticity. Distinct genes encode two isoforms of the GABA-synthesizing enzyme, glutamic acid decarboxylase (GAD). The larger 67-kD protein (GAD67) is localized to cell somata and dendrites, providing a constitutive concentration of GABA throughout the cell by a transporter release mechanism (5). In the absence of GAD67, mice die at birth with GABA concentrations less than 10% of those found in the brains of wild-type littermates (6). The 65-kD isoform (GAD65) is

found primarily in the synaptic terminal (7), where it is anchored to vesicles and serves as a reservoir of inactive GAD that can be recruited when additional GABA synthesis is required (8). The smaller isoform of GAD may therefore be specialized to respond to rapid changes in synaptic demand during intense neuronal activity.

We hypothesized that a loss of the GAD65 isoform would selectively reduce fast, intrinsic inhibitory transmission. The deleterious pathology of completely blocking GABAergic synapses pharmacologically could thus be avoided. Indeed, mice carrying a targeted disruption of the GAD65 gene (GAD65 KO mice) survive and develop typical gross cortical morphology and normal adult GABA concentrations because the expression of GAD67 is unaltered (9). Biochemical analysis of various brain regions (Fig. 1A) revealed that unlike in the adult, GAD65 contributes significantly to total GABA concentrations during early postnatal development (10). Consistent with our results, GAD65 expression is elevated in wild-type (WT) animals younger than 3 weeks of age during periods of active synaptogenesis (11). To determine whether the stimulated release of GABA is in fact compromised by loss of the GAD65 protein, we used microdialysis to directly monitor GABA output from the binocular zone of visual cortex *in vivo* (12). Although the basal extracellular concentrations were stable and similar between mutant and wild-type mice, maximal GABA release in response to a brief high-potassium depolarization was significantly reduced in GAD65 KO mice (Fig. 1B).

A physiological consequence of GAD65 deficiency in the visual cortex was enhanced activation in response to visual stimulation. Neurons modulate the expression of certain immediate-early genes in response to sensory stimuli (13). GAD65 KO mice at the peak of the critical period for plasticity exhibited a supernormal sensitivity to visual stimulation after an initial period of dark-rearing (Fig. 1C), as as-

T. K. Hensch, M. Fagiolini, N. Mataga, Laboratory for Neuronal Circuit Development, Brain Science Institute RIKEN, 2-1 Hirosawa, Wako-shi, Saitama 351-0198, Japan. M. P. Stryker, Department of Physiology, University of California, San Francisco, CA 94143, USA. S. Baekkeskov and S. F. Kash, Department of Medicine and Microbiology/Immunology, Hormone Research Institute, University of California, San Francisco, CA 94143, USA.

*To whom correspondence should be addressed. E-mail: hensch@postman.riken.go.jp

REPORTS

essed with the use of *zif268* expression as an endogenous marker of neuronal activity (14). Extracellular single-unit recordings (15) from the binocular zone of visual cortex lacking GAD65 in vivo further revealed a tendency for prolonged

discharge only as light-bar stimuli exited the cell's receptive field (Fig. 2A). However, this did not appear to affect the development of all other visual parameters tested, including spontaneous activity, habituation, retinotopic organization,

orientation and direction selectivity, or receptive field size (Fig. 2, B and C).

The effects of monocular deprivation result from the competitive interaction between left and right eye inputs that first converge at the level of the primary visual cortex (1). The distribution of ocular dominance in the binocular zone of GAD65 KO mice was identical to that of wild-type mice (Fig. 3, A and B, left panels). The response to a 4-day period of monocular vision beginning between postnatal day 25 (P25) and P27 was, however, strikingly different. Wild-type cells shifted their responsiveness in favor of the open eye, as expected (Fig. 3A). Mice lacking GAD65 showed no change in eye preference after deprivation, and cells continued to respond better to contralateral eye input (Fig. 3B).

To determine whether GAD65 KO mice carried a general defect in activity-dependent plasticity, we prepared coronal slices of visual cortex and assayed long-term depression (LTD) and potentiation (LTP) in layer 2/3 of the binocular zone (16). Both wild-type and mutant synaptic responses were persistently potentiated or depressed for periods of up to 1 hour after theta-burst or low-frequency stimulation, respectively (Fig. 3C). These simple correlation-based LTP-LTD paradigms (17) thus failed to predict the immunity to sensory disruption in the intact GAD65 KO mice. Other forms of synapse modification in vitro may better correlate with competitive plasticity in vivo.

The cortical plasticity defect in vivo could have been due to a developmental alteration of the cortical circuit or its plasticity machinery, caused by the lack of GAD65 earlier in development. Alternatively, the absence of plasticity might be an acute consequence of the reduction of cor-

Fig. 1. (A) GABA content in isolated homogenates of young (P18–19; eight WT and six GAD65 KO mice) and adult brains (>5 months old; six WT and four KO mice), normalized to young WT values by region. Open (WT) and solid (KO) bars represent means \pm SEM ($*P < 0.05$, $**P < 0.005$; *t* test). **(B)** Maximal GABA output from WT and GAD65 KO binocular zone in response to brief depolarizing stimuli (100 mM KCl, arrow; three mice, each >P27; $*P < 0.05$, *t* test). Columns are individual 30-min dialysis fractions \pm SEM normalized to mean before high K⁺ perfusion (basal GABA \pm SD = 0.18 \pm 0.02 pmol per fraction for WT, 0.22 \pm 0.11 pmol per fraction for KO). **(C)** Expression of activity-dependent immediate-early gene *zif268* in P28 visual cortex by 90 min of photostimulation (D+P) after 5 days of dark-rearing (D). Northern blots were normalized to house-keeping gene G3PDH (left panel). Sensitivity to visual stimulation in WT and KO mice is shown as (D+P)/D *zif268* expression ratio (right panel, mean \pm SEM, three mice each; $*P < 0.01$, *t* test).





Fig. 2. (A) Prolonged discharge of single-unit activity after visual stimulation in extracellular recordings from GAD65 KO visual cortex. Upper panel: Raster plot of spike response to four sweeps each of computer-generated light-bar stimuli moving across the visual field (5 °/s) in opposite directions (90° and 270°). Trials were randomly interleaved with periods of no stimulus presentation (Spont.). Despite crisp spike onset, neuronal response continued beyond the outer edge of the presumptive receptive field (RF), as defined by stimuli moving in the opposite direction. Lower panel: Proportions of WT and KO cells exhibiting prolonged discharge to stimuli exiting their RF ($n = 670$ and 1112 cells, 28 and 45 mice, respectively; $P < 0.0001$, *t* test). **(B)** Retinotopic organization of WT (open squares) and GAD65 KO (solid diamonds) primary visual cortex. RF-center azimuths are plotted versus electrode position relative to vertical meridian. Inset: Correlation coefficients for three WT and four KO regressions ($P = 0.2$, *t* test). **(C)** RF size (left panel, 82 WT and 79 KO cells; $P = 0.1$, *t* test) and proportion of cells exhibiting preference for light-bar stimuli of particular orientation (right panel, $n = 670$ and 1112 cells, 28 WT and 45 KO mice, respectively; $P = 0.3$, *t* test).




tical inhibition during the critical period. We thus attempted to rescue the plasticity defect in vivo by briefly enhancing inhibition in mutant animals. The postsynaptic impact of reduced GABA release could be compensated with benzodiazepine agonists. These drugs increase the open probability and channel conductance of GABA_A receptors selectively in a use-dependent manner, as they are inert in the absence of synaptic GABA release (18). Moreover, benzodiazepine binding sites are associated with intrinsic cortical elements rather than with thalamocortical axons or other subcortical inputs (19).

To identify the precise locus where rapid GABAergic transmission is required for plasticity in vivo, we administered diazepam locally into one hemisphere of the brains of the mutant mice during a period of monocular eyelid suture. The localization of diazepam to the infused visual cortex was biochemically confirmed (20), whereas it remained virtually undetectable in the adjacent temporal cortex, frontal regions, or opposite hemisphere (Fig. 4A, left). Monocular deprivation produced a complete ocular dominance shift in the infused mutant visual cortex (Fig. 4A, right), whereas no rescue of plasticity was observed by administering vehicle solutions in the same side or diazepam into the opposite hemisphere (Fig. 4B). Global diazepam treatment by intraventricular injection during a concurrent 4-day monocular deprivation also restored plasticity to GAD65 KO mice comparable to that of the wild-type mice [contralateral bias index (CBI) = 0.47 ± 0.03 versus 0.43 ± 0.03; 103 and 109 cells, 5 KO and 6 WT animals, respectively; *P* = 0.4, *t* test] (15). Taken together, fast inhibitory transmission via GABA_A-mediated circuitry intrinsic to visual cortex is both necessary and sufficient to detect an imbalance in activity between competing inputs from the two eyes, and had not been irreparably damaged by GAD65 deletion.

If intracortical inhibition determines the threshold for detecting competition, then the residual GABA found in visual cortex lacking GAD65 (Fig. 1) should allow limited plasticity with an extreme imbalance of input. We maximized the contrast between excitatory input from the two eyes by completely silencing the retinal activity of one eye with tetrodotoxin (TTX) injections for 4 days during the critical period (16). A significantly stronger ocular dominance shift was observed with TTX than with monocular eyelid suture in wild-type mice (mean CBI = 0.33 ± 0.04 versus 0.49 ± 0.02; 4 and 10 mice, respectively; *P* < 0.001, *t* test). This drastic imbalance in input activity produced detectable changes in only some GAD65 KO mice [mean CBI = 0.54 ± 0.06; 4 mice (21)], significantly weaker than in wild-type animals (*P* < 0.05, *t* test) and more variable

Fig. 3. (A) Ocular dominance scores for WT mice were normally skewed in favor of the contralateral eye (left panel, 66 cells, three mice). Four days of monocular eyelid suture (solid circle) during the critical period shifted ocular dominance toward the open, ipsilateral eye (open circle), yielding a balanced distribution (right panel, 132 cells, six mice; *P* < 0.0001, χ^2 test). Plasticity in rodents is less pronounced than in other mammals (24). **(B)** Ocular dominance distribution of nondeprived GAD65 KO mice is similar to WT (left panel, 99 cells, five mice; *P* = 0.2, χ^2 test). Four days of monocular vision yielded no shift in favor of the open eye (right panel, 108 cells, five mice; *P* = 0.4 vs. normal KO and *P* < 0.0001 vs. deprived WT, χ^2 test). As an indicator of the relative response to the two eyes, the CBI (16), shown in the upper right corner of each distribution, takes values of 0 and 1 for complete ipsilateral or contralateral eye dominance, respectively. **(C)** LTP and LTD in WT (open circles) and KO (solid circles) visual cortical slices after theta-burst (left panel) or low-frequency (1 Hz, right panel) stimulation, respectively [8 slices, seven mice from each group (LTP) and 7 slices, six WT and seven KO mice (LTD); *P* = 0.2 for both comparisons at 20 min, *t* test].




than in deprived mutants receiving diazepam (Fig. 4C).

Local cortical circuits may then instruct the reorganization of both intra- and thalamocortical connections that are known to accompany monocular deprivation (1). A delicately balanced recruitment of excitation and inhibition detects the disparity between competing sensory inputs and drives experience-dependent changes in visual cortex. Complete restoration of plasticity in GAD65 KO mice was achieved by use-dependent potentiation of GABA_A responses that produced only a modest, albeit significant, reduction of undamped recurrent excitation (54 ± 7% versus 74 ± 5% of cells exhibiting prolonged discharge, 18 diazepam-treated and 45 untreated mice; *P* < 0.03, *t* test). Molecular cues [such as neurotrophins (22)] must be exquisitely tuned to spatiotemporal patterns of activity for subtle modulation of stimulated GABA release to have such profound effects on plasticity. Notably, the diffuse activation of GABA_A receptors with muscimol to silence cortical activity reverses plasticity (3), whereas the selective enhancement of intrinsic inhibitory transmission with diazepam sharpens it even in wild-type animals (Fig. 4C). Competitive refinement of visual

cortex thus requires a threshold level of GABAergic transmission, making it fundamentally different from LTP or LTD models that are impeded by inhibition (23) and consistent with recent evidence dissociating in vivo from in vitro plasticity (24).

The many cortical interneuron subtypes operate and mature in a lamina-specific fashion (25) while responding differentially to monocular deprivation, neuromodulators, and input from other GABAergic cells (26). Only a specific subset of this extremely elaborated network may sculpt cortical activity to define the connections that will undergo refinement. GAD65 localizes preferentially to GABAergic terminals targeted to dendrites rather than to cell somata (8, 27). Moreover, fast (GABA_A) or slow (GABA_B) synaptic inhibition may originate from morphologically distinct classes of interneuron (28). Diazepam rescue of plasticity in GAD65 mutants focuses attention on GABA_A connections with appropriate subunit composition for benzodiazepine binding (20). We have identified an animal model in which plasticity of the intact visual cortex is robustly disrupted. Our findings provide a tool for dissecting the unique local circuit properties that drive experience-dependent plasticity.

Fig. 4. (A) Local diazepam treatment of visual cortex restores plasticity to GAD65 KO mice in vivo. Left panel: Directed infusion restricted diazepam from regions beyond the visual cortex ipsilateral to the cannula (solid circles), as well as throughout the opposite hemisphere (open circles, mean \pm SEM, three mice). Right panel: Ocular dominance shifted fully in the KO binocular zone exposed to diazepam concurrent with a brief period of monocular deprivation during the critical period (114 cells, four mice; $P < 0.0001$, χ^2 test vs. nondeprived KO; compare with Fig. 3B). **(B)** Neither vehicle treatment (left panel, 99 cells, four mice; $P < 0.0001$, χ^2 test vs. diazepam KO above) nor diazepam infusion into the hemisphere opposite the recording site (right panel, 77 cells, three mice; $P < 0.0001$, χ^2 test vs. diazepam KO above) restored the effect of monocular deprivation to GAD65 KO mice. **(C)** Monocular TTX injections for 4 days during the critical period produced significantly less plasticity in KO than in WT mice ($P < 0.05$, t test). Untreated and vehicle-treated monocularly deprived (MD) mice are grouped together, as are ventricle- and cortex-infused diazepam-treated animals. Shaded region indicates the range of nondeprived CBIs for both WT and KO mice. Each symbol represents one animal.



References and Notes

1. T. N. Wiesel and D. H. Hubel, *J. Neurophysiol.* **26**, 1003 (1963); C. R. Olson and R. D. Freeman, *ibid.* **38**, 26 (1975); C. J. Shatz and M. P. Stryker, *J. Physiol.* **281**, 267 (1978); A. Antonini and M. P. Stryker, *Science* **260**, 1819 (1993); C. J. Shatz, *Neuron* **5**, 745 (1990).
2. K. E. Kratz, P. D. Spear, D. C. Smith, *J. Neurophysiol.* **39**, 501 (1976); F. H. Duffy, S. R. Snodgrass, J. R. Burchfiel, J. L. Conway, *Nature* **260**, 256 (1976); C. Blakemore and M. J. Hawken, *J. Physiol. (London)* **327**, 463 (1982); A. M. Sillito, J. A. Kemp, C. Blakemore, *Nature* **291**, 318 (1981); G. D. Mower and W. G. Christen, *Dev. Brain Res.* **45**, 211 (1989).
3. H. O. Reiter and M. P. Stryker, *Proc. Natl. Acad. Sci. U.S.A.* **85**, 3623 (1988); Y. Hata and M. P. Stryker, *Science* **265**, 1732 (1994).
4. A. S. Ramoa, M. A. Paradiso, R. D. Freeman, *Exp. Brain Res.* **73**, 285 (1988); T. O. Viden, N. W. Daw, R. C. Collins, *Brain Res.* **371**, 1 (1986).
5. M. G. Erlander, N. J. K. Tillakaratne, S. Feldblum, N. Patel, A. J. Tobin, *Neuron* **7**, 91 (1991); S. Feldblum, M. G. Erlander, A. J. Tobin, *J. Neurosci. Res.* **34**, 689 (1993); D. L. Kaufman, C. R. Houser, A. J. Tobin, *J. Neurochem.* **56**, 720 (1991).
6. H. Asada et al., *Proc. Natl. Acad. Sci. U.S.A.* **94**, 6496 (1997); B. G. Condie, G. Bain, D. I. Gottlieb, M. R. Capecchi, *ibid.*, p. 11451.
7. Y. Guo, I. V. Kaplan, N. G. F. Cooper, G. D. Mower, *Dev. Brain Res.* **103**, 127 (1997); T. Fukuda, Y. Aika, C. W. Heizmann, T. Kosaka, *J. Comp. Neurol.* **395**, 177 (1998).

8. A. Reetz et al., *EMBO J.* **10**, 1275 (1991); D. L. Martin, S. B. Martin, S. J. Wu, N. Espina, *J. Neurosci.* **11**, 2725 (1991).
9. H. Asada et al., *Biochem. Biophys. Res. Commun.* **229**, 891 (1996); S. F. Kash et al., *Proc. Natl. Acad. Sci. U.S.A.* **94**, 14060 (1997).
10. Mice carrying a functional disruption of GAD65 were generated by insertion of a neomycin-resistance cassette into exon 1 of the gene on the C57BL/6 background, as described (9). Homogenates of various brain regions prepared from mice at different ages were analyzed for GABA content using high-performance liquid chromatography (HPLC; Eicom) with fluorescence detection (excitation, 340 nm; emission, 440 nm; L-1000, Hitachi) after ophthalmaldehyde (OPA) reagent derivatization by a computer-controlled autoinjector with column switching (CMA 200). Amino acids were separated on a Cosmosil 5C18 reversed-phase column (36°C; Nacal Tesque, Japan). The mobile phase was 50 mM sodium acetate buffer with 25% acetonitrile (pH 5.9) [N. Mataga, K. Imamura, Y. Watanabe, *Brain Res.* **551**, 61 (1991)].
11. K. F. Greif, M. G. Erlander, N. J. K. Tillakaratne, A. J. Tobin, *Neurochem. Res.* **16**, 235 (1991); K. F. Greif et al., *Dev. Biol.* **153**, 158 (1992).
12. or extracellular GABA measurements in vivo, mice (>P27) were mounted on a stereotaxic frame, and a microdialysis probe (membrane length 1 mm; CMA 11, CMA/Microdialysis AB, Sweden) was implanted into the binocular region of visual cortex. Artificial cerebrospinal fluid (ACSF) was perfused (1.0 μ l/min) and samples were taken every 30 min for GABA analysis (10) using a microfraction collector (CMA 140). ACSF contained 119 mM NaCl, 2.5

mM KCl, 1.3 mM MgSO₄, 1.0 mM NaH₂PO₄, 26.2 mM NaHCO₃, 2.5 mM CaCl₂, and 11 mM glucose. After equilibration (60 min), five or six basal values were collected, then ACSF containing 100 mM KCl was perfused for 30 min, followed by five or six recovery samples. Dead volume from probe output to fraction collector was 8.4 μ l (~8 min).

13. L. Kaczmarek and A. Chaudhuri, *Brain Res. Rev.* **23**, 237 (1997); P. F. Worley et al., *Proc. Natl. Acad. Sci. U.S.A.* **88**, 5106 (1991).
14. WT and KO mice (six animals each) were reared in total darkness for 5 days at the peak of the critical period (P23–28). The mice in the dark-reared group were killed immediately in the dark room, and the remainder were exposed to light from a photostimulator (1 Hz, 20 J, 90 min; NEC). Northern (RNA) blot analysis for *zif268* expression in the binocular zone was performed as described [N. Mataga et al., *Neurosci. Lett.* **218**, 149 (1996)], quantified by densitometer (Molecular Dynamics), and normalized to the housekeeping gene G3PDH.
15. Mice were prepared blind to genotype for electrophysiological recording in vivo under Nembutal (50 mg/kg, Abbot)/chlorprothixene (0.2 mg, Sigma) anesthesia using standard techniques (24) [J. A. Gordon and M. P. Stryker, *J. Neurosci.* **16**, 3274 (1996)]. For each animal, five to eight cells (>75 μ m apart) were recorded in each of four to six vertical penetrations spaced evenly (>200 μ m intervals) across the mediolateral extent of primary visual cortex to map the monocular and binocular zones and avoid sampling bias. Receptive fields of isolated single units were plotted on a tangent screen with a hand-held projection lamp or computer-generated stimuli. Cells were assigned ocular dominance scores according to the seven-point classification scheme of Hubel and Wiesel (7). A weighted average of the bias toward one eye or the other, the CBI, was calculated for each binocular zone according to the formula $CBI = [(n_1 - n_2) + (2/3)(n_2 - n_6) + (1/3)(n_3 - n_5) + N]/2N$, where N is the total number of cells and n_x is the number of cells with an ocular dominance score equal to x . For monocular deprivation experiments, eyelid margins were trimmed and sutured under halothane anesthesia at the peak of the critical period (for 4 days beginning at P25–27). In other cases, repeated intraocular injections (<350 nl) of TTX (3 mg/ml, Sankyo) or citrate vehicle (0.5%, Sigma) solution were made into one eye. TTX efficacy was confirmed by pupil dilation over a 4-day period. All recordings were contralateral to the deprived eye.
16. Coronal slices (400 μ m) through the binocular zone were prepared blind to genotype during the critical period for monocular deprivation effects (P24–33) and maintained at 33°C in oxygenated ACSF (95% O₂/5% CO₂; pH 7.4). Changes in the magnitude of extracellular field potential peak amplitude were recorded with a 1 M NaCl electrode (1 to 3 megohms) inserted into layer 2/3 in response to theta-burst or low-frequency (1 Hz) stimulation of layer 4, as described (24). Both size (half-maximal amplitude, >1 mV) and shape (half-width, <5 ms) of extracellular field potentials did not differ between genotypes and were routinely confirmed to be synaptic by glutamate receptor antagonists applied to the bath at the end of each experiment.
17. J. P. Rauschecker, *Physiol. Rev.* **71**, 587 (1991); T. Tsumoto, *Prog. Neurobiol.* **209**, 209 (1992); A. Kirkwood, M. G. Rioult, M. F. Bear, *Nature* **381**, 526 (1996); W. Singer, *Science* **270**, 758 (1995); L. C. Katz and C. J. Shatz, *ibid.* **274**, 1133 (1996).
18. R. E. Study and J. L. Barker, *Proc. Natl. Acad. Sci. U.S.A.* **78**, 7180 (1981); J. F. Tallman and D. W. Gallagher, *Annu. Rev. Neurosci.* **8**, 21 (1984); R. L. MacDonald and R. W. Olsen, *ibid.* **17**, 569 (1994); C. J. Rogers, R. E. Twyman, R. L. MacDonald, *J. Physiol. (London)* **475**, 69 (1994); M. Eghbali, J. P. Curmi, B. Birnir, P. W. Gage, *Nature* **388**, 71 (1997).
19. C. Shaw, C. Aoki, M. Wilkinson, G. Prusky, M. Cynader, *Dev. Brain Res.* **37**, 67 (1987); Z. Xiang, J. R. Huguenard, D. A. Prince, *J. Physiol. (London)* **506**, 715 (1998); E. Sigel and A. Buhr, *Trends Pharmacol. Sci.* **18**, 425 (1997); E. Costa, *Annu. Rev. Pharmacol. Toxicol.* **38**, 321 (1998).
20. In rescue experiments, either diazepam (2 mg/ml, Wako) or vehicle (50% propylene glycol, Wako) solution was injected daily into the ventricles (1.5 μ l each side) beginning 1 or 2 days before and concurrent with 4 days of

Synaptic Segregation at the Developing Neuromuscular Junction

Wen-Biao Gan and Jeff W. Lichtman*

Throughout the developing nervous system, competition between axons causes the permanent removal of some synaptic connections. In mouse neuromuscular junctions at birth, terminal branches of different axons are intermingled. However, during the several weeks after birth, these branches progressively segregated into nonoverlapping compartments before the complete withdrawal of all but one axon. Segregation was caused by selective branch atrophy, detachment, and withdrawal; the axon branches that were nearest to the competitor's branches were removed before the more distant branches were removed. This progression suggests that the signals that mediate the competitive removal of synapses must decrease in potency over short distances.

Competitive synaptic rearrangements change the number and strength of axonal connections with target cells in many parts of the nervous system (1). At each neuromuscular junction, for example, motor axons compete until all but one are removed (2). By iontophoretic application of lipophilic dye (3, 4), we observed synaptic competition between individual axon terminals at the developing neuromuscular junction of the mouse.

We began by looking at neuromuscular junctions during the second postnatal week [postnatal days 9 and 10 (P9-10)], when most muscle fibers (88%) were singly innervated and the rest (12%) were innervated by two axons (5). At the singly innervated junctions ($n = 242$; 20 muscles), the entire postsynaptic receptor territory was overlain by one axon, as observed by staining with lipophilic dyes or with nerve-specific antibodies (6). However, at the remaining 12% of the junctions, a labeled axon occupied only part of the receptor territory (Fig. 1, A through D). Subsequent staining with antibodies to neurofilaments or with tetanus toxin (6) showed that, in each of these cases ($n = 33$), the remaining part of the neuromuscular junction was innervated by another axon (Fig. 1A). In these dually innervated junctions, the labeled terminals of each individual axon occupied a variably sized but contiguous region extending from one edge of the junction toward the center (Fig. 1, A through D). Thus, in the second postnatal week, each axon's synaptic territory is confined to a separate region of the junction.

By contrast, at earlier stages (P1-3), a single axon typically occupies multiple small regions that are distributed throughout the neuromuscular junction (Fig. 1, E and F). As

expected, the sites that are occupied by different axons are intermingled ($n = 26$) (Fig. 2A) (7). The difference in the distribution of axon terminals between P1-3 and P9-10 could mean that the areas that are occupied by different axons segregate from each other over time. Alternatively, a small number of junctions may already be segregated at birth, and these junctions may retain multiple innervation longer than intermingled ones. To explore these possibilities, we used different lipophilic dyes to separately label each axon that innervated the same neuromuscular junction, and we measured the separation between the terminal areas that were occupied by the two competing axons (8) (Fig. 2, A through C). We found no junctions at birth in which axons had completely nonoverlapping territories. Thereafter, both the absolute separation (the distance between two centroids) and the relative separation [the distance normalized to the total length of the junction (Fig. 2E)] of the terminal areas gradually increased. In most cases, the separation was only partial at P3-4 (Fig. 2, B and E) and was complete by P7-10 (Fig. 2, C and E). By P16-17, when very few junctions (1.4%) were still multiply innervated, axon territories in all of the multiply innervated junctions (8 out of 8) showed a high degree of separation with an uninnervated gap in between (9) (Fig. 2, D and E). Therefore, neuromuscular junctions become progressively compartmentalized as development proceeds.

To learn how this segregation of terminals occurred, we examined multiply innervated junctions at P5-7 before segregation was complete in all junctions ($n = 62$). We found that, at multiply innervated junctions, individual axons often (30 out of 62) possessed two types of branches, thick and thin (Fig. 3, A and B). The thin axon branches were always nearest to or within the territory occupied by the competing axon, whereas the thickest branches were located on the perimeter of the junction at the great-

monocular deprivation. A >100-fold saturating dose of diazepam was determined by recording GABA_A responses in visual cortical slices (16), then infused to ensure adequate drug diffusion in vivo. Local rescue was achieved with low-flow osmotic minipumps (0.5 μ l/hour; Alzet 1007, Alza) containing drug or vehicle solution connected to cannulae (30 gauge) that were stereotactically implanted into one hemisphere under sterile surgical conditions 2 days before eyelid suture. Control experiments demonstrated that continuous infusion of TTX or dye solutions from osmotic minipumps could be restricted to ipsilateral visual cortex without disturbing the gross morphology of the binocular zone at least 1.5 mm away from the cannula tip. Drug was infused continuously for 4.5 hours at a high rate (3 μ l/hour) to determine the spread of diazepam. In agreement with its known rapid breakdown in vivo [L. A. Berrueta, B. Gallo, F. Vicente, *J. Pharm. Biomed. Anal.* **10**, 109 (1992)], diazepam was not detectable after 1 week of pumping at low rates. A diazepam detection assay was developed using reversed-phase HPLC (UV detection at 240 nm) after organic extraction: Brain tissue (~20 mg) was sonicated with ethyl acetate (400 μ l) and centrifuged (8000g, 15 min), the supernatant extracted on a mechanical shaker (1 M tris-HCl, pH 9.0, 20 min), and the organic layer evaporated under a gentle N₂ stream. The residue was reconstituted in ethyl acetate (100 μ l) and back-extracted with 6 N HCl (40 μ l); the aqueous phase was adjusted to neutral (3 M tris buffer, pH 11.6) and injected onto an HPLC apparatus. The mobile phase was 50 mM sodium phosphate buffer (pH 6.4) with 38% acetonitrile and 0.1% triethylamine.

21. Similarly weak and variable plasticity was observed in GAD65 KO versus WT mice that had experienced deprivations spanning the entire critical period (P17-36), despite the longer temporal integration and potential additional contribution of anatomical changes to the functional disconnection of deprived eye input (7). $CB1 = 0.54 \pm 0.04$ versus 0.41 ± 0.02 ; six KO and four WT mice, respectively ($P < 0.05$, t test).
22. F. Zafra, E. Castren, H. Thoenen, D. Lindholm, *Proc. Natl. Acad. Sci. U.S.A.* **88**, 10037 (1991); H. Thoenen, *Science* **270**, 593 (1995); A. K. McAllister, D. C. Lo, L. C. Katz, *Neuron* **17**, 1057 (1996); T. Bonhoeffer, *Curr. Opin. Neurobiol.* **6**, 119 (1996); R. Sala et al., *Eur. J. Neurosci.* **10**, 2185 (1998); L. C. Rutherford, S. B. Nelson, G. G. Turrigiano, *Neuron* **21**, 521 (1998).
23. A. Artola and W. Singer, *Nature* **330**, 649 (1987); M. F. Bear and A. Kirkwood, *Curr. Opin. Neurobiol.* **3**, 197 (1993); S. M. Dudek and M. J. Friedlander, *Neuron* **16**, 1097 (1996); K. D. Miller, *ibid.* **17**, 371 (1996).
24. T. K. Hensch and M. P. Stryker, *Science* **272**, 554 (1996); T. K. Hensch et al., *J. Neurosci.* **18**, 2108 (1998).
25. J. DeFelipe, *Cereb. Cortex* **3**, 273 (1993); Y. Gonchar and A. Burkhalter, *ibid.* **7**, 347 (1997); Y. Kang, T. Kaneko, H. Ohishi, K. Endo, T. Araki, *J. Neurophysiol.* **71**, 280 (1994); J. F. M. Van Bredenoord and W. J. Spain, *ibid.* **74**, 1149 (1995); H. J. Luhmann and D. A. Prince, *ibid.* **65**, 247 (1991); Y. Komatsu, *Dev. Brain Res.* **8**, 136 (1983).
26. R. K. Carder, S. S. Leclerc, S. H. C. Hendry, *Cereb. Cortex* **6**, 271 (1996); Y. Kawaguchi, *J. Neurophysiol.* **78**, 1743 (1997); D. Parra, A. I. Gulyas, R. Miles, *Neuron* **20**, 483 (1998); Y. Kawaguchi and T. Shindou, *J. Neurosci.* **18**, 6963 (1998); Z. Xiang, J. R. Huguenard, D. A. Prince, *Science* **281**, 985 (1998); G. Tamas, P. Somogyi, E. Buhl, *J. Neurosci.* **18**, 4255 (1998).
27. A. Gulyas, R. Miles, N. Hajos, T. F. Freund, *Eur. J. Neurosci.* **5**, 1729 (1993); E. Buhl, K. Halasy, P. Somogyi, *Nature* **368**, 823 (1994); I. Soltesz, D. K. Smetters, I. Mody, *Neuron* **14**, 1273 (1995); R. Miles et al., *ibid.* **16**, 815 (1996); G. Tamas, E. H. Buhl, P. Somogyi, *J. Physiol. (London)* **500**, 715 (1997).
28. Y. Kawaguchi and Y. Kubota, *Cereb. Cortex* **7**, 476 (1997); L. S. Benardo, *J. Physiol. (London)* **476**, 203 (1994); Y. Kawaguchi, *Exp. Brain Res.* **88**, 33 (1992); S. Sugita, S. W. Johnson, R. A. North, *Neurosci. Lett.* **134**, 207 (1992); M. Segal, *Brain Res.* **511**, 163 (1990); A. M. Thomson and D. C. West, *Soc. Neurosci. Abstr.* **23**, 379.7 (1997).

29. We thank S. Fujishima and K. Hartman for excellent technical assistance.

27 May 1998; accepted 19 October 1998

Department of Anatomy and Neurobiology, Washington University School of Medicine, 660 South Euclid Avenue, Box 8108, St. Louis, MO 63110, USA.

*To whom correspondence should be addressed. E-mail: jeff@thalamus.wustl.edu