

Role of α -Dystroglycan as a Schwann Cell Receptor for *Mycobacterium leprae*

Anura Rambukkana,* Hiroki Yamada, George Zanazzi, Todd Mathus, James L. Salzer, Peter D. Yurchenco, Kevin P. Campbell, Vincent A. Fischetti

α -Dystroglycan (α -DG) is a component of the dystroglycan complex, which is involved in early development and morphogenesis and in the pathogenesis of muscular dystrophies. Here, α -DG was shown to serve as a Schwann cell receptor for *Mycobacterium leprae*, the causative organism of leprosy. *Mycobacterium leprae* specifically bound to α -DG only in the presence of the G domain of the $\alpha 2$ chain of laminin-2. Native α -DG competitively inhibited the laminin-2-mediated *M. leprae* binding to primary Schwann cells. Thus, *M. leprae* may use linkage between the extracellular matrix and cytoskeleton through laminin-2 and α -DG for its interaction with Schwann cells.

Pathogenic bacteria are adapted to exploit a variety of host cell functions, and host cell receptors mostly serve as the initial target for bacterial interaction with a specific cell type (1, 2). However, not much is known about the bacterial receptors in the nervous system and how bacteria interfere with these neuronal cell receptor-associated functions. *Mycobacterium leprae*, the causative organism of leprosy, is an intracellular pathogen that invades the Schwann cell of the peripheral nervous system (3). During infection, *M. leprae* causes significant damage to peripheral nerves leaving patients with disabilities and deformities (4). Although antibiotic therapy is an effective cure of leprosy, it does not reverse the nerve function loss in these patients (5). Understanding the mechanisms of *M. leprae*-Schwann cell interaction may yield new therapeutic strategies for the prevention of nerve damage.


Dystroglycan (DG), a component of the dystrophin-glycoprotein complex, is a laminin receptor encoded by a single gene and cleaved by posttranslational processing into two proteins, peripheral membrane α -DG and transmembrane β -DG (6). Whereas α -DG interacts with laminin-2 in the basal lamina, β -DG appears to bind to dystrophin-containing cytoskeletal proteins in muscles and peripheral nerves

(7). DG is involved in agrin- and laminin-induced acetylcholine receptor clustering at neuromuscular junctions (8), morphogenesis (9), early development (10), and the pathogenesis of muscular dystrophies (6, 7). The loss or a defect of laminin-2- α -DG interaction causes certain types of muscular dystrophy and peripheral neuropathy (11, 12). We recently showed that laminin-2 in the basal lamina of the Schwann cell-axon unit serves as an initial target for *M. leprae* interaction with peripheral nerves (13). Laminin-2, which comprises $\alpha 2$, $\beta 1$, and $\gamma 1$ chains (14), anchors to Schwann cells through laminin receptors (15). α -DG serves as a receptor on the Schwann cell that interacts with laminin-2 in the basal lamina surrounding the Schwann cell-axon unit (7, 16).

Fig. 1. *Mycobacterium leprae* binding to immobilized α -DG is mediated by the G domain of the laminin- $\alpha 2$ chain. (A) Quantification of *M. leprae* binding to immobilized native α -DG purified from peripheral nerves (PN- α DG) (black bars) and skeletal muscles (muscle- α DG) (gray bars) in the presence of recombinant fragments of the laminin- $\alpha 2$ chain. *Mycobacterium leprae* (5×10^8) was preincubated with rLN $\alpha 2$ G or rLN $\alpha 2$ (VI-IVb)' (10 μ g/ml), and the mixtures were overlaid onto immobilized α -DG (50 μ g/ml). The number of adherent *M. leprae* (ML) within a 0.25 mm² grid area of each well was quantified after 60 min of incubation, and the data were expressed as the mean \pm SD values from five to six wells. Three additional experiments gave similar results. The inset is a laminin-2 molecule showing the location of the NH₂-terminal (VI-IVb)' fragment and the COOH-terminal G domain of the $\alpha 2$ chain. (B) Coomassie blue-stained SDS-PAGE gel showing the purified native α -DG preparations and recombinant laminin- $\alpha 2$ chain fragments used in the study: peripheral nerve α -DG (120 kD), muscle α -DG (156 kD), rLN- $\alpha 2$ (VI-IVb)' (116 kD), and rLN- $\alpha 2$ G (120 kD).

To determine the role of α -DG in the *M. leprae* interaction with Schwann cells, we first examined the binding of *M. leprae* (17) to native α -DG purified from peripheral nerves (18) in the presence or the absence of recombinant fragments of the laminin- $\alpha 2$ chain (19) (Fig. 1, A and B). In a solid-phase assay (20), *M. leprae* bound to immobilized α -DG only in the presence of the COOH-terminal fragment of the laminin- $\alpha 2$ chain (rLN $\alpha 2$ G; Fig. 1A). *Mycobacterium leprae* also bound to muscle α -DG only in the presence of rLN- $\alpha 2$ G (Fig. 1A). *Mycobacterium leprae* binding to α -DG of both peripheral nerve and muscle was increased by >95% with a concentration of 10 μ g/ml (0.1 μ g per well) of rLN $\alpha 2$ G. Even higher concentrations (100 μ g/ml) of the NH₂-terminal r(VI-IVb)' fragment of laminin- $\alpha 2$ chain or the G domain of the laminin- $\alpha 1$ chain (rLN $\alpha 1$ G) had no effect on *M. leprae* binding to α -DG (110 ± 24 and 130 ± 31 bacteria per 0.25 mm², respectively), suggesting that the G domain of the laminin- $\alpha 2$ chain specifically mediated *M. leprae* binding to α -DG. Thus, LN $\alpha 2$ G has two binding sites, one for *M. leprae* and the other for α -DG, and the G domain forms a bridge between *M. leprae* and α -DG. Additionally, the activity of merosin/ $\alpha 2$ laminins (a mixture of laminin-2 and laminin-4) on *M. leprae*- α -DG interaction yielded results similar to those with rLN $\alpha 2$ G at equal molar ratio (1010 ± 110 and 1290 ± 161 bacteria per 0.25 mm², respectively).

Comparison of rLN $\alpha 2$ G-mediated *M. leprae* binding to native (Fig. 1B) versus fusion proteins of α -DG (Fig. 2, B and C) showed that the bacteria strongly bound only to the native α -DG in a concentration-dependent manner (Fig. 2A). Because the native confor-



A. Rambukkana and V. A. Fischetti, Laboratory of Bacterial Pathogenesis and Immunology, Rockefeller University, New York, NY 10021, U.S.A. H. Yamada and K. P. Campbell, Howard Hughes Medical Institute, (HHMI), Department of Physiology and Biophysics, and Department of Neurology, University of Iowa College of Medicine, Iowa City, IA 52242, U.S.A. G. Zanazzi and J. L. Salzer, Departments of Cell Biology and Neurology and the Kaplan Cancer Center, New York University Medical Center, New York, NY 10016, U.S.A. T. Mathus and P. D. Yurchenco, Department of Pathology, Robert Wood Johnson Medical School, Piscataway, NJ 08854, U.S.A.

*To whom correspondence should be addressed. E-mail: rambuka@rockvax.rockefeller.edu

REPORTS

mation of α -DG seems unnecessary for the laminin interaction, as denatured DG also binds laminins (7) and *M. leprae*+rLN α 2G

(21), glycosylation is the most likely post-translational modification that contributes to α -DG's interaction with *M. leprae* through

the LN- α 2G domain. Because the rLN α 2G-mediated *M. leprae* binding to α -DG was sensitive to periodate (Fig. 2D), the carbohydrate moieties of α -DG are likely important for rLN α 2G-mediated *M. leprae* interactions. Although α -DGs are differentially glycosylated in different tissues (for example, peripheral nerve and muscle) of the same species, which contributes to different molecular sizes (22) (Fig. 1B), glycosylation of α -DG in a given tissue (for example, peripheral nerve) is almost identical in mouse, rabbit, cow, and human (6, 22–24). Further characterization of the rLN α 2G-mediated *M. leprae*- α -DG interaction showed that this binding is completely abolished by EDTA, indicating the crucial role of calcium for the interaction of the G domain with α -DG. Moreover, the lack of inhibitory effects of heparin on rLN α 2G-mediated *M. leprae* binding to α -DG (Fig. 2E) suggests that the heparin binding site of the G domain of the laminin-2 molecule is different from the α -DG binding site. Thus, the G domain is the α -DG binding site of the laminin-2 molecule, and this interaction is dependent on calcium and is largely mediated by the carbohydrate moieties of α -DG. This may be of significant physiological relevance in muscular dystrophies because the loss of laminin-2 interaction with α -DG is critical for the disease pathogenesis (6, 7).

To determine whether peripheral nerve α -DG serves as a Schwann cell receptor for *M. leprae*, we purified Schwann cells from rat sciatic nerves (25). These primary Schwann cells strongly expressed α -DG but showed almost no deposition of LN α 2G in early cultures (Fig. 3C) (21); they were also devoid of certain laminin receptors, for example, the integrin β 4 subunit (25). Because *M. leprae* binds to





Fig. 3. α -DG is involved in rLN- α 2G-mediated *M. leprae* adherence to rat Schwann cells. (A) *Mycobacterium leprae* adherence to primary Schwann cells in the presence or the absence of rLN- α 2G. Purified Schwann cells grown for 3 days without forskolin were inoculated with *M. leprae*, which were preincubated with either rLN- α 2G or BSA (10 μ g/ml). The number of *M. leprae*-bound Schwann cells were expressed per 100 cells, and data were presented as mean \pm SD from three experiments. (B) Competitive inhibition of rLN- α 2G-mediated *M. leprae* binding to primary rat Schwann cells by native peripheral nerve α -DG. rLN- α 2G-coated *M. leprae* were preincubated with increasing concentrations of purified α -DG and the mixture was added onto Schwann cells. Cell-bound *M. leprae* were detected by acid-fast labeling and values are presented as percent binding of control (mean \pm SD) obtained from three experiments. Net rLN- α 2G-mediated *M. leprae* adherence to Schwann cells [(*M. leprae*+rLN- α 2G) - (*M. leprae*+BSA)] in the absence of α -DG was considered as 100% binding. (C) α -DG expression on primary Schwann cells purified from rat sciatic nerve as shown by immunofluorescence labeling by mAb IIH6C4 to α -DG. (D) *Mycobacterium leprae* adherence to human Schwann cells in the presence or absence of rLN- α 2G. Experiments were performed in similar conditions as in (A). (E) Competitive inhibition of rLN- α 2G-mediated *M. leprae* binding to human Schwann cells by native α -DG. Experiments were performed in similar conditions as in (B). (F) α -DG expression on immortalized human Schwann cells as detected by immunofluorescence with mAb IIH6C4.



REPORTS

rLN α 2G with high affinity (13) and exogenous rLN α 2G efficiently mediated *M. leprae* adherence to primary Schwann cells (Fig. 3A) (26), this model system enabled us to study the involvement of α -DG in G domain-mediated *M. leprae*-Schwann cell interactions independent of the influence of other regions of the laminin-2 molecule. Using this system in competition assays (27), we found that rLN α 2G-mediated *M. leprae* binding to Schwann cells was inhibited by preincubation of the *M. leprae*+rLN α 2G complex with native α -DG in solution (median inhibitory concentration, IC_{50} = 160 μ g/ml) (Fig. 3B). Fusion proteins of α -DG had no inhibitory effect on *M. leprae* binding to Schwann cells (21). Thus, it is likely that the carbohydrate moieties of α -DG are involved in G domain-mediated *M. leprae* interaction with Schwann cells.

To investigate whether the above findings of α -DG on primary rat Schwann cells holds equally true for humans, we used immortalized human Schwann cells for the binding and invasion assays. Monoclonal antibody (mAb) IIH6C4 (6), which is specific for α -DG, strongly reacted with human Schwann cells (Fig. 3F), and the pattern of α -DG expression on the dorsal surface of the live cells was in the form of microclusters as found for

primary rat Schwann cells (Fig. 3, C and F). Although *M. leprae* alone showed almost no binding to human Schwann cells, preincubation of *M. leprae* with a 10 μ g/ml concentration of rLN α 2G resulted in a >90% increase of cellular adherence (Fig. 3D). This binding was inhibited by preincubation of the *M. leprae*+rLN α 2G complex with native α -DG (IC_{50} = 250 μ g/ml) (Fig. 3E). The low inhibitory effect of α -DG on *M. leprae*+rLN α 2G binding to human Schwann cells, as compared with primary rat Schwann cells (Fig. 3B), may be due to the increased expression of other molecules (or secretory products) on the transformed human Schwann cells that mask the inhibitory effect of native α -DG. Nevertheless, the data suggest the involvement of α -DG in LN α 2G-mediated *M. leprae* adherence in both rat and human Schwann cells. However, our data do not exclude other mechanisms of *M. leprae* adherence of Schwann cells because purified α -DG was unable to compete 100% for rLN α 2G-mediated *M. leprae* adherence, suggesting the participation of other Schwann cell laminin receptors.

To substantiate further the involvement of α -DG as a Schwann cell receptor for *M. leprae*, we examined the effect of rLN α 2G-coated *M. leprae* on the distribution of α -DG

receptors on live primary Schwann cells (28). α -DG is a mobile receptor on muscle cells and forms clusters when it interacts with matrix proteins (28). In control primary Schwann cell cultures, α -DG labeling, as determined by mAb IIH6C4, was equally distributed as microclusters on the dorsal surface of live Schwann cells (immunolabeled as individual dots; Fig. 4A). Microclusters (each dot) on unstimulated Schwann cells may represent single α -DG molecule, as suggested previously for muscle cells (28). When Schwann cells were challenged with rLN α 2G-coated *M. leprae* (after removing free rLN α 2G), α -DG labeling on most of the Schwann cells appeared as aggregates or macroclusters with different sizes and shapes over the entire dorsal cell surface (Fig. 4, B and C). Bacterial challenge caused a dramatic decrease of microclusters and increased the number of macroclusters (compare Fig. 4A with Fig. 4, B and C), which suggests that macroclusters were derived from microclusters. C and D of Fig. 4 demonstrate the colocalization of α -DG receptor clusters and *M. leprae* on the same Schwann cell. Because *M. leprae* alone failed to induce cluster formation (21), the clustering of α -DG on Schwann cells appeared to be contributed by the *M. leprae*-bound LN α 2G. These data strongly suggest that the α -DG participates in the LN α 2G-mediated *M. leprae* interaction with Schwann cells.

Pathogenic bacteria are particularly adapted to exploit host cell functions (1, 2). In peripheral nerves, α -DG appears to link extracellular laminin-2 to the intracellular cytoskeleton through β -DG and associated proteins (6–8, 16). In addition to playing a structural role, this system also regulates host cell functions (6, 7). Present data suggest that *M. leprae* is adapted to exploit this matrix-cytoskeleton link of the peripheral nervous system for its own benefit; *M. leprae* adhere and possibly invade Schwann cells and subsequently interfere with neural cell functions associated with this system.




Fig. 4. α -DG receptor clustering on Schwann cells induced by rLN α 2G-coated *M. leprae*. (A) Immunofluorescence of α -DG showing its distribution on the dorsal surface of live primary Schwann cells before bacterial challenge. The areas corresponding to the nuclei of cells are out of focus. (B and C) Representative examples showing different forms of macroclusters of α -DG 3 hours after bacterial challenge. Comparison of (A) with (B) and (C) reveals that rLN α 2G-coated *M. leprae* induced an extensive aggregation of α -DG receptors on the dorsal surface of Schwann cells. (C and D) Colocalization of α -DG receptor clusters and *M. leprae* on the same Schwann cell as detected by immunodouble labeling with mAbs to α -DG (IIH6C4) (C) and *M. leprae*-specific PGL-1 (D). The arrows mark the α -DG clustering site approximately corresponding with Schwann cell-bound *M. leprae*. All labeling was viewed with a 100 \times oil immersion objective.

References and Notes

1. B. B. Finlay and P. Cossart, *Science* **276**, 718 (1997).
2. S. Falkow, *Cell* **65**, 1099 (1991).
3. G. L. Stoner, *Lancet* **10**, 994 (1979).
4. C. K. Job, *Int. J. Lepr.* **57**, 532 (1989).
5. WHO Weekly Epidemiological Record, 20 September 1995, p. 269.
6. O. Ibraghimov-Beskrovnaia et al., *Nature* **355**, 696 (1992); J. M. Ervasti and K. P. Campbell, *Cell* **66**, 1121 (1991); K. P. Campbell, *ibid.* **80**, 675 (1995).
7. J. M. Ervasti and K. P. Campbell, *J. Cell. Biol.* **122**, 809 (1993); A. Suzuki et al., *Eur. J. Biochem.* **220**, 283 (1994); H. Yamada et al., *J. Neurochem.* **66**, 1518 (1996); B. Yang et al., *J. Biol. Chem.* **270**, 11711 (1995).
8. S. H. Gee, F. Montanaro, M. H. Lindenbaum, S. Carbonetto, *Cell* **77**, 675 (1994); J. T. Campanelli, S. L. Roberds, K. P. Campbell, R. H. Scheller, *ibid.*, p. 663; M. A. Bowe, K. A. Deyst, J. D. Leszyk, J. R. Fallon, *Neuron* **12**, 1173 (1994); J. Sugiyama, D. C. Bowen, Z. W. Hall, *ibid.* **13**, 103 (1994).
9. M. Durbeek et al., *J. Cell. Biol.* **130**, 79 (1995).

10. R. A. Williamson *et al.*, *Hum. Mol. Genet.* **6**, 831 (1997).
11. Y. Sunada, S. M. Bernier, C. A. Kozak, Y. Yamada, K. P. Campbell, *J. Biol. Chem.* **269**, 13729 (1994); H. Xu, X. R. Wu, U. M. Wewer, E. Engvall, *Nature Genet.* **8**, 297 (1994); Y. Sunada, S. M. Bernier, A. Utani, Y. Yamada, K. P. Campbell, *Hum. Mol. Genet.* **4**, 1055 (1995).
12. V. Straub and K. P. Campbell, *Curr. Opin. Neurol.* **10**, 168 (1997).
13. A. Rambukkana, J. L. Salzer, P. D. Yurchenco, E. I. Tuomanen, *Cell* **88**, 811 (1997).
14. I. Leivo and E. Engvall, *Proc. Natl. Acad. Sci. U.S.A.* **85**, 1544 (1988); R. E. Burgeson *et al.*, *Matrix Biol.* **14**, 209 (1994); R. Vuolteenaho *et al.*, *J. Cell Biol.* **124**, 381 (1994).
15. A. M. Mercurio, *Trends Cell Biol.* **5**, 419 (1995).
16. H. Yamada *et al.*, *J. Biol. Chem.* **271**, 23418 (1996).
17. *Mycobacterium leprae* was purified from armadillos and provided by P. J. Brennan (Colorado State University, Fort Collins, CO). Each isolate was tested for acid-fast labeling and *M. leprae*-specific phenolic glycolipid-1 (PGL-1) reactivity with a auramine-rhodamine Bacto TB Fluorescent Stain Kit (Difco, Detroit, MI) and mAb to native PGL-1, respectively, before the assays.
18. The bovine peripheral nerve α -DG was purified as described [(16); H. Yamada, T. Shimizu, T. Tanaka, K. P. Campbell, K. Matsumura, *FEBS Lett.* **352**, 49 (1994); J. M. Ervasti, S. D. Kahl, K. P. Campbell, *J. Biol. Chem.* **266**, 9161 (1991)]. The rabbit skeletal muscle α -DG was purified by using the same method but with KCl-washed heavy microsomes of rabbit skeletal muscle as a starting material. α -DG fusion proteins B and D (FP-B and FP-D) were prepared as described (6).
19. Recombinant (r) LN- α 2G, rLN- α 2(VI-IVb)', and rLN- α 1G fragments were prepared with a baculovirus expression system as previously described [(13); P. D. Yurchenco, U. Sung, M. D. Ward, Y. Yamada, J. J. O'Rear, *J. Biol. Chem.* **268**, 8356 (1993)]. Human merosin (laminin-2 and laminin-4) was a gift from M. Paulsson. The purity of DG preparations and recombinant fragments of laminins were analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting with antibodies specific for each fraction as described (13, 18, 19).
20. *Mycobacterium leprae* binding to α -DG was determined with a solid-phase bacterial adherence assay (13) by using immobilized native α -DG purified from peripheral nerves or skeletal muscles, or recombinant α -DG. Terasaki plates were coated overnight with α -DGs (50 μ g/ml, 0.5 μ g per well) or bovine serum albumin (BSA) as a negative control. *Mycobacterium leprae* (5×10^8 bacteria/ml) suspension was preincubated with rLN α 2G or LN α 2(VI-IVb)' (10 μ g/ml, 0.1 μ g per well) or BSA for 1 hour at 37°C. After blocking the nonspecific binding with BSA, 10 μ l of the *M. leprae* mixture was added to each well and incubated for 1 hour at 37°C. Unbound bacteria were removed by washing with DPBS and wells were fixed with 2.5% glutaraldehyde (Sigma). Adherent *M. leprae* was detected by acid-fast labeling, counted, and expressed as described (13). The effect of heparin and EDTA on rLN α 2G-mediated *M. leprae* binding to α -DG was determined similarly by incubating the bacterial mixture with 10 mM EDTA or heparin (1 mg/ml). The effect of periodate treatment was evaluated by preincubation of increasing concentrations of sodium periodate with native α -DG before the addition of *M. leprae*+rLN α 2G. Periodate and EDTA treatment did not detach the α -DG from wells because no difference was found in antibody activity to α -DG before and after treatment as detected by enzyme-linked immunosorbent assay with polyclonal antibodies to α -DG.
21. A. Rambukkana *et al.*, data not shown.
22. J. M. Ervasti, A. L. Burwell, A. L. Geissler, *J. Biol. Chem.* **272**, 22315 (1997).
23. O. Ibraghimov-Beskrovnyaya *et al.*, *Hum. Mol. Genet.* **2**, 1651 (1993); H. Yamada and K. Matsumura, unpublished data.
24. A. Chiba *et al.*, *J. Biol. Chem.* **272**, 2156 (1997); T. Sasaki, H. Yamada, K. P. Campbell, and T. Endo, unpublished data.
25. Schwann cells were isolated from neonatal rat sciatic

nerve, purified, and amplified as described [S. Einheber, T. A. Milner, F. Giancotti, J. L. Salzer, *J. Cell Biol.* **123**, 1223 (1993)]. Human Schwann cells were purified and immortalized as described [J. L. Rutkowski, J. S. Rhim, K. W. C. Peden, G. I. Tennekoon, in *Neoplastic Transformation in Human Cell Systems in Vitro*, J. S. Rhim and A. Dripschillo, Eds. (Humana, Totowa, NJ, 1991), pp. 343–346]. Schwann cells were plated onto poly-L-lysine-coated eight-well Lab-Tek chamber slides (Nunc) or 12-mm cover slips and cultured without forskolin to prevent the deposition of laminin-2. These primary rat Schwann cells and human Schwann cells were found to be 100% pure as determined by antibody to S-100 antigen.

26. Primary rat Schwann cells and immortalized human Schwann cells were used for both adherence and invasion assays because they are devoid of LN α -2G. The *M. leprae* adherence assay to Schwann cells was previously described (13).

27. For competitive inhibition assays, rLN α 2G-coated *M. leprae* were preincubated with increasing concentrations of native α -DG for 3 hours at 37°C, after which the mixture was added onto Schwann cells, and the adherence assays were performed as described (13). The number of acid-fast-labeled bacteria were quantified, and values were presented as the mean percent binding of controls. The net rLN α 2G-mediated *M. leprae* adherence to Schwann cells was considered as 100%.

28. Light microscopy and immunofluorescence of Schwann cells were performed as described (13). Characterization of mAb IIH6C4 to α -DG and affinity-purified rabbit polyclonal antibody (pAb) to human rLN- α 2G were described previously (7, 13). The mAb F47-21 to native PGL-1 was a gift from A. H. J.

Kolk (Royal Tropical Institute, Amsterdam). The pAb to S-100 was from Sigma. α -DG detection and clustering studies were performed as previously reported [M. W. Cohen, C. Jacobson, P. D. Yurchenco, G. E. Morris, S. Carbonetto, *J. Cell Biol.* **136**, 1047 (1997)]. For bacterial-induced α -DG clustering, *M. leprae* was preincubated with rLN- α 2G for 1 hour at 37°C, and the mixture was centrifuged and the pellet was resuspended in phosphate-buffered saline (PBS) to avoid the contact of free rLN- α 2G with Schwann cells. These rLN- α 2G-coated *M. leprae* were added onto primary Schwann cells as described in adhesion assays. Cultures were then stained live and fixed with 2.5% glutaraldehyde before processing for fluorescence microscopy. In live Schwann cells, α -DG labeling is restricted to the dorsal surface because IIH6C4 immunoglobulin M mAb is unable to reach the ventral cell surface due to its large size. Colocalization of α -DG and *M. leprae* was performed by double immunofluorescence with mAb IIH6 and mAb to *M. leprae* PGL-1.

29. We thank P. J. Brennan for providing *M. leprae* through a National Institute of Allergy and Infectious Disease/NIH contract, M. Zsack for graphics, and S. Terlow for *M. leprae* preparations. We also thank E. Tuomanen for the initial support and encouragement for this study. Supported by grants from the United Nations Development Programme/World Bank/World Health Organization Special Program for Research in Tropical Diseases and NIH (A.R., V.A.F., J.L.S., and P.Y.). H.Y. was supported by an American Heart Association fellowship and by the Mizutani Foundation. K.P.C. is an HHMI investigator.

18 August 1998; accepted 4 November 1998

Identification of α -Dystroglycan as a Receptor for Lymphocytic Choriomeningitis Virus and Lassa Fever Virus

Wei Cao, Michael D. Henry, Persephone Borrow, Hiroki Yamada, John H. Elder, Eugene V. Ravkov, Stuart T. Nichol, Richard W. Compans, Kevin P. Campbell, Michael B. A. Oldstone*

A peripheral membrane protein that is interactive with lymphocytic choriomeningitis virus (LCMV) was purified from cells permissive to infection. Tryptic peptides from this protein were determined to be α -dystroglycan (α -DG). Several strains of LCMV and other arenaviruses, including Lassa fever virus (LFV), Oliveros, and Mobala, bound to purified α -DG protein. Soluble α -DG blocked both LCMV and LFV infection. Cells bearing a null mutation of the gene encoding DG were resistant to LCMV infection, and reconstitution of DG expression in null mutant cells restored susceptibility to LCMV infection. Thus, α -DG is a cellular receptor for both LCMV and LFV.

Arenaviruses consist of several causative agents of fatal human hemorrhagic fevers (1, 2). Among these pathogens, LFV causes an estimated 250,000 cases and more than 5000 deaths annually (1, 3). LCMV, the prototype arenavirus, has been studied primarily in its natural rodent host as a model of viral immunology and pathogenesis (4).

To initiate infection, the LCMV glycoprotein GP-1 anchors the virus to the cell surface through a proteinaceous receptor (5, 6),

which by a virus overlay protein blot assay (VOPBA) (7) was identified as a single high molecular weight glycoprotein (5). The presence of the receptor protein correlated directly with a cell's susceptibility to LCMV attachment and infection (Fig. 1). Its broad migration pattern on SDS-polyacrylamide gels is likely to reflect the heterogeneity in cell type-specific posttranslational modifications (5). In addition to murine cells, a broad range of rodent and primate cells express the same protein (5) (Fig.