

15. A. Bendelac et al., *Science* **268**, 863 (1995); S. Cardell et al., *J. Exp. Med.* **182**, 993 (1995).
16. Splenocytes from naïve and SPZ-primed class II^{-/-} donors were bulk cultured in flat-bottomed plates in the presence or absence of PpGPI and iM4 for 4 days. The proportion of NKT cells expressing intracellular IL-4 was measured by flow cytometry after subtraction of background staining by isotype-matched fluorescein isothiocyanate-conjugated control monoclonal antibody R35-38.
17. D. B. Moody, et al., *Science* **278**, 283 (1997).
18. H. Chen and W. E. Paul, *J. Immunol.* **159**, 2240 (1997); H. Chen, H. Huang, W. E. Paul, *ibid.* **158**, 5112 (1997).
19. S. T. Smiley, M. H. Kaplan, M. J. Grusby, *Science* **275**, 977 (1997).
20. L. Brossay et al., *J. Immunol.* **159**, 1216 (1997); L. Brossay et al., *ibid.* **160**, 3681 (1998).
21. S. H. Park, J. H. Roark, A. Bendelac, *ibid.*, p. 3128.
22. S. A. Porcelli, *Adv. Immunol.* **59**, 1 (1995); A. Bendelac, M. N. Rivera, S.-H. Park, J. H. Roark, *Annu. Rev. Immunol.* **15**, 535 (1997).
23. P. Matzinger, *Annu. Rev. Immunol.* **12**, 991 (1994).
24. M. F. Good, S. Kumar, L. H. Miller, *Immunol. Today* **9**, 351 (1988).
25. Monoclonal antibodies were to CD1.1 (1B1), CD4 (H129.19), CD16 (2.4G2), NK1.1 (PK136), V β 8.1/8.2 (KJ16), V α 14 (CMS-1), I-A^b (M5/114), H-2K^b/H2-D^b (28-8-6), H-2 (M1/42), and IL-4 (11B11, BVD6-24G2, BVD4-1D11).
26. P. J. Romero et al., *Eur. J. Immunol.* **18**, 1951 (1988).
27. M. W. Clarke, R. W. Olafson, T. W. Pearson, *Mol.*

Biochem. Parasitol. **17**, 19 (1985); L. Schofield and F. Hackett, *J. Exp. Med.* **177**, 145 (1993); S. D. Tachado et al., *Proc. Natl. Acad. Sci. U.S.A.* **94**, 4022 (1997).

28. We thank M. Kronenberg for the CD1-transfectants and monoclonal antibody 1B1 and M. Taniguchi for CMS-1. Supported by the United Nations Development Program/World Bank/World Health Organization Special Program for Research and Training in Tropical Diseases, the Australian Academy of Science (L.S.), NIH grant GM 41071 (B.F.-R. and S.C.), INSMED (B.F.-R.), NIH grant AI-40171, and a gift from the Mathers Foundation (M.J.G.). M.J.M. is a Wellcome Trust Senior Research Fellow. M.J.G. is a Scholar of the Leukemia Society of America.

10 August 1998; accepted 3 December 1998

Grassland Vegetation Changes and Nocturnal Global Warming

Richard D. Alward,* James K. Detling, Daniel G. Milchunas

Global minimum temperatures (T_{MIN}) are increasing faster than maximum temperatures, but the ecological consequences of this are largely unexplored. Long-term data sets from the shortgrass steppe were used to identify correlations between T_{MIN} and several vegetation variables. This ecosystem is potentially sensitive to increases in T_{MIN} . Most notably, increased spring T_{MIN} was correlated with decreased net primary production by the dominant C_4 grass (*Bouteloua gracilis*) and with increased abundance and production by exotic and native C_3 forbs. Reductions in *B. gracilis* may make this system more vulnerable to invasion by exotic species and less tolerant of drought and grazing.

There is general consensus that there is an anthropogenic warming signal in the long-term climate record (1). Over land, this is primarily due to average annual minimum temperatures (T_{MIN}) having increased at twice the rate of maximum temperatures (T_{MAX}) (1, 2). At the global scale, these increases in T_{MIN} are related to increases in global cloudiness (1, 3). Experiments with agricultural plants and insect pests suggest important roles for T_{MIN} in influencing plant and insect development (4, 5). However, there has been little research on the consequences of elevated T_{MIN} for natural ecosystems (6, 7). If elevated T_{MIN} leads to longer growing seasons, net primary production and carbon sequestration may increase as a consequence (8). However, the opposite may occur if elevated T_{MIN} leads to increased plant and microbial nocturnal respiration rates without a compensatory

increase in photosynthesis. Additionally, elevated T_{MIN} could shift competitive interactions among C_3 (cool-season) and C_4 (warm-season) plants.

It is important to identify features of ecosystems that are sensitive to changes in T_{MIN} . To date, most modeling efforts and experimental manipulations investigating ecosystem responses to climate change have assumed that future warming will occur primarily during the day or uniformly over the diurnal cycle. This assumption clearly is not valid on a global level nor at most regional scales (2). Furthermore, there is no a priori reason to assume that ecosystems will respond similarly to changes in T_{MIN} and T_{MAX} . To investigate potential ecological consequences of elevated T_{MIN} , we examined a 23-year data set for correlations between temperature [T_{MIN} , T_{MAX} , and mean annual temperature (T_{AVE}) ($T_{\text{AVE}} = (T_{\text{MIN}} + T_{\text{MAX}})/2$)] and both the abundance and aboveground net primary productivity (ANPP) of several key plant species and functional groups found at the Central Plains Experimental Range (9) in northeastern Colorado.

We identified seasonal and annual trends in T_{MIN} and T_{MAX} to determine whether asymmetric diurnal temperature increases held true for this site (10). The densities of most species were determined by counting all individuals within permanently marked quad-

rats (11). Harvests at time of peak standing crop were used as estimates of ANPP (12, 13). Plants in the shortgrass steppe are commonly water-limited, and variation in precipitation could obscure plant responses to gradually changing temperatures (9, 14). Therefore, we included annual and seasonal precipitation totals, in addition to annual and seasonal minimum and maximum temperatures, as variables for stepwise regression model selection (15). We constructed linear models to evaluate significant correlations between these variables and ANPP or plant species density (16).

Mean annual temperatures (T_{AVE}) have increased by an average of $0.12^\circ\text{C year}^{-1}$ at this site since 1964 ($P = 0.0001$, $R^2 = 0.52$). During this period, T_{MAX} increased $0.085^\circ\text{C year}^{-1}$ (Fig. 1A), whereas T_{MIN} increased $0.15^\circ\text{C year}^{-1}$ (Fig. 1B). We limited further analyses of temperature to the period beginning in 1970, when standardized monitoring of vegetation density was initiated. Since 1970, T_{AVE} has risen over 1.3°C , largely due to a significant increase in T_{MIN} of $0.12^\circ\text{C year}^{-1}$ ($P = 0.003$; $R^2 = 0.44$). However, there was no significant trend for T_{MAX} ($P = 0.49$). Averages of seasonal minimum temperatures since 1970 also exhibited significant warming, with similar trends in winter ($0.17^\circ\text{C year}^{-1}$, $P = 0.0013$, $R^2 = 0.40$), spring ($0.16^\circ\text{C year}^{-1}$, $P = 0.0007$, $R^2 = 0.43$), and summer T_{MIN} ($0.12^\circ\text{C year}^{-1}$, $P = 0.004$, $R^2 = 0.33$). No significant trends were detected in fall T_{MIN} ($P = 0.64$, $R^2 = 0.01$). Annual precipitation (Fig. 1C) varied from 230 to 480 mm and has also exhibited a significant linear increase since 1970 (6 mm year^{-1} , $P = 0.007$, $R^2 = 0.30$). However, there were no significant correlations between annual or seasonal T_{MIN} and annual or seasonal precipitation ($P > 0.1$).

Since 1983 (12), ANPP of *Bouteloua gracilis*, the dominant C_4 grass of the shortgrass steppe, declined over time ($-12.2 \text{ g m}^{-2} \text{ year}^{-1}$; $P = 0.002$; $R^2 = 0.78$), and was negatively correlated with average spring T_{MIN} (Fig. 2A). ANPP of the most abundant C_3 forb, *Sphaeralcea coccinea*, was negatively correlated with winter T_{MIN} (Fig. 2B). In contrast, ANPP of

R. D. Alward, Graduate Degree Program in Ecology and Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA. J. K. Detling, Department of Biology and Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA. D. G. Milchunas, Department of Rangeland Ecosystem Science and Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA.

*To whom correspondence should be addressed. Present address: School of Biological Sciences, University of Nebraska, Lincoln, NE 68588, USA. E-mail: ralward1@unl.edu

both the C_3 sedge *Carex eleocharis* (Fig. 2C) and of all C_3 forbs combined (Fig. 2D) was positively correlated with fall and summer T_{MIN} , respectively. Plant density was also correlated with T_{MIN} . Exotic forb density was positively correlated with spring T_{MIN} (Fig. 2E), whereas the density of the C_3 grass *Sitanion hystrix* was positively correlated with winter T_{MIN} (Fig. 2F).

The relationships between T_{MIN} and vegetation revealed by these analyses highlight potential effects of climate change on natural ecosystems. This shortgrass steppe site has experienced increases in T_{MIN} over the past few decades that are similar to trends found by others at larger spatial and temporal scales (2). For each 1°C increase in average spring T_{MIN} , ANPP of the dominant grass decreased by nearly one-third (Fig. 2A). Because this one species (*B. gracilis*) represents an average of 66% of total ANPP and nearly 90% of the total basal cover (9), this result has serious implications for both the structure and function of the shortgrass steppe, if its productivity is causally related to T_{MIN} . *Bouteloua gracilis* is a drought- and grazing-tolerant species that makes up as much as 40% of the diet of cattle on the shortgrass steppe (17). A major reduction in this species could have

substantial consequences for livestock production if it were not replaced by other palatable species. Also of concern is the increase in exotic forb density, because invasive exotic plants are already recognized as a threat to the structure and function of numerous natural ecosystems (18); increasing T_{MIN} may exacerbate this threat.

Elevated T_{MIN} may have direct, but counterbalancing, effects on ANPP and the abundance of plants through mechanisms such as increased rates of carbon assimilation due to warmer mornings, accelerated carbon loss through increased rates of respiration due to warmer nights, and differential effects on C_3 - versus C_4 -photosynthesizing plants. Positive correlations between T_{MIN} and both forb ANPP and exotic plant densities (Fig. 2, D and E) support the hypothesis that increased production will be observed in some plants, and the negative correlation between T_{MIN} and *B. gracilis* ANPP (Fig. 2A) is consistent with the increased respiration hypothesis. In addition to direct effects on rates of plant physiological processes, increases in T_{MIN} could affect plant growth indirectly through changes in the length of growing seasons through increased duration of the frost-free period or changes in the availability of soil water. An increase in season duration would be expected to primarily benefit cool-season plants that are growing most rapidly, and preemptively consuming resources, early and late in the growing season. The positive correlation between spring T_{MIN} and exotic C_3

forbs (Fig. 2E) and between fall T_{MIN} and *C. eleocharis* (Fig. 2C) is consistent with this hypothesis.

Some of the correlations could be the result of effects of elevated T_{MIN} on biotic interactions. Some plants may be increasing (Fig. 2, C through F) in response to the decrease in *B. gracilis* ANPP and the consequent increase in availability of space, nutrients, or water, rather than because of any direct effects of elevated T_{MIN} on these plants. Alternatively, if increased T_{MIN} benefits the growth of C_3 plants (Fig. 2, C through F), this could subsequently result in negative effects on the C_4 *B. gracilis* (Fig. 2A). Such a scenario might occur if cool-season plants were able to reduce available soil moisture before the period of rapid growth of warm-season plants. Intertrophic interactions might also be affected. If increased developmental and consumption rates by insects in response to elevated T_{MIN} (4) are common, increased herbivory could alter plant responses to climate change.

Without a clear causal link, there is no compelling evidence to eliminate factors other than increased T_{MIN} as causes of observed changes in ANPP and plant densities. Unfortunately, most experiments and models designed to investigate climate change effects have focused on manipulating T_{MAX} or have assumed equal contributions by T_{MIN} and T_{MAX} toward achieving an increase in T_{AVE} . The outcomes of such experiments may not realistically predict the future structure and




Fig. 1. Summary of climate data for the Central Plains Experimental Range site. (A) Average annual T_{MAX} . The heavy line is the significant linear trend in T_{MAX} [$T_{\text{MAX}} = -150 + 0.085(\text{year})$; $P = 0.001$; $R^2 = 0.36$]. (B) Average annual T_{MIN} . The heavy line is the significant linear trend in T_{MIN} [$T_{\text{MIN}} = -299 + 0.15(\text{year})$; $P = 3.3 \times 10^{-8}$; $R^2 = 0.68$]. (C) Total annual precipitation. The horizontal dashed line identifies the average annual precipitation (323 mm) at this site since 1939.




Fig. 2. Vegetation correlations with seasonal average T_{MIN} . (A) *Bouteloua gracilis* and spring T_{MIN} [ANPP = $288 - 33.1(T_{\text{MIN}})$; $P = 0.039$; $R^2 = 0.48$]. (B) *Sphaeralcea coccinea* and winter T_{MIN} [ANPP = $0.149 - 0.77(T_{\text{MIN}})$; $P = 0.038$; $R^2 = 0.48$]. (C) *Carex eleocharis* and fall T_{MIN} [ANPP = $13.6 + 2.44(T_{\text{MIN}})$; $P = 0.019$; $R^2 = 0.56$]. (D) Native forb (herbaceous dicots) and summer T_{MIN} [ANPP = $-29.5 + 2.99(T_{\text{MIN}})$; $P = 0.028$; $R^2 = 0.52$]. (E) Exotic (nonnative) forb density and spring T_{MIN} [density = $0.008e^{0.71(T_{\text{MIN}})}$; $P = 0.014$; $R^2 = 0.46$]. (F) *Sitanion hystrix* density and winter T_{MIN} [density = $6.4e^{0.33(T_{\text{MIN}})}$; $P = 0.002$; $R^2 = 0.57$]. Methods for obtaining density and ANPP data are described in (11, 12).

REPORTS

dynamics of ecosystems if climate change continues to be manifested primarily as increases in T_{MIN} . There is a need for experiments that define the relationship between T_{MIN} , plant abundance, and ANPP and that identify mechanisms behind the relationship.

References and Notes


- Intergovernmental Panel on Climate Change (IPCC), *Climate Change 1995: The Science of Climate Change*, J. T. Houghton et al., Eds. (Cambridge Univ. Press, Cambridge, 1995).
- G. Kukla, T. Karl, M. Riches, Eds., *Asymmetric Change of Daily Temperature Range: Proceedings of the International MINIMAX WORKSHOP*, College Park, MD, 27 to 30 September 1993 [U.S. Department of Energy (DOE), Washington, DC, 1994]; T. R. Karl, D. R. Easterling, R. W. Knight, P. Y. Hughes, in *Trends '93: A Compendium of Data on Global Change*, T. A. Boden, D. P. Kaiser, R. J. Sepanski, F. W. Stoss, Eds. (ORNL/CDIAC-65, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN, 1994), pp. 686–736; D. R. Easterling et al., *Science* **277**, 364 (1997).
- A. Henderson-Sellers, *GeoJournal* **27**, 255 (1992); A. Dai, A. D. Del Genio, I. Y. Fung, *Nature* **386**, 665 (1997).
- Y. Yang and N. E. Stamp, *Oecologia* **104**, 225 (1995).
- M. Seddigh, G. D. Jolliffe, J. H. Orf, *Crop Sci.* **29**, 400 (1989); C. A. Zeiher, P. W. Brown, J. C. Silvertooth, N. Matumba, N. Milton, in *Cotton: A College of Agriculture Report*, J. Silvertooth, Ed. (University of Arizona, Tucson, AZ, 1994), pp. 89–96; P. Manunta and M. B. Kirkham, *J. Agron. Crop Sci.* **176**, 267 (1996).
- H. A. Mooney, G. W. Koch, C. B. Field, in *Asymmetric Change of Daily Temperature Range*, G. Kukla, T. Karl, M. Riches, Eds. (DOE, Washington, DC, 1994), pp. 467–484.
- R. A. Kerr, *Science* **255**, 682 (1992); *ibid.* **268**, 1567 (1995); *ibid.* **270**, 1565 (1995).
- M. B. Coughenour and D.-X. Chen, *Ecol. Appl.* **7**, 802 (1997); R. B. Myneni, C. D. Keeling, C. J. Tucker, G. Asrar, R. R. Nemani, *Nature* **386**, 698 (1997).
- For site descriptions, see D. G. Milchunas, W. K. Lauenroth, P. L. Chapman, M. K. Kazempour, *Vegetatio* **80**, 11 (1989); W. K. Lauenroth and D. G. Milchunas, in *Ecosystems of the World*, R. T. Coupland, Ed. (Elsevier, Amsterdam, 1991), vol. 8A, chap. 11.
- Daily T_{MIN} and T_{MAX} were used to calculate monthly averages; monthly data were used to calculate seasonal averages identified as winter (January–March), spring (April–June), summer (July–September) and the previous fall (October–December).
- D. G. Milchunas, W. K. Lauenroth, P. L. Chapman, M. K. Kazempour, *J. Veg. Sci.* **1**, 375 (1990); D. G. Milchunas and W. K. Lauenroth, *Ecology* **73**, 593 (1992); *Ecol. Appl.* **5**, 452 (1995).
- In August or September of each year, all vegetation within 15 0.25-m² circular quadrats was harvested, sorted, and dried. We restricted analyses of ANPP to data collected in 1983–1992, because the method used during these years was different from that used before 1983.
- The following species and functional groups were represented in the analyses: *Artemisia frigida*, *Aristida longiseta*, *Bouteloua gracilis*, *Carex eleocharis*, *Kochia scoparia*, *Opuntia polyacantha*, *Sitanion hystrix*, and *Sphaeralcea coccinea*, warm-season grasses, cool-season grasses and sedges, annual grasses, all grasses and sedges combined, forbs (native herbaceous dicots), exotic (nonnative) forbs, dwarf shrubs, and total biomass from all species. Species richness was also analyzed.
- O. E. Sala, W. J. Parton, L. A. Joyce, W. K. Lauenroth, *Ecology* **69**, 40 (1988); D. G. Milchunas, J. R. Forwood, W. K. Lauenroth, *J. Range Manage.* **47**, 133 (1994).
- The timing of vegetation measurements determined appropriate precipitation variables. Plant density was measured in early July. Although in July, August, and September potential evapotranspiration exceeds precipitation, these dry months would not affect July plant density. Thus, in addition to total annual (January–December) precipitation, "seasonal" precipitation totals included "useful" annual precipitation (October–June) and winter (October–March) and spring (April–June) precipitation. However, ANPP was measured in September; thus the seasonal precipitation totals used in the ANPP analyses included total annual (January–December), useful annual (October–September), winter (October–March), spring (April–June), and summer (July–September) precipitation.
- Significant correlations were identified with a stepwise model-building routine [SAS, version 6.11 (SAS Institute, Cary, NC, 1996); PROC REG with STEPWISE method; enter and exit $\alpha = 0.1$]. Plant density values were log-transformed. We report the single factor that explained the most variation in each species' responses.
- R. W. Rice and M. Vavra, *US IBP Grassland Biome Technical Report No. 103* (Colorado State University, Fort Collins, CO, 1971).
- D. A. Wedin and D. Tilman, *Science* **274**, 1720 (1996); A. N. Cohen and J. T. Carlton, *ibid.* **279**, 555 (1998); T. J. Stohlgren et al., *Ecol. Monogr.*, in press.
- We thank T. J. Minnick for bringing the data sets to our attention and H. W. Polley for comments that improved the manuscript. Support included a NASA Graduate Student Fellowship in Global Change Research to R.D.A., the Shortgrass Steppe Long Term Ecological Research Project (NSF grants DEB-9632852 and BSR-8114822), a National Park Service Agreement (1268-2-9004, CEGR-R92-0043,174), and NSF grant DEB-9708596. The Central Plains Experimental Range is administered by the U.S. Department of Agriculture's Agricultural Research Service.

28 August 1998; accepted 3 November 1998

Enhance your AAAS membership with the Science Online advantage.

Science ONLINE

- Full text Science**—research papers and news articles with hyperlinks from citations to related abstracts in other journals before you receive *Science* in the mail.
- ScienceNOW**—succinct, daily briefings, of the hottest scientific, medical, and technological news.
- Science's Next Wave**—career advice, topical forums, discussion groups, and expanded news written by today's brightest young scientists across the world.
- Research Alerts**—sends you an e-mail alert every time a *Science* research report comes out in the discipline, or by a specific author, citation, or keyword of your choice.
- Science's Professional Network**—lists hundreds of job openings and funding sources worldwide that are quickly and easily searchable by discipline, position, organization, and region.
- Electronic Marketplace**—provides new product information from the world's leading science manufacturers and suppliers, all at a click of your mouse.



All the information you need... in one convenient location.

Visit Science Online at <http://www.scienceonline.org>, call 202-326-6417, or e-mail membership2@aaas.org for more information.

AAAS is also proud to announce site-wide institutional subscriptions to Science Online. Contact your subscription agent or AAAS for details.



AMERICAN ASSOCIATION FOR THE
ADVANCEMENT OF SCIENCE