

$\text{Fe}_8$  complexes have a total spin  $S = 10$ . Below a few kelvin, thermal fluctuations cannot overcome the anisotropy barrier on the scale of hours, and quantum physics sets in. Rather than by rotation, magnetization variation proceeds by transitions across the discrete energy levels approximately described by the magnetic quantum number  $M$ . In a spectacular manifestation of quantum tunneling, the tunnel splitting between  $M = -S$  and  $S - n$  (where  $n$  is an integer between 0 and  $S$ ) oscillates as a function of the applied magnetic field, as a result of interference of two tunneling paths connecting the two quantum levels. Moreover, the oscillation phase depended on the parity of  $n$ , providing solid proof for a quantum mechanic effect.

Magnetic structures slightly larger than the single domain case are most likely to be used in applications in the near future. Claude Chappert proposed calling such structures that are not single domain but have a limited number of degrees of freedom "coherent spin structures." The entire structure of these samples can be computed with micromagnetic theory and compared with experiments. One example is the

prospective basic element for magnetic random access memory (MRAM) (5), a rectangle several hundreds of nanometers in lateral size and a few nanometers thick. The material is magnetically soft (no anisotropy) so that the magnetization lies in the element plane, oriented along the long edge with some deviation at the ends (see figure, right panel). An IBM team recently measured the switching of such an element with nanosecond time resolution by magnetotransport measurements (9). Comparison was made with the predictions of the Landau-Lifshitz-Gilbert equation, which describes the evolution of the local magnetization and contains a relaxation or damping term. Many mechanisms contribute to this damping, such as magnon excitation and intrinsic damping for asymmetric ions. The Gilbert damping parameter  $\alpha$  is therefore often adjusted to experimental values. With some care, it can be extracted from the ferromagnetic resonance (FMR) linewidth, but the bulk FMR value is lower than that extracted from experiments involving an inhomogeneous magnetic structure. For example, Ono *et al.* (2) recently measured the domain wall velocity in a 500 nm by 20 nm

NiFe wire and extracted  $\alpha = 0.6$ , which is rather large. Hopefully, structures with reduced dimensions will help us to understand damping better by restricting the available loss channels. Industry also requires that damping be controlled, because MRAM elements should reverse as fast as possible with no overshoot.

All these experiments on high-spin molecules, nanoparticles, or small elements such as wires, are concerned with magnetization relaxation or damping. The many different approaches to magnetism—classical and quantum, continuous and discrete—can guide each other to understand and use magnetism at dimensions that have only recently become accessible to experiments.

#### References

1. W. Wernsdorfer and R. Sessoli, *Science* **284**, 133 (1999).
2. T. Ono *et al.*, *ibid.*, p. 468.
3. S. A. Majetich and Y. Jin, *ibid.*, p. 470.
4. W. Wernsdorfer *et al.*, *J. Appl. Phys.* **78**, 7192 (1995).
5. W. J. Gallagher *et al.*, *ibid.* **81**, 3741 (1997).
6. S. Guéron, <http://xxx.lanl.gov/abs/cond-mat/9904248>.
7. R. H. Kodama and A. E. Berkowitz, *Phys. Rev. B* **59**, 6321 (1999).
8. W. Wernsdorfer *et al.*, *Phys. Rev. Lett.* **79**, 4014 (1997).
9. R. H. Koch *et al.*, *ibid.* **81**, 4512 (1998).

#### RETROSPECTIVE

## Rolf Landauer (1927–1999)

Charles Bennett

**R**olf Landauer, noted for his contributions to the theory of electrical conductivity, mesoscopic phenomena, and the physics of information, died of brain cancer on 27 April at age 72.

Born into a German Jewish family in Stuttgart in 1927, he emigrated to the U.S. during the 1930s, received his Ph.D. from Harvard in 1950, and began working for IBM in 1952. During his long career at IBM, he was awarded the American Physical Society's Buckley Prize and the Institute of Electrical and Electronics Engineers' Edison Medal and was elected to the U.S. National Academies of Sciences and Engineering, to the American Academy of Arts and Sciences, and to the European Academy of Science.

Rolf Landauer's work on electrical transport theory has found important applications in almost all areas of mesoscopic physics. In a pioneering paper in 1957 and subsequent contributions, he arrived at a simple and elegant formulation for electrical conductance

in terms of scattering. His observation that elastic scattering does not break the coherence of quantum mechanical wave functions led to the idea of normal persistent currents. The Landauer approach now stands along with the Kubo formula as one of the most powerful ways to understand conduction.

During the 1960s, as director of IBM Research's Solid State Sciences Division, Rolf Landauer oversaw the development of the injection laser and promoted large-scale integration, before returning to full-time research in 1969.

Landauer did more than anyone else to establish the physics of information processing as a serious subject for scientific inquiry. This achievement grew out of his deep conviction that "information is physical." He had little use for purely mathematical ideas that could not be given an operational meaning or physical embodiment. This passion led him to pursue the thermodynamics of information and to eventually overturn the long-held belief that each elementary information-processing act requires an expenditure of work at least as great as the mean thermal energy. In his landmark 1961 and subsequent papers, Landauer established that this is true only for operations like erasure that cannot be un-

done, whereas other operations have no intrinsic, irreducible thermodynamic cost. This discovery led to the theory of reversible computers and communication channels, as well as our modern understanding of Maxwell's Demon, whose obedience to the Second Law of Thermodynamics is enforced by what is now called Landauer's principle, the thermodynamic cost of forgetting.

Landauer was critical of portraying each new physical phenomenon as "tomorrow's revolutionary computer technology." Recently, he doubted whether quantum computers could ever be made stable enough for their promised advantages to materialize. Advocates of quantum computation responded with ingenious error correction, eventually earning his grudging respect.

But he was no simplistic opponent of scientific fashion. Recently, he contrasted the narrowness of many recent sessions on quantum mechanical entanglement, with the diversity of the first conference seriously addressing the fundamental physics of computation, held at MIT in 1981: "The narrowing represents real progress; we now understand more about what counts." But, at the same time, the narrowing strengthens fashionability in science. A carefully selected group reinforces its existing values, and declares to science journalists that their stuff is what really counts. I hope that quantum information can receive the attention it deserves, without eclipsing [...] other questions."

The author is at the T. J. Watson Research Center, IBM, Yorktown Heights, NY 10598, USA. E-mail: bennetc@watson.ibm.com. This retrospective was written with the help of Alan Fowler, Joseph Imry, and Seymour Keller.