### REPORTS

achieve a productive and substrate-specific catalytic complex, and this induced-fit mechanism provides a direct opportunity for allosteric regulation. In analogy to an allosteric dimer, the active site of AC is formed at the interface between dyad-related homologous domains. Thus, AC provides two symmetrically related binding sites for the homologous G proteins  $G_s\alpha$  and  $G_i\alpha$  (4, 5). We suggest that  $G_s\alpha$  facilitates collapse of the active-site loops from  $C_{1a}$  and  $C_{2a}$  around the substrate ATP to form an competent active site, whereas  $G_i\alpha$  hinders such collapse and stabilizes an open, inactive conformation of the enzyme.

#### **References and Notes**

- R. K. Sunahara, C. W. Dessauer, A. G. Gilman, Annu. Rev. Pharmacol. Toxicol. 36, 461 (1996); D. M. F. Cooper, Ed., Advances in Second Messenger and Phosphoprotein Research, vol. 32 (Lippincott-Raven, Philadelphia, PA, 1998).
- R. K. Sunahara, C. W. Dessauer, R. E. Whisnant, C. Kleuss, A. G. Gilman, J. Biol. Chem. 272, 22265 (1997).
- R. E. Whisnant, A. G. Gilman, C. W. Dessauer, *Proc. Natl. Acad. Sci. U.S.A.* **93**, 6621 (1996); S.-Z. Yan, D. Hahn, Z.-H. Huang, W.-J. Tang, *J. Biol. Chem.* **271**, 10941 (1996).
- 4. C. W. Dessauer, J. J. Tesmer, S. R. Sprang, A. G. Gilman, J. Biol. Chem. **273**, 25831 (1998).
- J. J. G. T. Tesmer, R. K. Sunahara, A. G. Gilman, S. R. Sprang, *Science* 278, 1907 (1997).
- D. L. Garbers and R. A. Johnson, J. Biol. Chem. 250, 8449 (1975); G. Zimmermann, D. Zhou, R. Taussig, *ibid.* 273, 19650 (1998).
- P. J. Artymiuk, A. R. Poirrette, D. W. Rice, P. Willett, Nature 388, 33 (1997).
- 8. S. Doublié, S. Tabor, A. M. Long, C. C. Richardson, T. Ellenberger, *ibid.* **391**, 251 (1998).
- 9. H. Huang, R. Chopra, G. L. Verdine, S. C. Harrison, Science 282, 1669 (1998).
- Y. Li, S. Korolev, G. Waksman, EMBO J. 17, 7514 (1998).
- N. Kaushik et al., Biochemistry **35**, 11536 (1996); A. Y. Woody, S. S. Eaton, P. A. Osumi-Davis, R. W. Woody, *ibid.*, p. 144.
- I. Shoshani, V. Boudou, C. Pierra, G. Gosselin, R. A. Johnson, unpublished data.
- 13. R. K. Sunahava and J. J. G. Tesmer, data not shown. 14. F. Eckstein, P. Romaniuk, W. Heideman, D. Storm,
- J. Biol. Chem. 256, 9118 (1981). 15. The recombinant C<sub>1a</sub> domain from canine type V AC, the  $C_{za}$  domain from rat type II AC, and bovine  $G_{sa}$ were expressed in E. coli and purified as previously described (30).  $G_{s\alpha}$  was activated with  $Mg^{2+} \cdot GTP\gamma S$ and then subjected to limited proteolysis by trypsin. Crystals of the ternary complex between  $C_{1a}$ ,  $C_{2a}$ , and  $G_{s\alpha}$  were grown as previously described (5). All crystals were harvested in cryoprotectant (5) to which was added various combinations of inhibitors and metal ions, described as follows: for AC  $\cdot \beta$ LddATP  $\cdot Mg$ , 5 mM MgCl<sub>2</sub> and 380 μM β-L-2',3'-dd-5'-ATP; for AC  $\cdot \beta$ LddATP  $\cdot$  Mn, 250  $\mu$ M MgCl<sub>2</sub>, 500  $\mu$ M MnCl<sub>2</sub>, and 380  $\mu$ M  $\beta$ -L-2',3'-dd-5'-ATP; for AC  $\cdot \beta$ Ldand 380  $\mu$ M g-L-2 ,3 - dd-5 -ATP; for AC +  $\beta$ Ld-dATP + Zn, 250  $\mu$ M MgCl<sub>2</sub>, 500  $\mu$ M ZnCl<sub>2</sub>, and 380  $\mu$ M  $\beta$ -L-2',3'-dd-5'-ATP; for AC + ATP $\alpha$ S-R<sub>p</sub> + Mn, 500  $\mu$ M MgCl<sub>2</sub>, 500  $\mu$ M MnCl<sub>2</sub>, and 1 mM ATP $\alpha$ S-R<sub>p</sub>. Crystals of AC +  $\beta$ LddATP + Mg, AC +  $\beta$ LddATP + Mn, and AC +  $\beta$ LddATP + Zn were soaked for 1 to 2 hours; the conformation of AC in these complexes was closed. AC  $\cdot$  ATP $\alpha$ S-R<sub>p</sub>  $\cdot$  Mn was soaked for 6 hours to achieve a closed conformation; 2-hour soaks resulted in structures of the open conformation of AC. AC complexes with inhibitors such as ATP  $\cdot$  Zn<sup>2+</sup>, ATP $\alpha$ S-S $_{P}$   $\cdot$  Zn<sup>2+</sup>, and 2'-iodo-ATP  $\cdot$ Mg<sup>2+</sup> also adopt the open conformation, and each metal site exhibits the same preference for either Mn<sup>2+</sup> or Zn<sup>2+</sup> as observed in the closed conformation (13). Strong electron density is observed for the  $\beta$  and  $\gamma$  phosphates of these inhibitors in

the presence of  $Mg^{2+}$  alone or of  $Mn^{2+}$  and  $Mg^{2+}$ , and for the entire triphosphate in the presence of  $Zn^{2+}$  and  $Mg^{2+}$ . Much weaker density corresponding to a purine ring is observed in the purine binding pocket of these complexes. However, the purine and triphosphate densities are not continuous, and a single model of ATP cannot be built to accommodate both purine and triphosphate binding sites.

- E. E. Kim and H. W. Wyckoff, J. Mol. Biol. 218, 449 (1991).
- 17. V. Derbyshire, J. K. Pinsonneault, C. M. Joyce, *Methods Enzymol.* **262**, 363 (1995).
- J. A. Piccirilli, J. S. Vyle, M. H. Caruthers, T. R. Cech, *Nature* **361**, 85 (1993).
- H. Pelletier, M. R. Sawaya, W. Wolfle, S. H. Wilson, J. Kraut, *Biochemistry* **35**, 12762 (1996); I. Slaby, B. Lind, A. Holmgren, *Biochem. Biophys. Res. Commun.* **122**, 1410 (1984); G. C. King, C. T. Martin, T. T. Pham, J. E. Coleman, *Biochemistry* **25**, 36 (1986); E. Ferrari *et al., J. Virol.* **73**, 1649 (1999).
- 20. S.-Z. Yan, Z.-H. Huang, R. S. Shaw, W.-J. Tang, J. Biol. Chem. 272, 12342 (1997).
- Y. Liu, A. E. Ruoho, V. D. Rao, J. H. Hurley, Proc. Natl. Acad. Sci. U.S.A. 94, 13414 (1997).
- W. S. Sheldrick and E. Rieke, Acta Crystallogr. B34, 2324 (1978); K. I. Varughese, C. T. Lu, G. Kartha, J. Am. Chem. Soc. 104, 3398 (1982); N. Padmaja, S. R. Ramakumar, M. A. Viswamitra, Bull. Chem. Soc. Jpn. 64, 1359 (1991).
- 23. C. W. Dessauer and A. G. Gilman, J. Biol. Chem. 272, 27787 (1997).
- W.-J. Tang, M. Stanzel, A. G. Gilman, *Biochemistry* 34, 14563 (1995).
- 25. J. J. Tesmer and S. R. Sprang, *Curr. Opin. Struct. Biol.* **8**, 713 (1998).
- L. S. Beese and T. A. Steitz, *EMBO J.* **10**, 25 (1991);
  T. A. Steitz, *Curr. Opin. Struct. Biol.* **3**, 31 (1993); H. Pelletier, M. R. Sawaya, A. Kumar, S. H. Wilson, J. Kraut, *Science* **264**, 1891 (1994).

- 27. S. Doublié, M. R. Sawaya, T. Ellenberger, *Structure* 7, R31 (1999).
- S. Doublié and T. Ellenberger, *Curr. Opin. Struct. Biol.* 8, 704 (1998).
- S. G. Sarafianos, V. N. Pandey, N. Kaushik, M. J. Modak, J. Biol. Chem. 270, 19729 (1995); V. N. Pandey, N. Kaushik, M. J. Modak, *ibid.* 269, 13259 (1994).
- 30. R. K. Sunahara *et al., J. Biol. Chem.* **273**, 16332 (1998).
- 31. Single-letter abbreviations for the amino acid residues are as follows: D, Asp; I, Ile; K, Lys; N, Asn; R, Arg,
- 32. A. G. W. Leslie, MOSFLM 5.50 for image plate data (1997).
- 33. S. Bailey, Acta Crystallogr. D50, 760 (1994).
- 34. A. T. Brünger et al., ibid. D54, 905 (1998).
- T. A. Jones, J. Y. Zou, S. W. Cowan, M. Kjeldgaard, *ibid*. A47, 110 (1991).
- 36. We thank F. Eckstein at the Max-Planck Institute (Göttingen) for pure ATP $\alpha$ S-R<sub>p</sub>; J. Collins for technical assistance; C. Brautigam for helpful discussion and for reading the manuscript; D. Coleman and the MacCHESS staff for their assistance with data collection at the Cornell High Energy Synchrotron Source (CHESS); and L. Esser for his assistance in preparing figures. R.K.S. was supported by a postdoctoral fellowship from the Medical Research Council of Canada. This work was supported by grants from the Agence Nationale de Recherche sur le SIDA (ANRS, France) to G.G.; by NIH grants DK38828 (R.A.J.), GM34497 (A.G.G.), and DK46371 (S.R.S.); by an Innovative Technology Grant from the Stony Brook Center for Biotechnology to R.A.J.; by Welch Foundation grants I-1271 (A.G.G.) and I-1229 (S.R.S.); and by the Raymond and Ellen Willie Distinguished Chair of Molecular Neuropharmacology (A.G.G.). Coordinates for the four models have been deposited in the Protein Data Bank with the codes 1CJT, 1CJU, 1CJV, and 1CJK.

26 April 1999; accepted 22 June 1999

# Staphylococcus aureus Sortase, an Enzyme that Anchors Surface Proteins to the Cell Wall

## Sarkis K. Mazmanian, Gwen Liu, Hung Ton-That, Olaf Schneewind\*

Surface proteins of Gram-positive bacteria are linked to the bacterial cell wall by a mechanism that involves cleavage of a conserved Leu-Pro-X-Thr-Gly (LPXTG) motif and that occurs during assembly of the peptidoglycan cell wall. A *Staphylococcus aureus* mutant defective in the anchoring of surface proteins was isolated and shown to carry a mutation in the *srtA* gene. Overexpression of *srtA* increased the rate of surface protein anchoring, and homologs of *srtA* were found in other pathogenic Gram-positive bacteria. The protein specified by *srtA*, sortase, may be a useful target for the development of new antimicrobial drugs.

Hospital isolates of *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Enterococcus faecalis* have become resistant to most, if not all, known therapeutic regimens (1). Many antibiotics, including penicillin and its derivatives, target the transpeptidation reaction of bacterial cell wall synthesis, which cross-links peptidoglycan strands (2). To search for other cell wall synthesis reactions that may serve as targets for antimicrobial therapy, we have focused on the anchoring of surface proteins to the peptidoglycan of Gram-positive bacteria.

Department of Microbiology and Immunology, UCLA School of Medicine, University of California, 10833 Le Conte Avenue, Los Angeles, CA 90095, USA.

<sup>\*</sup>To whom correspondence should be addressed. Email: olafs@ucla.edu

Surface proteins not only promote interaction between the invading pathogen and animal tissues, but also provide ingenious strategies for bacterial escape from the host's immune response (3). In the case of *S. aureus* protein A, immunoglobulins are captured on the microbial surface and camouflage bacteria during the invasion of host tissues (4). Protein A is cleaved by a transpeptidase, sortase, between the threonine and the glycine of a conserved LPXTG motif (5). The carboxyl group of threonine

Fig. 1. Isolation of a staphylococcal mutant defective in cell wall sorting of surface proteins. (A) Primary structure of the surface protein precursor SEB-SPA<sub>490-524</sub>, a fusion between enterotoxin B (SEB) and COOH-terminal protein A (SPA) sequences. P1 is directed across the cytoplasmic membrane by an NH2-terminal leader peptide, and is then cleaved to generate P2. P2 bears a COOH-terminal sorting signal that includes an LPXTG motif, a hydrophobic domain (black bar), and a positively charged tail (boxed +). The sorting signal of P2 is cleaved at the LPXTG motif, and the mature protein (M) is linked to the cell wall. (B) Staphylococcus aureus ts mutants were screened for the accumulation of P2 by labeling with [35S]Met. SM317 and SM329 accumulate more P2 than does the wild-type (WT) strain OS2. (C) Pulse-chase analysis of SEB-SPA490-524 anchoring in wild-type and mutant strains.

Fig. 2. Structure of surface protein anchor in strain SM317. (A) Primary structure of SEB-MH<sub>6</sub>-CWS and its linkage to the cell wall. The glycan strands of the staphylococcal cell wall consist of a repeating disaccharide Nacetylmuramic acid- $(\beta 1-4)-N$ -acetylglucosamine (MN-GN) in which the lactyl of muramic acid is linked to the wall peptide (L-Ala-D-iGln-L-Lys-D-Ala). Wall peptides are linked to the pentaglycine cross-bridge, which tethers the  $\varepsilon$ -amino of L-Lys to the carboxyl of D-Ala. The cell wall can be cut at specific sites with the enzymes muramidase (solid arrow) and  $\phi$ 11 hydrolase (open arrows). (B) SEB-MH<sub>6</sub>-CWS was solubilized by digesting the staphylococcal cell wall with muramidase (mutanolysin) and purified by affinity chromatography on Ni-NTA resin. The polypeptide was cleaved with cyanogen bromide (CnBr), and COOHterminal anchor peptides were purified by another affinity chromatography step. The structure of the mutanolysinreleased anchor peptides of strain SM317 was analyzed by MALDI-MS (6). (C) Muramidase-released anchor peptides were digested with  $\phi$ 11 hydrolase and analyzed by MALDI-MS. The observed ions represent anchor peptide linked to cell wall tetrapeptide (m/z)2236) and murein-disaccharide tetrapeptide (m/z 2715 and 2757).

## REPORTS

is amide-linked to the amino group of the pentaglycine cross-bridge, thereby tethering the COOH-terminal end of protein A to the bacterial cell wall ( $\delta$ ). This reaction, called cell wall sorting, is strikingly similar to the penicillin-sensitive transpeptidation reaction, and is likely to occur in most Grampositive bacteria (7).

To identify cell wall sorting mutants, we mutagenized *S. aureus* strain OS2 with nitrosoguanidine ( $\delta$ ). Temperature-sensitive (ts) mutants were identified and 1000 were trans-





formed with pSEB-SPA490-524, a plasmid encoding a reporter protein that allows measurement of surface protein anchoring (9). The SEB-SPA<sub>490-524</sub> precursor (P1) is exported from the cytoplasm, and its NH2-terminal leader peptide is removed to generate the P2 intermediate (Fig. 1A) (10). P2 is cleaved by sortase at the LPXTG motif to generate the mature, surface-anchored protein (M). After labeling with [35S]Met for 5 min, the reporter protein in strain OS2 was distributed as follows: P1 (5%), P2 (19%), and M (76%) (Fig. 1B) (11). We used this assay to screen 1000 ts mutants and identified two strains in which there was aberrant accumulation of P2 (47% in SM317 and 26% in SM329) (Fig. 1B). Pulse-chase analysis (12) revealed that in strain OS2, P2 was cleaved and anchored within 2 min, whereas in strain SM317 these events required more than 10 min (Fig. 1C). In strain SM329, P2 processing required 3 min, suggesting a mild defect in cell wall sorting (13).

Previous work showed that mutations in *S. aureus fem* genes slow the anchoring of surface proteins to the cell wall (14). These genes are thought to specify enzymes that catalyze the addition of glycines to the  $\varepsilon$ -amino of lysine within the peptidoglycan precursor lipid II (15). To examine whether the SM317 and SM329 mutants were defective in cell wall synthesis, we tested their sensitivity to lysostaphin, an enzyme that cuts the pentaglycine cross-bridges of the staphy-



**Fig. 3.** Overexpression of *srtA* reduces P2 accumulation in wild-type *S. aureus* strain OS2 and in two mutant strains, SM317 and SM329. (**A**) Transformants of a multicopy plasmid library in strain SM317 were screened by labeling with [<sup>35</sup>S]Met for a decrease in the accumulation of P2. Plasmid pGL1834 contains the *srtA* gene cloned into pC194-mcs. (**B**) The *srtA* gene of strain SM317 (pGL1898) or of strain OS2 (pGL1897) was transformed into SM317 and analyzed for P2 processing. (**C**) All three strains were transformed with pGL1834 and subjected to pulse-chase analysis (*12*).

lococcal cell wall (16). In contrast to fem mutants, which are resistant to lysostaphin (17), strains SM317 and SM329 were sensitive at concentrations that also inhibited growth of wild-type staphylococci, indicating that their sorting defects are not caused by a mutationally altered cell wall cross-bridge (18). To measure cell wall synthesis, we grew the wild-type and mutant strains in minimal medium containing [3H]Lys or [3H]Leu. Because lysine (but not leucine) is a component of the cell wall, the ratio of [<sup>3</sup>H]Lys:[<sup>3</sup>H]Leu incorporation into acid-precipitable and protease-resistant murein polymer is a measure of cell wall synthesis (19). Wild-type S. aureus displayed a ratio of 30, and the inhibition of cell wall synthesis by vancomycin reduced this ratio to 1.5. Strains SM317 and SM329 displayed ratios of 18 and 19, respectively, which indicates that the accumulation of P2 in strain SM317 is not caused by a defect in cell wall synthesis.

To determine the cell wall anchor structure of surface proteins in strain SM317, we introduced into cells a plasmid (pHTT4) specifying the reporter protein SEB-MH<sub>e</sub>-CWS (Fig. 2A) (6). The cell wall was purified and digested with mutanolysin, which hydrolyzes the glycan strands (6). Mutanolysin-released surface protein was purified by chromatography on nickelnitrilotriacetic acid (Ni-NTA) and cleaved at Met with cyanogen bromide. COOH-terminal peptides bearing cell wall anchor structures were purified by a second affinity chromatography step and analyzed by matrix-assisted laser desorption ionization mass spectrometry (MALDI-MS) (Fig. 2). A series of ion signals with regularly spaced mass increments was revealed. These measurements are consistent with increasing numbers of peptidoglycan subunits linked to the COOH-terminal threonine of surface protein (20). If surface protein is tethered to cross-linked peptidoglycan of strain SM317, digestion of muramidase-solubilized anchor peptides with φ11 hydrolase should produce anchor peptide linked to murein tetrapeptide and disaccharide-tetrapeptide (21) (Fig. 2). This was tested, and the doubly digested anchor peptides generated the predicted ion signals (Fig. 2C). Thus, surface proteins of S. aureus SM317 are tethered to cross-linked peptidoglycan in a manner that is indistinguishable from the anchor structure of polypeptides in wild-type staphylococci.

**Fig. 4.** Deduced amino acid sequence of the *srtA* gene (25). The  $NH_2$ -terminal hydrophobic membrane anchor sequence is boxed. The single Cys is shaded. The *srtA* gene of strain SM317 carries two mutations, one in codon 35 replacing Asp (GAT) with Gly (GGT), and another in codon 180 replacing Thr (ACA) with Ala (AGA). The altered amino acids are

These results suggest that the accumulation of P2 in strain SM317 is caused by a defect in sortase.

We reasoned that overexpression of sortase from a multicopy plasmid should reduce the concentration of P2 in both wild-type and mutant S. aureus. A plasmid library of 2000 random DNA fragments from S. aureus OS2 was screened for sequences that reduce the accumulation of P2 in strain SM317 and two plasmids, pGL1631 and pGL1834, were identified (Fig. 3) (22). Transformation with pGL1834 reduced the P2 concentration in strain SM317 by 35%, in strain SM329 by 14%, and in wild-type S. aureus OS2 by 9%. All three strains showed a rapid increase in P2 processing (Fig. 3C). We mapped the critical S. aureus sequences in pGL1631 and pGL1834 to a gene that we named srtA (surface protein sorting A) (18, 23).

The *srtA* gene specifies a protein of 206 amino acids with a potential NH<sub>2</sub>-terminal signal peptide/membrane anchor sequence and a presumed active-site cysteine at position 184, consistent with the observation that the cell wall sorting reaction is sensitive to reagents that modify sulfhydryl groups (Fig. 4) (10). Database searches revealed that *srtA* homologs are present in *Actinomyces naeslundii, Bacillus subtilis, Enterococcus faecalis, Staphylococcus aureus, Streptococcus mutans, Streptococcus pneumoniae*, and *Streptococcus pyogenes*. All *srtA* homologs displayed absolute conservation of the cysteine-encoding codon 184 (18).

To examine whether the defect in cell wall sorting of S. aureus SM317 is caused by a srtA mutation, we amplified the genes from S. aureus OS2 and SM317 by PCR, and cloned them into a multicopy vector, which was then transformed into S. aureus SM317 (18, 23). The wild-type (OS2) srtA gene reduced the accumulation of P2 by 32%, whereas the mutant had no effect (Fig. 3B). DNA sequence analysis revealed mutations in codons 35 and 180 in the SM317 srtA gene. Interestingly, multicopy expression of wild-type srtA (pGL1894) did not fully complement the ts growth phenotype of SM317. The transformed mutant grew at 42°C but slower than the wild type, suggesting that the conditional lethal phenotype of S. aureus SM317 is not caused solely by the mutations in srtA.

Together, our data reveal that *srtA* encodes sortase, the transpeptidase that anchors surface proteins to the bacterial cell wall

MKKWTNRI<u>MT IAGVVLILVA AYLFA</u>KPHID 30 NYLHDKDKDE KIEQYDKNVK EQASKDKKQQ 60 AKPQIPKDKS KVAGYIEIPD ADIKEPVYPG 90 PATPEQLNRG VSFAEENESL DDQNISIAGH 120 TFIDRPNYQF TNLKAAKKGS MVYFKVGNET 150 RKYKMTSIRD VKPTDVGVLD EQKGKDKQLT 180 LITCDDYNEK TGVWEKRKIF VATEVK\* 206

indicated in bold. The DNA sequence of plasmid pGL1834 has been submitted to GenBank (accession number AF162687).

(24). In principle, purified SrtA protein can be used to screen for compounds that inhibit cell wall sorting, a strategy that may lead to new therapies for human infections caused by Gram-positive bacteria.

### **References and Notes**

- H. S. Gold and R. C. Moellering, N. Engl. J. Med. 335, 1445 (1996); K. Hiramatsu et al., Lancet 350, 1670 (1997); K. Sieradzki, R. B. Roberts, S. W. Haber, A. Tomasz, N. Engl. J. Med. 340, 517 (1999).
- J. L. Strominger, Johns Hopkins Med. J. 133, 63 (1973); C. T. Walsh, Science 261, 308 (1993).
- 3. W. W. Navarre and O. Schneewind, *Microbiol. Mol. Biol. Rev.* **63**, 174 (1999).
- M. Uhlén, B. Guss, B. Nilsson, F. Götz, M. Lindberg, J. Bacteriol. 159, 713 (1984).
- 5. W. W. Navarre and O. Schneewind, *Mol. Microbiol.* **14**, 115 (1994).
- H. Ton-That, K. F. Faull, O. Schneewind, J. Biol. Chem. 272, 22285 (1997); W. W. Navarre, H. Ton-That, K. F. Faull, O. Schneewind, *ibid*, 273, 29135 (1998).
- O. Schneewind, A. Fowler, K. F. Faull, Science 268, 103 (1995).
- 8. Staphylococci (10<sup>12</sup> colony-forming units) were treated with N-methyl-N'-nitro-N-nitrosoguanidine (0.2 mg/ml) for 45 min at 30°C, and mutagenesis was quenched by the addition of 2 volumes of 100 mM sodium phosphate (pH 7.0). About 80% of the mutagenized population was killed, and the mutational frequency of rifampicin-resistant *rpoB* mutations was increased to  $1.2 \times 10^{-4}$ . Ts mutants were selected three times in succession by growing the mutagenized population in tryptic soy broth (TSB) at 42°C and treating with penicillin G (8  $\mu$ g/ml) for 2 hours. Colonies were formed at 30°C, streaked on TSB agar, and examined for growth at 42°C.
- O. Schneewind, D. Mihaylova-Petkov, P. Model, *EMBO* J. 12, 4803 (1993).
- 10. H. Ton-That and O. Schneewind, J. Biol. Chem., in press.
- 11. Staphylococcus aureus was grown in TSB supplemented with chloramphenicol (cm; 10 µg/ml) or tetracycline (tet; 2 µg/ml) at 30°C until the culture reached an optical density at 660 nm of 0.6. Cells were incubated at 42°C for 20 min, centrifuged at 15,000g for 3 min, and washed with 1 ml of prewarmed minimal medium (12). Cells were labeled with 50 µCi of [35S]-Promix (Amersham) for 5 min, and surface protein processing was guenched by addition of 75 µl of 100% trichloroacetic acid (TCA). TCA precipitates were collected by centrifugation, washed in acetone, and dried. Samples were suspended in 1 ml of 0.5 M tris-HCl (pH 7.5) and incubated for 1 hour at 37°C after addition of 50 µl of lysostaphin (2 mg/ml). Proteins were again precipitated with TCA and washed with acetone; after immunoprecipitation with  $\alpha$ -enterotoxin B, they were analyzed by 14% SDS-polyacrylamide gel electrophoresis and Phosphorlmager.
- 12. O. Schneewind, P. Model, V. A. Fischetti, *Cell* **70**, 267 (1992).
- 13. In pulse-labeling experiments, strain SM329 had a more severe defect in cell wall sorting when grown in minimal medium rather than TSB. The reason for this phenomenon is not known.
- H. Ton-That, H. Labischinski, B. Berger-Bächi, O. Schneewind, J. Biol. Chem. 273, 29143 (1998).
- 15. B. Berger-Bächi, Trends Microbiol. 2, 389 (1994).
- C. A. Schindler and V. T. Schuhardt, Proc. Natl. Acad. Sci. U.S.A. 51, 414 (1964).
- U. Kopp, M. Roos, J. Wecke, H. Labischinski, Microb. Drug Resist. 2, 29 (1996).
- Lysostaphin sensitivity data, mapping data, and the sequence alignment can be viewed at *Science* Online (www.sciencemag.org/feature/data/1041555.shl).
- D. Boothby, L. Daneo-Moore, G. D. Shockman, *Anal. Biochem.* 44, 645 (1971).
- Ion signals of mutanolysin-solubilized anchor peptides correspond to H<sub>6</sub>AQALPET-Gly<sub>5</sub> linked to cell wall tetrapeptide (predicted mass 2235; observed 2236), pentapeptide (predicted mass 2306; observed 2307), N,O6-diacetyl-MurNac-GlcNac tetrapeptide

(predicted mass 2755, observed 2755), N,O6-diacetyl-MurNac-GlcNac pentapeptide (predicted mass 2828, observed 2826), murein-tetrapeptide-mureinpentapeptide (predicted mass 3990, observed 3995), (murein-tetrapeptide)<sub>2</sub>-murein-pentapeptide (predicted mass 5194; observed 5196), and (mureintetrapeptide)<sub>4</sub> (predicted mass 6285, observed 6286).

- W. W. Navarre, H. Ton-That, K. F. Faull, O. Schneewind, J. Biol. Chem. 274, 15847 (1999).
- 22. Plasmid pGL4 contains the coding sequence of SEB-SPA<sub>490-524</sub>, which was released from pSEB-SPA<sub>490-524</sub> by Eco RI–Bam HI digestion and inserted into the pT181 derivative pWil5. S. aureus SM317 (pGL4) was grown on TSB tet agar. A plasmid library of S. aureus OS2 chromosomal DNA was obtained by partial digestion with Sau 3A1. DNA fragments of 3 to 5 kb were purified and

cloned into Bam HI–digested pC194-mcs, which contains the multiple cloning site of pUC19 inserted into the Hind III site of pC194. SM317 (pGL4) was transformed with the pC194-mcs plasmid library and transformants were selected on TSB tet-cm agar.

- 23. The DNA insertions of pGL1631 and 1834 were mapped and sequenced by synthesizing oligonucleotide primers. The primers for the amplification of srtA from the chromosomal DNA of S. aureus strains OS2 (pGL1897) and SM317 (pGL1898) were 5'-AAG-GATCCAAAAGGACCGGTATACATTGC-3' and 5'-AAGGATCCTACCTTTTCCTCTAGCTGAAG-3'.
- 24. In another report we show that purified SrtA protein catalyzes the in vitro transpeptidation of substrates bearing an LPXTG motif (H. Ton-That, G. Liu, S. K. Mazmanian, K. F. Faull, O. Schneewind, in preparation).

# Phosphorylation and Sequestration of Serotonin Transporters Differentially Modulated by Psychostimulants

## Sammanda Ramamoorthy and Randy D. Blakely\*

Many psychotropic drugs interfere with the reuptake of dopamine, norepinephrine, and serotonin. Transport capacity is regulated by kinase-linked pathways, particularly those involving protein kinase C. (PKC), resulting in transporter phosphorylation and sequestration. Phosphorylation and sequestration of the serotonin transporter (SERT) were substantially impacted by ligand occupancy. Ligands that can permeate the transporter, such as serotonin or the amphetamines, prevented PKC-dependent SERT phosphorylation. Nontransported SERT antagonists such as cocaine and antidepressants were permissive for SERT phosphorylation but blocked serotonin effects. PKC-dependent SERT sequestration was also blocked by serotonin. These findings reveal activitydependent modulation of neurotransmitter reuptake and identify previously unknown consequences of amphetamine, cocaine, and antidepressant action.

sensitizes monoaminergic synapses to subsequent psychostimulant challenge (9), which may involve modulated protein kinase cascades (10). Alterations in SERT activity and binding site density (11) and SERT gene polymorphisms (12) have implicated the transporter in anxiety, depression, suicide, autism, and substance abuse. Recent findings with transgenic mice (13) support an important role for 5-HT and SERTs in the behavioral actions of cocaine and amphetamine.

SERT expression can be rapidly modulated by receptor stimulation, second messenger production, and kinase activation (14-16). Suppression of SERT activity accompanying protein kinase C (PKC) activation (17) arises from a loss of 5-HT uptake capacity (V<sub>max</sub>). The loss in 5-HT uptake capacity correlates with a loss of surface-expressed SERTs (17), similar to the PKC modulation of homologous  $\gamma$ -aminobutyric acid (GABA), DA, and NE transporters (18). PKC activators and phosphatase inhibitors induce SERT phosphorylation (19) with a similar time course and kinase antagonist sensitivity as observed for changes in 5-HT transport.

- 25. Abbreviations for the amino acid residues are as follows: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr.
- 26. We thank W. W. Navarre for help in determining the surface protein anchor structure of strain SM317, and D. Missiakas, P. Model, M. Russel, and members of our laboratory for critical reading of this manuscript. S.K.M. was supported by the Predoctoral Training Program in Genetic Mechanisms at UCLA (T32GM07104). H.T.-T. was supported by the Microbial Pathogenesis Training Grant at UCLA (AI 07323). Work in O.S.'s laboratory is supported by grant AI33987 from the National Institute of Allergy and Infectious Diseases.

6 May 1999; accepted 28 June 1999

We investigated whether the regulation of SERTs was influenced by transport and whether SERT ligands differentially influenced SERT regulation. Figure 1A shows that PKCmediated SERT phosphorylation in transfected human embryonic kidney-293 (HEK-293) cells was substantially diminished if assayed in the presence of the transported neurotransmitter, 5-HT. SDS-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of immunoprecipitates from  $[^{32}P]PO_{4}$ -labeled cell extracts (20) revealed a three- to fivefold stimulation of human SERT (hSERT) phosphorylation after phorbol 12-myristate 13-acetate (β-PMA) application. This stimulation was abolished by coapplication of PKC antagonists. In the presence of 5-HT (1 µM), SERT phosphorylation triggered by phorbol esters was also substantially blunted. At low concentrations of B-PMA (for example, 10 nM), 5-HT essentially abolished phorbol ester-induced SERT labeling. At 200 nM β-PMA, where labeling of SERTs is maximal, we consistently achieved 40 to 60% inhibition of SERT phosphorylation at maximal concentrations of 5-HT (1  $\mu$ M) with a median effective concentration (EC50) of 70 nM (Fig. 1B).

If the actions of 5-HT on SERT phosphorylation arise as a consequence of transport, then an intrinsic homeostatic loop might be present to link transporter expression to extracellular amine availability, and coincubation with SERT antagonists should block this effect. Indeed, the SERT-selective tricyclic antidepressant imipramine  $(1 \ \mu M)$  or the SSRIs paroxetine (1 µM) and citalopram (1 µM) blocked the ability of 5-HT to limit PKC-dependent SERT phosphorylation (Fig. 1C). This effect was selective for SERT antagonists, as neither the DAT inhibitor GBR-12909 nor the NET antagonist nisoxetine could affect the ability of 5-HT to blunt SERT phosphorylation (Fig. 1C). There are no known 5-HT receptor subtypes on HEK-293 cells, and to our knowledge, 5-HT does not induce acute changes in cyclic adenosine 5'-monophosphate, inositol trisphosphate, or intracellular Ca<sup>2+</sup> levels in these cells. Moreover, the ability of 5-HT to diminish PKCdependent SERT phosphorylation was not

Serotonin [5-hydroxytryptamine (5-HT)] is a platelet-stored vasoconstrictor that also acts as a transmitter in the nervous system to modulate a wide spectrum of behaviors (1). The actions of 5-HT are terminated by active transport (2). Whereas 5-HT actions are mediated by >15 different types of receptors, a single 5-HT transporter (SERT) is responsible for extracellular 5-HT clearance (3). SERT activity is blocked by cocaine and tricyclic antidepressants. Serotonin-selective reuptake inhibitors (SSRIs) like fluoxetine (Prozac) preferentially block SERTs and enhance serotonergic signaling in affective disorders (2, 4). The amphetamines are substrates for SERTs, as well as for dopamine (DA) and norepinephrine (NE) transporters (DATs and NETs, respectively) (5) and can trigger SERT-mediated release of 5-HT (5-8). Repeated administration of amphetamines

Department of Pharmacology and Center for Molecular Neuroscience, School of Medicine, Vanderbilt University, Nashville, TN 37232–6420, USA.

<sup>\*</sup>To whom correspondence should be addressed. Email: randy.blakely@mcmail.vanderbilt.edu