

Alarmed at such talk, Representative Vern Ehlers (R-MI) last week began rallying support behind civilian science. "We are concerned that funding for science may take a back seat" to housing, veterans' health care, and education programs, he wrote in a 20 March draft of a letter to colleagues that is aimed at the House leadership. "We ask you, in the strongest words possible, to assign a high priority to basic scientific research." There is evidence that the leadership is listening. Ehlers and Representative Rush Holt (D-NJ), both physicists, successfully lobbied last week for civilian R&D spending to get a tiny sliver more—some \$100 million—of the discretionary budget pie. Although appropriators are not bound by those numbers, "it shows there is political interest" in funding science, one House staffer says.

So despite the inevitable posturing and election-year rhetoric, many observers insist that R&D will ultimately prevail. Last year, for example, NASA's spending panel, after strong pressure from the White House, found a way to restore steep cuts in space science. "The outlook for federal R&D in 2001 is highly favorable," states a preview of the annual budget report published by the American Association for the Advancement of Science (which publishes *Science*). "It seems almost certain," the report predicts, that science and technology will achieve funding levels equal to or even surpassing those requested by Clinton. Even more certain, however, is that it will take a lot of political horse trading for that conventional wisdom to prevail.

—ANDREW LAWLER

## GAMMA RAY ASTRONOMY

### Aging NASA Satellite Headed for Fiery End

NASA has decided to euthanize an ailing but still functioning satellite rather than risk a slim chance that it could spin out of control and crash in a populated area. Last week NASA described plans for a late-spring maneuver that will cause the Compton Gamma Ray Observatory (CGRO) to enter the atmosphere and break apart across a swath of the Pacific Ocean.

The satellite is equipped with three gyroscopes, one of which failed late last year (*Science*, 21 January, p. 403). Although it needs only two to perform its scientific duties, a second failure—a possibility the Goddard Space Flight Center in Greenbelt, Maryland, puts at 10% in the next 3 years—would make the craft more difficult to con-


trol. "It was my decision [to put the craft down]," says Edward Weiler, NASA's associate administrator for space science, noting that his decision follows the original contingency plan. Although the craft is still productive, he says, NASA was faced with trying to calculate "how many papers are worth an increased risk to human life?"

Launched in 1991, the CGRO has already lasted almost twice its intended 5-year life-span. It mapped the sky's gamma rays, found the first gamma ray pulsars, and determined that gamma ray bursts originate beyond the Milky Way and are likely signs of the largest explosions since the big bang. Researchers have published more than 1000 papers based on CGRO data, including a paper last week linking some gamma ray sources to a new class of mysterious objects (see next story).

CGRO's scientific instruments will be turned off on 26 May. Five days later, a Goddard team will execute the first of four burns designed to shove the craft out of its orbit. The final two burns are planned for 3 June. Most of the 15,000-kilogram spacecraft will disintegrate and burn up when it hits the atmosphere, but 30 to 40 pieces will survive, ranging from bolt-sized to some possibly as heavy as 1000 kilograms. The detritus is expected to scatter in an area 25 kilometers wide and 1550 km long in the Pacific Ocean, about 4000 km southeast of Hawaii.

CGRO's demise will temporarily close a window onto gamma rays, which are blocked by the atmosphere. "Astronomers around the world will be quite disappointed," says CGRO project scientist Neil Gehrels. In 2003 NASA plans to launch a satellite, Swift, that will monitor gamma ray bursts, and in 2005 GLAST will focus on high-energy gamma ray phenomena. In the meantime, says Gehrels, "we have 62 days, 19 hours, and 42 minutes left" before the receivers are shut down. "We're going to make the best use of every one of those minutes."

—LAURA HELMUTH



**Ker-plunk.** Satellite debris should splash down on 3 June somewhere within a 1550-kilometer swath of the eastern Pacific Ocean.

## GAMMA RAY ASTRONOMY

### Sky Survey Finds Mysterious Strangers

Amid the diffuse bath of gamma rays coming from the galaxy, about 200 point sources—tiny gamma ray beacons—twinkle within the haze. For 2 decades, astronomers have been puzzling over what they are. Now astronomers at NASA's Goddard Space Flight Center in Greenbelt, Maryland, have doubled the mystery: They have discovered that these point sources come in two different varieties.

Neil Gehrels and his colleagues at Goddard analyzed the data from more than 4 years of observations with the EGRET telescope aboard the Compton Gamma Ray Observatory. Most of the sources lie along the galactic plane, but a few dozen lie in the middle latitudes, as much as 40 degrees above and below the galaxy's equator. In last week's issue of *Nature*, the team reported that the sources away from the galactic plane turned out to be surprisingly dim. "We didn't know what to make of the middle-latitude sources," says Gehrels. "The new population is much weaker than ones along the plane."

The mid-latitude sources appear to lie in the so-called Gould belt, a broad, expanding ring of stars, gas, and dust about 3000 light-years across. If so, they are our galactic neighbors. The Gould belt formed 30 million or 40 million years ago, probably when a series of stars exploded or some other powerful event disrupted the matter in our region of the galaxy. Clouds of gas, shocked outward, created a burst of new, massive stars. The bright sources in the galactic plane, in contrast, are likely to be tens of times farther away and thus, in absolute terms, vastly more powerful.

Isabelle Grenier, an astronomer at the University of Paris VII, finds the argument compelling. "What [Gehrels] has shown is that there is a clear difference between those out of the plane and those along the Milky Way," she says. "Now we're all convinced that there are two populations." Grenier also agrees that the weak sources are located in the Gould belt. "If you look at the distribution, you can really find that it follows the Gould belt," she says.

If Gehrels's team is correct, then the weak mid-latitude gamma ray sources and the intense equatorial sources may be entirely different types of objects. "The luminosity is so different between the two populations, there must be very different physical mechanisms," says Gehrels. Perhaps the mid-latitude sources are gamma ray pulsars, rapidly spinning geriatric stars that for some reason bathe us in gamma rays rather than the usual lower frequency radio waves. (If