

Determining the 3D Structure of HIV-1 Protease

In his News article "When Pharma merges, R&D is the dowry" (special issue on Drug Discovery, 17 Mar., p. 1952), Bruce Agnew writes that "Merck researchers were the first to determine the three-dimensional structure of the HIV-1 protease enzyme in 1989," and Roger Perlmutter is quoted as saying, "we published that structure so that everybody else could work on it, too." However, these statements do not accurately reflect the course of events.

The human immunodeficiency virus-type 1 (HIV-1) protease structure determined crystallographically by Merck researchers using recombinantly expressed HIV-1 protease was published in *Nature* in early 1989 (1). This structure was of the unliganded (empty active site) enzyme and was seriously flawed because the low resolution of the data led to an incorrect tracing of the polypeptide chain at the dimer interface. In any event, only the coordinates of the carbon alpha atoms of the main chain were deposited with the Protein Data Bank (PDB). Such limited data for the unliganded enzyme were of little, if any, use to researchers undertaking structure-based drug design.

The first complete and correct structure of the HIV-1 protease was determined crystallographically at the National Cancer Institute (NCI) using enzyme prepared by total chemical synthesis in Kent's laboratory at the California Institute of Technology, and the structure was published in August 1989 (2). The more important structure of an HIV-1 protease-ligand complex was also determined at NCI, again using enzyme prepared by total chemical synthesis in Kent's laboratory at Caltech with a substrate-derived inhibitor prepared by Marshall's laboratory at Washington University at St. Louis. That structure was published in December 1989 (3). These structures of the synthetic enzyme were of high resolution and of good quality, providing an appropriate target for structure-based drug design. The full coordinates for both structures were immediately deposited in the PDB and were made freely available to researchers.

Stephen Kent

Gryphon Sciences, 250 East Grand Avenue, South San Francisco, CA 94080, USA. E-mail: skent@gryphonsci.com

Garland R. Marshall

Institute for Biomedical Computing, Center for Molecular Design, Washington University, St. Louis, MO 63110-1012, USA. E-mail: garland@gpc.wustl.edu

Alexander Wlodawer

NCI-Frederick Cancer Research Facility, Frederick, MD 21702, USA. E-mail: wlodawer@ncifcrf.gov

References

1. M. A. Navia *et al.*, *Nature* **337**, 615 (1989).
2. A. Wlodawer *et al.*, *Science* **245**, 616 (1989).
3. M. Miller *et al.*, *Science* **246**, 1149 (1989).

Response

As Kent, Marshall, and Wlodawer make plain, numerous groups (including their own) made contributions to the determination of the structure of the HIV-1 protease. There appear to be no serious issues of contention between us. Merck Research Laboratories made public a structural analysis of the HIV-1 protease and deposited the data in the PDB in early 1989. The structure was not "seriously flawed," although we readily acknowledge that it was incomplete. It provided the best, and at the time the only, representation of the structure of the HIV-1 protease. Resolution of the alpha chain backbone was a fundamental first step.

Crystallographic analysis is typically iterative, and subsequent work by Kent, Marshall, and Wlodawer clearly provided substantive and more detailed information. The important point, as I indicated in Agnew's article, is that the initial publication of structural data by Navia *et al.* accelerated the development of protease inhibitors by several pharmaceutical companies, to the general benefit of patients suffering from HIV infection.

Roger M. Perlmutter

Executive Vice President, Worldwide Basic Research and Preclinical Development, Merck Research Laboratories, Whitehouse Station, NJ 08889, USA

Annotation of the Human Genome

The News article "Are sequencers ready to 'annotate' the human genome?" by Elizabeth Pennisi (special issue on the *Drosophila* Genome, 24 Mar., p. 2183) is especially timely and provocative. Pennisi mentions two ideas: a small group gathering at a centralized annotation jamboree, or a distributed, Web-based system that would allow anyone to contribute annotations with a "smart browser" that would merge all efforts. I favor the essence of the second proposal because it provides a more democratic and more "biological" approach to an all-important problem.

There is, however, a third approach for annotating the human genome (providing at least the putative start, stop, and structure of each gene) that is, in a sense, already extant: extend the capabilities of the biological science literature. The current journal system is decentralized, yet most research articles adhere to common standards that make them ideal for annotation: (i) Each article associates a bit of annotation with a distinct time and place and with specific, responsible parties. (ii) Attentive scholarly referencing and footnoting provide a way to connect bits of annotation and allow for

continuous "updates." (iii) Peer review and editing provide a proven quality-control mechanism. (iv) Publication is an established indicator of scientific productivity; consequently, scientists already have an incentive to provide the information, whereas database submissions are often regarded as a chore.

The main drawback of current journal article formats is that they are not very "computer-parseable," or suitable for bulk annotation of thousands of genes. However, by adding sections of highly structured text to each article (that is, extended keywords and using a controlled vocabulary) and linking subparts of an article to relevant database identifiers, one can envision how a "literature annotation standard" could readily be interpreted by computers. Furthermore, if an article could be linked to a large "supplementary materials" data file with simple annotations for many genes (for example, lists of all the membrane proteins in the *Caenorhabditis elegans* genome), one would have a mechanism for bulk annotation. Further standardization could be achieved if the article described defined ways in which the data file might be updated over time and if the supplementary materials were refereed and evaluated with the text of the article.

Mark Gerstein

Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT 06520-8114, USA. E-mail: mark.gerstein@yale.edu

Window on the Early Earth

Carl Zimmer's News Focus article "Ancient continent opens window on the early Earth" (17 Dec., p. 2254) highlights interpretations of a very old fragment of Canadian continental crust by a team headed by Wouter Bleeker and Richard Stern. This team has substantially advanced understanding of the early geologic evolution of the Slave Province, but Zimmer attributes solely to these scientists the model of an ancient protocontinent overlain by a shallow-water sedimentary sequence in the western Slave Province, and collision of this protocontinent with a younger arc terrane 2.7 to 2.6 billion years ago. Virtually the same model was conceived 15 years before and published in reputable journals, which is nowhere mentioned in the article.

The model attributed to Bleeker and Stern stems from years of work, including more than 9 months in the field, mapping the distribution of and determining basic geological relationships between ancient basement rocks and surrounding units (1). The boundaries of this old crustal fragment were defined on the basis of field relationships and a limited number of uranium-lead (U-Pb) dates, and the name "Anton terrane" was proposed along with its interpretation as an ancient continent (2). The boundaries of the old continent have only been slightly modified on the map pre-